

An Educational Program Using A Bladder Model To Improve Diagnostic Cystourethroscopy Skills

C Kilpatrick, L Chohan, G Siddiqui, E Reichman, F Orejuela

Citation

C Kilpatrick, L Chohan, G Siddiqui, E Reichman, F Orejuela. *An Educational Program Using A Bladder Model To Improve Diagnostic Cystourethroscopy Skills*. The Internet Journal of Medical Simulation. 2008 Volume 3 Number 1.

Abstract

Objective: To determine if an educational program with the use of a bladder model, constructed out of a rubber ball and balloon, improved surgical skills requisite in the performance of diagnostic cystourethroscopy. **Methods:** We created a bladder model out of a rubber ball and balloon. We invited all Obstetrics and Gynecology residents at The University of Texas Houston to participate in the study. We assigned participants to receive the intervention, interactive training and practice on a bladder model, or the usual education and teaching. All participants had baseline knowledge assessed using an Objective Structured Assessment of Technical Skills (OSATs) in August and November of 2008. The OSATs consisted of a Global Ratings Scale with pass/fail grade, task specific checklist of 11 items, written questionnaire, and time required to complete a diagnostic cystourethroscopy on the bladder model. We assessed post intervention knowledge and skills in a similar fashion in February and May of 2009. Statistical analyses were performed using the SAS statistical package. A p value of $<.05$ was considered statistically significant.

Results: Forty one of 45 (91.1%) residents agreed to participate in the study. Twenty-five residents (60.9%) had complete information with pre and post intervention measurements. A trend in improved performance was seen in those that received the intervention versus control although none reached statistical significance. Inter-rater reliability was greatest for the pass/fail portion of the global rating scale.

Conclusion: This model may be useful in training obstetric and gynecology residents in diagnostic cystourethroscopic procedures.

INTRODUCTION

Injury to the urinary tract during gynecologic surgery is not uncommon. In some series the ureteral injury rate is as high as 26.8/1000, with bladder injury at 29.1/1000¹. Diagnostic cystourethroscopy at the time of difficult gynecologic cases or cases where ureteral integrity is questioned has been shown to reliably identify ureteral injury^{2,3,4}. As the population ages it is likely that the number of cases for stress urinary incontinence, and pelvic organ prolapse will increase⁵. Currently, training in diagnostic cystourethroscopy of gynecology residents varies across the nation.

We designed an Objective Structured Assessment of Technical Skills (OSATS) using a bladder model, constructed out of a rubber ball and balloon, for teaching requisite skills necessary in the performance of diagnostic cystourethroscopy. The purpose of this study was to determine whether an educational program, assessed using the OSATS, improved the surgical skills necessary for the

performance of diagnostic cystourethroscopy. Our hypothesis was that the use of this model and education provided in association with the use of this model would improve surgical skills compared to the group that did not receive training using the model.

MATERIALS AND METHODS

We invited all residents of both obstetrics and gynecology training programs at The University of Texas Health Science Center at Houston (UTHSCH) to participate in the study. We fashioned a low cost bladder model out of a rubber ball and balloon. The rubber ball was approximately 4 cms in diameter, and in the center of the ball we created a 5mm urethra with the drill. The balloon fit snugly over the rubber ball, as depicted in the figure, and we attached the balloon and ball to the simulation table with a Velcro strap. We marked two areas in the balloon representing the ureteral orifi based on published anatomic criteria⁶, and also an area marking the dome of the bladder.

We created a written questionnaire and 11 point task specific checklist (Figure 1) in regard to diagnostic cystourethroscopy. We created the checklist based on essential steps necessary to perform a diagnostic cystourethroscopy and those highlighted in a contemporary text⁷. The written questionnaire assessed the learners' knowledge in regard to diagnostic cystourethroscopy.

Figure 1

Figure 1: Task Specific Checklist

Procedure Checklist for Bladder Training, OSATS-C		
Item	Performed	Not Performed
Recognizes and assembles the correct parts of the cystoscope		
Connects the cystoscope to the light source		
Connects the fluid source to the cystoscope		
Has the assistant hang the fluid source no greater than 60 cms above the bladder model		
Connects the camera to the eyepiece of the telescope		
Performs a white balance		
Ensures the tube is flushed		
Ensures the correct ports are closed on the cystoscope, and water leakage does not occur		
Applies lubricant to the tip of the sheath prior to insertion		
Instills 100-300 ccs of fluid		
Correctly identifies bladder markers on survey of bladder model in correct anatomic position		

We divided the participants into an intervention group, which consisted of a one hour interactive presentation followed by two hours of practice using the bladder model, and control group, which received the usual education and teaching. The usual education and teaching was that performed in the operating room during diagnostic cystourethroscopy. This training likely varied based on the attending physician supervising the case but was performed on all cases in which there was the possibility of ureteral injury. Resident physicians were the primary surgeons on all cases performed in the operating room. The residents were divided by the first author of the study based on post graduate year of training. Prior to dividing the residents into groups the residency directors were polled to determine the number of diagnostic cystoscopic procedures performed, and no statistical differences were noted between participants.

We obtained informed consent of all participants and then performed baseline testing of all participants. We assessed knowledge with the written questionnaire. We then timed and evaluated the participants while performing a diagnostic cystourethroscopy on the bladder model using the global rating scale with pass/fail grade⁸ (Figure 2), and task specific checklist. We allowed the participants ten minutes to complete the task on the bladder model. The global rating scale consists of a pass/fail grade and a five point likert scale with seven categories: respect for tissue, time and motion, instrument handling, knowledge of instruments, flow of

operation, use of assistants, and knowledge of specific procedure. The global rating scale is a validated scale to measure resident surgical performance. The task specific checklist, which along with the global rating scale comprised the OSATS, was not validated.

Figure 2

Figure 2: Global Rating Scale

GLOBAL RATING SCALE OF OPERATIVE PERFORMANCE					
Please circle the number corresponding to the candidate's performance in each category, irrespective of training level.					
Respect for Tissue:					
1	2	3	4	5	
Frequently used unnecessary force on tissue or caused damage by inappropriate use of instruments		Cautious handling of tissue but occasionally caused inadvertent damage		Consistently handled tissues appropriately with minimal damage	
Time and Motion:					
1	2	3	4	5	
Many unnecessary motions		Efficient movements but some unnecessary pauses		Clear economy of movement and maximum efficiency	
Instrument Handling:					
1	2	3	4	5	
Repetitively makes mistakes or performs routine with instruments by inappropriate use of instruments		Consistent use of instruments but occasionally appeared stiff or awkward		Fluid moves with instruments and an air-water motion	
Knowledge of Instruments:					
1	2	3	4	5	
Frequently asked for wrong instrument or used inappropriate instrument		Knew names of most instruments and used appropriate instrument		Thoroughly familiar with the instruments and their names	
Flow of Operation:					
1	2	3	4	5	
Frequently stopped operating and moved around in and out of room		Uninterrupted smooth forward planning with reasonable progression of thoughts		Uninterrupted planned course of operation with efficiency from first one motion to the next	
Use of Assistants:					
1	2	3	4	5	
Consistently ignored assistants, prodded or talked to one assistant		Appropriate use of assistant most of the time		Strategically used assistants to the best advantage at all times	
Knowledge of Specific Procedures:					
1	2	3	4	5	
Deficient knowledge, missed specific, instructions at most steps		Knew all important steps of operation		Uninterrupted familiarity with all aspects of operation	
OVERALL ON THIS TASK, SHOULD THE CANDIDATE:					
				FAIL	PASS

Each participant had two evaluators, and four evaluators total participated in the study. Prior to participating in the study the evaluators met to discuss the evaluation tools so as to be uniform in evaluation interpretation. We deemed failure of the task as not being able to complete the task within ten minutes. All of the evaluators in the study were blind to group allocation except the first author.

We performed the baseline testing in August and November of 2008. We provided the intervention in December 2008, or usual teaching, followed by post intervention testing. We performed post intervention testing in the same manner as baseline testing in February and May of 2009. The participants were not blind to group assignment.

We obtained data in four categories at each evaluation: the written questionnaire, task specific checklist, global rating scale ranging from one to five, and the time to complete the task with a maximum of 600 seconds. We recorded the pass/fail portion of the global rating scale.

Statistical analyses were performed using the SAS statistical package. Inter-rater reliability was measured using the Pearson correlation coefficient test for continuous variables. Chi square and Fisher's exact test when appropriate were

performed to compare nominal variables. The Wilcoxon sum test was used to compare the continuous variables between cohorts, and the Wilcoxon matched-pairs signed rank test within cohorts.


We obtained IRB approval for the study from UTHSCH and report the data using the CONSORT guidelines for randomized controlled trials⁹.

RESULTS

Forty one out of 45 (91.1%) residents agreed to participate in the study. Twenty-one residents were assigned to the intervention group and 20 to the usual teaching. Twelve participants in the intervention group and 13 in the control group had complete information with pre and post intervention measurements. Eight of the 12 residents assigned to the intervention group received the intervention (66.7%). Figure 3 is a flowchart demonstrating participant allocation.

Figure 3

Figure 3: Flow chart



The price of the rubber balls and balloons were negligible, and the drill and drill bits used to create a 'urethra' in the rubber ball about \$40.

There were no statistically significant differences in postgraduate year of training between the residents in the intervention group and the control ($p=0.93$).

Participants were analyzed in the group to which they were assigned. We discovered no statistically significant differences in pre test or post test scores between those who were assigned to the intervention and those who were not. Table 1 lists the scores on the task specific checklist, global rating scale, questionnaire, and time to complete task at baseline (pretest) and after intervention (post test). We also

found no significant differences in pass/fail grade at baseline or after intervention between the control and intervention group.

Figure 4

Table 1.

Table 1: Pre and post intervention test results between cohorts

	Pretest			Posttest		
	Score±SD			Score±SD		
Evaluation	Control	Intervention	P value	Control	Intervention	P value
Tool	Group	Group		Group	Group	
	(N=13)	(N=12)		(N=13)	(N=12)	
Task Specific Checklist	57.1±22.9	45.1±21.9	0.20	76.6±18.0	70.1±15.1	0.16
Global Rating Scale	3.1±1.1	2.3±1.0	0.06	3.77±0.8	3.95±0.8	0.62
Time to Complete (sec)	441±145	506.4±105.9	0.17	356±117.3	344±154.5	0.58
Questionnaire	50±14	46±9	0.28	49±11	55±9	0.08

Table 2 outlines the results within cohorts and noted a statistically significant improvement from the pretest to post test within the intervention group in regard to the task specific checklist (.004), global rating scale (.004), time to complete in seconds (.006), and questionnaire (.01). We noted a statistical significant difference in the control group in the task specific checklist (.03) and a trend toward improved performance in the other tasks but this did not reach statistical significance.

Figure 5

Table 2.

Table 2: Pre and post intervention test results within cohorts

	Pretest	Post test	P value	Pretest	Post test	P value
	Score±SD	Score±SD		Score ±SD	Score±SD	
Evaluation	Control	Control		Intervention	Intervention	
Tool	Group	Group		Group	Group	
	(N=13)	(N=13)		(N=12)	(N=12)	
Task Specific Checklist	57.1±22.9	76.6±18.0	0.03	45.1±21.9	70.1±15.1	0.004
Global Rating Scale	3.1±1.1	3.77±0.8	0.08	2.3±1.0	3.95±0.8	0.004
Time to Complete (sec)	441±145	356±117.3	0.16	506.4±105.9	344±154.5	0.006
Questionnaire	50±14	49±11	0.8	46±9	55±9	0.01

We also analyzed the difference in improvement (pretest to post test scores) between the intervention and control group for the task specific checklist (mean 0.25 vs 0.19, p=0.549), global rating scale (mean 1.55 vs 0.63, p=0.057), and reduction in time to complete the task (mean 162 secs vs 85 secs, p=0.276). Although there was a trend toward improved performance in all areas measured there was no statistical significance achieved. The global rating scale was close to statistical significance.

Inter-observer reliability on pre/post test results for the task specific checklist (0.78); global rating scales (0.87) and pass/fail grade (1.0) were high.

DISCUSSION

Surgical education has changed over the last two decades. As early as the 1970's, surgeons had developed task specific checklists in order to evaluate operative skills¹⁰. With the work hour restrictions, emphasis from the ACGME to implement standardized teaching for surgical skills, and a move toward more competency-based surgical education, simulation learning is a necessary tool in resident education. Recent literature in regard to obstetric and gynecologic surgical simulation training would support its efficacy^{11,12,13}. Also, given the likelihood that the number of procedures that require diagnostic cystourethroscopy to ensure ureteral/bladder integrity will increase, we felt an educational program to teach this skill was necessary.

There are a number of strengths to our study including those associated with the use of a model versus observation of real time surgery. Our model is inexpensive. This is in contrast to a similar study used to evaluate diagnostic cystourethroscopy skills among residents where the costs associated with the model neared \$4,000¹⁴. We had no operating room time costs. We avoided patient safety issues, and the model was standardized; the trainee was able to complete the task without concern for any patient related factor that would require an attending physician to step in and complete the task. Also, the time involved in the interactive session was not prohibitive, an hour for didactics and a couple of hours for practice.

Much like most of the gynecology simulation literature, we did not compare performance on the model in relation to live operating room performance. We cannot say whether superior performance on the bladder model leads to improved technical performance in the operating room. We believe the statistically significant increase observed in the control group in regard to performance on the task specific

checklist and the trend toward improved overall performance can be explained by a number of factors; recent attention in the current literature to the importance of diagnostic cystourethroscopy, increase in the number of diagnostic cystourethroscopies performed at our institution during the study period, and involvement in the study itself may have motivated control group participants to better their performance (Hawthorne effect).

Given the resident rotation schedules, participants were not truly randomized. However, we doubt this was a source of bias as the numbers of senior and junior residents in each arm were similar, as evidenced by no statistical differences in scores at baseline testing. Participation was initially strong but waned due to a conflict in the laboratory space which changed the post intervention testing dates. We did not monitor the number of cystoscopes performed by study participants, possibly affecting post intervention scores. Given the low numbers, any significant conclusions proved difficult. Also, the evaluators knew the participants possibly introducing bias into the scores.

We observed those that were randomized to the intervention had statistically significant improvements in all evaluation categories, while the control group only showed an improvement in the task specific checklist. We believe that training using the balloon model and didactics that we have outlined improves the skills necessary for diagnostic cystourethroscopy.

We believe this economical model is useful in training residents in diagnostic cystourethroscopic procedures. A larger sample size is required to show if training improves skill level. We have incorporated this OSATs with bladder model into our residency curriculum.

References

1. Gilmour DT, Dwyer PL, Carey MP. Lower urinary tract injury during gynecologic surgery and its detection by intraoperative cystoscopy. *Obstet Gynecol.* 1999; 94(5):883-9.
2. Kwon CH, Goldberg RP, Koduri S, et al. The use of intraoperative cystoscopy in major vaginal and urogynecologic surgeries. *Am J Obstet Gynecol.* 2002; 187(6):1466-71.
3. Jabs CF, Drutz HP. The role of intraoperative cystoscopy in prolapse and incontinence surgery. *Am J Obstet Gynecol.* 2001; 185(6):1368-71.
4. Ibeanu OA, Chesson RR, Echols KT, et al. Urinary Tract Injury During Hysterectomy Based on Universal Cystoscopy. *Obstet Gynecol.* 2009; 113(1): 6-10.
5. Lubner KM, Boero S, Choe JY. The demographics of pelvic floor disorders: current observations and future projections. *Am J Obstet Gynecol.* 2001; 184(7):1496-501.
6. Rahn DD, Bleich AT, Wai CY et al. Anatomic relationships of the distal third of the pelvic ureter, trigone, and urethra in unembalmed female cadavers. *Am J Obstet Gynecol.* 2007 Dec;197(6):668.e1-4.
7. Dwyer PD: Atlas of Urogynecological Endoscopy, 1st edition. London, Informa Health Care, 2007, 189 pp.
8. Reznick R, Regehr G, MacRae H, et al. Testing technical skill via an innovative "bench station" examination. *Am J Surg.* 1997 Mar;173(3):226-30.
9. Altman DG, Schulz KF, Moher D, et al. The Revised CONSORT Statement for Reporting Randomized Trials: Explanation and Elaboration. *Annals of Internal Medicine.* 2001; 134(8): 663-694.
10. Kopta JA. An approach to the evaluation of operative skills. *Surgery.* 1971; 70(2):297-303.
11. VanBlaricom AL, Goff BA, Chinn M, et al. A new curriculum for hysteroscopy training as demonstrated by an objective structured assessment of technical skills (OSATS). *Am J Obstet Gynecol.* 2005; 193(5):1856-65.
12. Lentz GM, Mandel LS, Goff BA. A six-year study of surgical teaching and skills evaluation for obstetric/gynecologic residents in porcine and inanimate surgical models. *Am J Obstet Gynecol.* 2005 Dec;193(6):2056-61.
13. Siddighi S, Kleeman SD, Baggish MS, et al. Effects of an educational workshop on performance of fourth-degree perineal laceration repair. *Obstet Gynecol.* 2007; 109(2):289-94.
14. Fialkow M, Mandel L, VanBlaricom A, et al. A curriculum for Burch colposuspension and diagnostic cystoscopy evaluated by an objective structured assessment of technical skills. *Am J Obstet Gynecol.* 2007; 197(5):1-6.

Author Information

Charlie C. Kilpatrick, MD

Department of Obstetrics and Gynecology and Emergency Medicine, University of Texas Health Science Center at Houston

Lubna Chohan, MD

Department of Obstetrics and Gynecology and Emergency Medicine, University of Texas Health Science Center at Houston

Gazala Siddiqui, MD

Department of Obstetrics and Gynecology and Emergency Medicine, University of Texas Health Science Center at Houston

Eric F. Reichman, PhD, MD

Department of Obstetrics and Gynecology and Emergency Medicine, University of Texas Health Science Center at Houston

Francisco J. Orejuela, MD

Department of Obstetrics and Gynecology and Emergency Medicine, University of Texas Health Science Center at Houston