

Total Salivary Glutathione Levels: Periodontitis in Smoker and non-Smoker

B Rai, R Jain, S Anand, S Kharb

Citation

B Rai, R Jain, S Anand, S Kharb. *Total Salivary Glutathione Levels: Periodontitis in Smoker and non-Smoker*. The Internet Journal of Laboratory Medicine. 2008 Volume 3 Number 2.

Abstract

Objective: To determine the association between Total glutathione levels, in saliva and presence of periodontitis. **Subjects and Methods:** Forty subject samples, as healthy control (n=16), periodontitis in smokers (n=11) and periodontitis in non-smokers (n=13) were recruited for the study. The total glutathione levels were analyzed in the saliva of smoking and non-smoking patients of periodontitis before and after scaling and root planning and healthy control. **Results and Conclusions:** Total glutathione of saliva were statistically significantly higher in subject of periodontitis in smoker as compared to non-smoker and salivary total glutathione concentrations were significant reduced following scaling and root planning in smoker as well as in nonsmoker. Total glutathione of saliva was affected by periodontal disease. Thus smoking and periodontitis compromised the antioxidant capacity of saliva in systemically healthy patients.

INTRODUCTION

The human inflammatory periodontal diseases are amongst the most common chronic diseases to affect adults. Antioxidants are present in all body fluids and tissues and protecting against endogenously formed free radicals. Antioxidant (AOX) enzymes such as superoxide dimutase and glutathione peroxidase provide protection within cells whilst low molecular weight scavenging antioxidants and AOX vitamins are present in extra cellular fluid. In addition, dietary driven components such as uric acid non protein thiols and glutathione also act as antioxidants¹. Total antioxidant activity has been reported to be reduced in saliva of patients with periodontitis relative to that in non-periodontitis subjects². Diagnosis of periodontal disease has been primarily based upon clinical and radiographic measures of periodontal tissue destruction. These parameters (clinical examination and radiographically) provide a measure of past destruction and are of limited use in early diagnosis³. Smokers are more susceptible to advanced and aggressive forms of periodontal disease than non-smokers⁴. It has been reported that smokers tend to respond less favourably to periodontal treatment⁵. Reactive oxygen species play an important role in cell signaling and metabolic processes and contribute to pathogenic processes in a variety of inflammatory disorders. All organisms possess a range of enzymatic and non-enzymatic antioxidant systems

to protect them against harmful oxidative reactions⁶⁷. Under certain conditions, an increase in oxidants and a decrease in antioxidants cannot be prevented, and the oxidant or antioxidant balance shifts towards the oxidative state. There is increasing evidence that oxidative stress is an important contributing factor in the pathogenesis of periodontal disease⁷⁸⁹¹⁰. Antioxidant molecules are present in all body fluids and tissues. Saliva possesses a variety of defense mechanisms responsible for the protection of oral cavity from oxidative attacks, including uric acid, vitamin C, glutathione, and others. Together with uric acid and albumin, ascorbic acid is among the major antioxidants in saliva¹²³⁴⁵⁶⁷⁸⁹¹⁰.

Hence, the present study was planned to determine the relationship between total glutathione levels in saliva in cases of healthy periodontium, periodontitis in smokers and non-smokers, and effect of scaling and root planning on salivary glutathione levels.

MATERIALS AND METHODS

The patients of periodontitis in smokers (11), periodontitis in non-smokers (13) and healthy periodontium (16), without any systemic disease, aged 30-50 years attending Jain Diagnostic clinic, New Delhi, India were selected. The clinical examination, invaluable measurement of both probing depths and bleeding on probing, performed by two

experienced examiners who measured the same clinical parameters throughout the study. During the examination and one week after scaling and root planing, paraffin wax stimulated whole saliva was collected, and samples were stored at -20 ° C until analyzed. Saliva were centrifuged at 8000g for 9 minutes, and levels of total salivary glutathione was determined by using a kinetic enzymatic recycling assay, according to the manufacturers instructions (Oxford Biomedical research, Oxford, MI, USA) ¹¹.

Differences in Total glutathione of saliva levels between groups at baseline were analyzed by the mann-whitney's U-test. Differences in Total glutathione of saliva samples, probing depths and bleeding on probing were analyzed using a student's t-test. All statistical analyses were performed using SPSS.

OBSERVATIONS AND RESULTS

The mean salivary total glutathione level in subjects with a clinically healthy periodontium was 3.6 ± 1.8 micromole ($p < 0.05$). In periodontitis in smokers and non-smokers involved teeth of hopeless prognosis, the mean total salivary glutathione levels were 5.7 ± 1.8 micromole ($p < 0.05$), 4.8 ± 1.3 micromole ($p < 0.05$) respectively. The mean probing depth and bleeding on probing were 4.82 ± 0.12 mm & 64.2 ± 0.5 % & 3.94 ± 0.14 mm & 54.3 ± 0.6 % & 1.92 ± 0.14 mm & 25.2 ± 0.4 % in periodontitis in smokers & non-smokers and healthy periodontium. Significant reductions in clinical recordings were obtained in smoker and non-smoker patients following periodontal treatment. Salivary total glutathione conc. were reduced following therapy in smoker as well as in non-smoker (Table-1, $P < 0.05$)

Figure 1

Table I Total Glutathione (micromole) , probing depth and bleeding in probing in periodontitis in smokers, periodontitis in non-smokers and healthy periodontium before and after treatment.

Parameters	Mean + SD		
	Periodontitis in non-smokers	Periodontitis in smokers	Healthy periodontium
Total salivary glutathione content (before treatment)	4.8 ± 1.3^a	$5.7 \pm 1.8^{*,a}$	$3.6 \pm 1.8^*$
Total salivary glutathione content (After one week of scaling and root planning)	3.9 ± 1.6^a	$5.1 \pm 1.5^{*,a}$	$3.6 \pm 1.8^*$
Probing depth (in mm) before treatment	3.94 ± 0.14^a	$4.82 \pm 0.12^{*,a}$	$1.92 \pm 0.14^*$
Probing depth (in mm) after treatment	3.21 ± 0.14^a	$4.41 \pm 0.16^{*,a}$	$1.91 \pm 0.15^*$
Bleeding on probing (in %) before treatment	54.3 ± 0.6^a	$64.2 \pm 0.5^{*,a}$	$25.2 \pm 0.4^*$
Bleeding on probing (in %) after treatment	34.6 ± 0.7^a	$57.4 \pm 0.6^{*,a}$	$25.3 \pm 0.5^*$

^a $p < 0.05$, as compared to healthy control.

^{*} $p < 0.05$, as compared to periodontitis in non-smokers.

DISCUSSION

Stimulated saliva contains a lower concentration of antioxidants but when flow rates are taken into account, antioxidant capacity is higher than in unstimulated saliva. Stimulated saliva has been used in the analysis of antioxidant ¹⁰. PMN has been observed to be in greater concentration at sites of gingival inflammation ¹¹. They postulated that O_2^- produced by PMN as part of the host immune response could cause host tissue oxidative damage if it were not matched by an increase in antioxidant concentration ¹². Number of studies have evaluated the use of various host derived factors in saliva for diagnosis of periodontal disease. ^{10,11,12} In the present study, higher level of total glutathione contents were observed in periodontitis patients and level were still higher in smokers as compared to non-smokers (table 1, $p < 0.05$). Our results are compared with those reported in the literature. ¹¹ It may be due to one potential mechanism is through tissue damage mediated by oxidative species originating from tobacco smoke and

tobacco induced inflammation ,in addition to the direct cigarette smoke –mediated depletion of antioxidants.Significant reductions in clinical recordings were obtained in smoker and non-smoker patients following periodontal treatment.Salivary total glutathione conc. were significantly reduced following therapy in smoker as well as in non-smoker (Table-1,P<0.05) while in previous study only no statistically reduction of glutathione levels in periodontal disease patients ¹¹ .The acute influence of smoking a single cigarette on concentrations of glutathione ,uric acid, and total antioxidant activity measured in saliva has been addressed previously ¹³ .

It has been reported that the possible effects of smoking and gingival inflammation on salivary antioxidants in gingivitis patients. They reported that no statistically significant difference was found in any of antioxidant indices between any of the groups ¹⁴ . Few authors determined that GCF antioxidant concentration was significantly lower in periodontitis subjects compared to healthy controls. Thus, periodontal disease has been suggested to be associated with reduced salivary antioxidant status and increased oxidative damage within oral cavity ^{11,12,13,14,15} . Since, saliva can be easily collected, measurement of Total salivary antioxidant capacity levels may prove to useful in identifying patients at risk of tooth loss. Moreover, a salivary analysis for periodontal diagnosis may prove a cost effective method for screening large populations. Further, studies involving larger group sizes and analysis of GCF total glutathione are required to address these questions.

References

1. Halliwell B. Reactive oxygen species in living systems : source, biochemistry, and role in human disease. American Journal of Medicine 1991; 91 : 14-22.
2. RaiB,Jain R,Anand SC,KharbS.8-hydroxydeoxyguanosine levels:periodontitis in smoker and nonsmoker:A pilot study.J Pak Dent Assoc.2006;15:89-90.
3. RaiB, Anand SC,Kharb S.Effect of scaling and root planning on salivary 8-hydroxydeoxyguanosine levels:periodontitis. J Pak Dent Assoc.2007;16(4):189-191.
4. Calsina G,Ramon JM,Echeverria JJ.Effects of smoking on periodontal tissues.Journal of Clinical Periodontology.2002;29:771-76.
5. Renvert S,Dahlen G,Wikstrom M.The clinical and microbiological effects of non-surgical periodontal therapy in smokers and non-smokers.Journal of Clinical Periodontology.1998;25:153-57.
6. Waddington RJ, Moseley R, Emberg G. Reactive oxygen species : a potential role in the pathogenesis of periodontol disease. Oral Dis 2000; 6 : 138-151.
7. Chapple IL. Reactive oxygen species and antioxidants in inflammatory diseases. J Clin Periodontol 1997; 24 : 287-96.
8. Becker W, Berg L, Becker BE. The long term evaluation of periodontal treatment and maintenance in 95 patients. Int J Periodontics Restorative Dent 1984; 4 : 54-71.
9. Moore S, Calder KAC, Millar NJ, Rice Evans CA. Antioxidant activity of saliva and periodontal disease. free Radical Research 1994; 21: 417-425.
10. Takane M, Sagano N, Ezawa T, Lichiyama T, Ito K. A Marker of oxidative stress in saliva : association with periodontally involved teeth of a hopeless prognosis. J Oral Science 2005; 47 : 53-57.
11. Buduneli N,Kardesler L,Isik H,Willis CS,Hawkins SI,Kinane DF,Scott DA.Effects of smoking and gingival inflammation on salivary antioxidant capacity.J clin Periodontol.2006;33:159-64.
12. Ohshima M, Fujikawa K, Akutagawa H, Kato T, Ito K, Otsuka K. Hepatocyte growth factor in saliva : a possible marker for periodontal disease status. J Oral Sci 2002; 44 : 35-39.
13. Brock GR, Butterworth CJ, Mathews JB, Chapple IL. Local and systemic total antioxidant capacity in periodontitis and health. J Clin Periodontol 2004; 31 : 515-521.
14. Wilton JM, Curtis MA, Gillett IR, Griffiths GS, Maiden MF, Sterne JA, Wilson DT, Johanson NW. Detection of high risk groups and individuals for periodontal diseases : laboratory markers from analysis of saliva. J Clin Periodontol 1989; 16 : 475-483.
15. Aurer A, Aurer-Kozelj J, Stavlijenic Rukavina A, Kalenic S, Ivic-Kardum M, Haban V. Inflammatory mediators in saliva of patients with rapidly progressive periodontitis during war stress induced incidence increase. Coll Antropol 1999; 23 : 117-124.

Author Information

Balwant Rai

Editor In Chief Internet Journal Of Dental Science, P.D.M.Dental College And Research Institute,Bahadugarh , Haryana (INDIA)

Rajnish Jain, (M.D.S.) Endodontics

Kalka Dental college (INDIA)

SC Anand, (M.D.S.) Oral & Maxillofacial Surgery and Orthodontics

P.D.M.Dental College And Research Institute,Bahadugarh , Haryana (INDIA)

Simmi Kharb, (M.D.) Bio-Chemistry

Pt. Bhagwat Dayal Sharma, Post Graduate Institute of Medical Science, Rohtak, Haryana (INDIA)