Identification and Characterization of a Putative Chemotaxis Protein, CheY, from the Oral Pathogen *Campylobacter rectus*

M J LaGier, I Bilokopytov, B Cockerill, D S Threadgill

Citation

M J LaGier, I Bilokopytov, B Cockerill, D S Threadgill. *Identification and Characterization of a Putative Chemotaxis Protein, CheY, from the Oral Pathogen Campylobacter rectus*. The Internet Journal of Microbiology. 2014 Volume 12 Number 1.

DOI: [10.5580/IJMB.21300](https://doi.org/10.5580/IJMB.21300)

Abstract

Campylobacter rectus is an understudied oral bacterium that contributes to periodontitis. Processes that contribute to the disease-causing capabilities of pathogens, such as chemotaxis, are largely unknown in *C. rectus*. The aim of this study was to better understand *C. rectus* chemotaxis, by examining the *C. rectus* genome for the presence of a *cheY* gene. CheY proteins play a part in chemotaxis by acting as two-component response regulators. Significantly, CheY proteins from several pathogens, including the related species *Campylobacter jejuni*, have been shown to contribute to bacterial virulence. Degenerate PCR, RT-PCR, sequence analyses, and structural modeling showed that *C. rectus* encodes a gene (*Cr-CheY*) which shares significant homology with previously characterized CheY proteins. Functional studies of a recombinant form of the protein supports a likely role of *Cr-CheY* in *C. rectus* chemotaxis. *Cr-CheY* is the first CheY characterized from the oral campylobacters.

INTRODUCTION

Periodontitis is an infectious disease of the supportive tissue of the teeth (Philstrom et al., 2005). It infects 10% of the population making it a relatively common infectious disease (Teng et al., 2002). Approximately 13% of patients develop severe forms of periodontitis, which, may result in tooth loss and systemic complications including an increased risk of pregnancy complication (Philstrom et al., 2005; Teng et al., 2002).

Campylobacter rectus is a Gram-negative, motile, oral bacterium that has been implicated as a cause of periodontitis (Columbo et al., 2006; Dzink et al., 1985; Lai et al., 1992). Women with periodontitis are seven times more likely to experience preterm labor than their healthy counterparts (Bobetsis et al., 2006; Offenbacher et al., 1996). Serological data has implied that *C. rectus* plays a role in the preterm labors of mothers with periodontitis (Madianos et al., 2001). Additionally, a pregnant mouse model has shown the association of *C. rectus* with a decreased survival of pups (Offenbacher et al., 2005; Yeo et al., 2005). Although these studies have established *C. rectus*

as an agent of periodontitis, the mechanisms of *C. rectus* pathogenesis are not well known.

The genes important to the pathogenesis of *C. rectus* have not been well characterized. A few studies have identified potential virulence factors in *C. rectus*, including toxin genes (*csxA* and *csxB*; *csxC* and *csxD*), a potential cell invasion gene (*ciaB*), and a surface array protein (*crsA* gene) thought to play a role in avoiding the host immune system (Braun et al., 1999; LaGier and Threadgill, 2008; Wang et al., 2000; LaGier and Threadgill, 2014). Within the campylobacters, virulence factors have been best characterized from the gastrointestinal pathogen *Campylobacter jejuni* (Young et al., 2007). In *C. jejuni*, chemotaxis is important to establishing disease, by allowing the bacteria to locate an optimal site for infection (Zautner et al., 2012). Among the chemotaxis proteins in *C. jejuni*, CheY is believed to act as a response regulator that interacts with flagellar motor proteins (specifically FliM, when phosphorylated by the signal-transducing histidine kinase CheA) to influence chemotaxis (Marchant et al., 2002). In addition, mutant *C. jejuni* lacking CheY do not efficiently colonize mice, and do not induce campylobacteriosis in ferret models (Yao et al., 1997).

In this study, a cheY gene from *C. rectus* was discovered and characterized (Cr-cheY) using a combination of bioinformatics, PCR, reverse transcriptase PCR (RT-PCR), and functional complementation experiments in *Escherichia coli*. Interestingly, *C. rectus* appears to be capable of moving within a host from one site to a distant site of infection; specifically, from the subcutaneous tissue to the placenta of pregnant mice (Offenbacher et al., 2005; Yeo et al., 2005). *C. rectus* is also capable of chemotactic behavior in vitro (Paster and Gibbons, 1986). Hence, chemotaxis may contribute to the systemic complications of *C. rectus*-related periodontitis. Additional characterization of *C. rectus* chemotaxis, including Cr-CheY, has the potential to identify new ways to disrupt this process.

MATERIALS AND METHODS

Bacterial strains and DNA isolation

Campylobacter curvus (ATCC 33273) and *Campylobacter rectus* strains 314 and 33238 (LaGier and Threadgill, 2008) were grown under standard anaerobic conditions. *C. curvus*, like *C. rectus*, is an oral bacteria associated with periodontitis (Muchach and Tanner, 2000). Bacteria were grown on tryptic soy blood agar with sodium formate (0.3%). Genomic DNA was isolated from bacterial pellets using CTAB (Ausubel et al., 1990). For the complementation studies, an isogenic mutant strain of *E. coli* lacking CheY function (RP5232, Δ cheY) and the corresponding wild-type strain (RP437, WT) were used (Bourret et al., 1990).

Degenerate PCR

Genomic DNA from *C. rectus* 314 and 33238 were PCR scanned for cheY genes using oligonucleotides 5'-GATATGCCTATYATNATGGTWAC-3' (CheY-DF) and 5'-TTAAWACTTGNGGMGTAAAGG-3' (CheY-DR). CheY-DF and CheY-DR were designed to amplify the entire cheY gene (Wren et al., 1992). Each 50 μ l PCR reaction contained 0.5 μ M of each primer. The cycling conditions used were as follows: 94°C for 1 minute, 45°C for 1 minute, and 72°C for 1 minute (10 cycles); followed by 30 cycles of 94°C for 1 minute, 50°C for 1 minute, and 72°C for 1 minute. Amplicons were purified (QIAquick gel extraction kit, Qiagen, USA) and cloned for sequencing into pCR2.1-TOPO (TOPO-TA kit, Invitrogen, USA).

RNA isolation and reverse transcription PCR

RNA was isolated from bacterial pellets using a QIAgen

miniprep kit (Qiagen, USA). The RNA was then used for RT-PCR using the Access RT-PCR kit (Promega, USA). Primers CheY-RTF1 (5'-ATCCTAGGTTTATATCG-3') and CheY-RTR1 (5'-TAGGTAAAGCCGAGGTC-3') were used for amplification of cheY transcripts.

Recombinant plasmid construction and expression in *E. coli*

The complete open reading frame of *C. rectus* cheY (strain 314, GenBank Accession #EU119866) was cloned (BamHI and HindIII sites) into *E. coli* expression vector pQE80L (Invitrogen, USA) to create pQE80L-Cr-CheY. The engineered plasmid contains the complete cheY gene with an in-frame, N-terminus histidine tag. The fidelity of the construct was checked by DNA sequencing. The plasmid pQE80L-Cr-CheY was transformed into *E. coli* RP5232 and RP437. The expression of pQE80L-Cr-CheY was confirmed by SDS-PAGE analysis (Ausubel et al., 1990), after induction of the protein by the addition of 1mM IPTG to overnight cultures of *E. coli* growing in Luria broth; supplemented with 100 μ g/mL ampicillin.

Swarming motility assays

E. coli swarming assays were carried out as previously described (Bourret et al., 1990; Li et al., 2006). Briefly, the WT (RP5232) and Δ cheY (RP437) strains transformed with pQE80L-Cr-CheY were grown overnight at 37°C in LB supplemented with 100 μ g/mL ampicillin. A 5 μ L sample (106 cells) of each culture was placed in the center of motility agar plates (Li et al., 2006) supplemented with 100 μ g/mL ampicillin and 0.0 or 10.0 μ M IPTG. In pQE80L-Cr-CheY, the expression of the recombinant protein is under the control of an IPTG-inducible promoter. The plates were incubated at 30°C. After 24 hours, swarming was assessed by measuring the distance moved (radii measurements) by *E. coli* cells from the original site of inoculation. Three independent experiments were performed and the swarming radii were averaged. Each experiment contained replicate plates for each condition tested. From each plate, three radii measurements were taken.

PCR-based site directed mutagenesis

The QuikChange II site directed mutagenesis kit (Stratagene, USA) was used to change a residue important to the predicted functionality of Cr-CheY. The kit was used according to the manufacturer's protocols, and the change in

residue was from a conserved aspartate to an alanine. The PCR template used was pQE80L-Cr-CheY, and CheY-MUTF1 (5' - TGCTTATCACCAACTGGA -3') and CheY-MUTR1 (5' -GCATGTTCCAGTCGGTGATA-3') were used as primers to introduce the point mutation. The intended change was confirmed by sequencing the resultant plasmid, pQE80L-Cr-CheY-MUT. This plasmid was subjected to motility assays after transformation into *E. coli* RP5232 and RP437.

Sequence analysis and bioinformatics

DNA chromatographs were edited and assembled using Vector NTI software (Invitrogen, USA). Obtained sequences were then used as templates for sequence homology searches using BLAST. Searches were performed using the default parameters for BLASTx (Altschul et al., 1990). Alignments were generated using T-Coffee (Thompson et al., 1994). The resulting alignment was used to generate a WebLogo (Crooks et al., 2004).

I-TASSER was used to predict a tertiary structure for Cr-CheY (Zhang, 2008).

RESULTS

Identification and expression of *C. rectus* cheY

Degenerate PCR was used to amplify a potential cheY gene from *C. rectus* genomic DNA. PCR yielded an amplicon of the expected size (366 base pairs, Figure 1A). Amplicons of the correct size were also detected from *C. rectus* 33238 and *Campylobacter curvus* 33273. Cr-CheY from strain 314, which is the most widely studied strain of *C. rectus*, was used in all subsequent experiments. RT-PCR showed the expression of Cr-cheY (Figure 1B), with an expected amplicon of 151 base pairs generated from isolated *C. rectus* RNA. DNA sequencing of the degenerate PCR amplicon revealed a Cr-cheY ORF of 366 base pairs. The 366 base pair sequence is available at NCBI (#EU119867.1). The open reading frame (ORF) included a start codon (GTG), stop codon (TAA), and translation of the ORF reveals a protein of 121 amino acids, with a predicted molecular weight of 13.7 Kilodaltons.

Figure 1

A. Agarose gel of *C. rectus* or *Campylobacter curvus* DNA amplified by degenerate PCR. An arrow marks 400 base pairs. B. Agarose gel of *C. rectus* strain 314 cDNA amplified with RT-PCR. No template = No template control, No RT = No Reverse-Transcriptase control. An arrow marks 200 base pairs.

In silico characterization of Cr-CheY

BLASTx analysis of Cr-CheY revealed a *C. rectus* gene sharing significant sequence identity and similarity to CheY proteins from campylobacters, including *C. jejuni* (NCBI #YP_179249.1; 88% identity, and E-value of 5e-71). A WebLogo (Figure 2, based on a T-Coffee alignment) of Cr-CheY showed homology with CheY proteins, including the presence of conserved residues that contribute to the function of CheY as a response regulator. In the model bacterium *E. coli*, CheY interacts with the flagellar motor switch protein FliM (when phosphorylated by the histidine kinase CheA) to influence chemotaxis (Bourret et al., 1990). Cr-CheY contains a conserved site for phosphorylation (Zautner et al., 2012) by CheA (aspartate, D53 in Cr-CheY; D71 in the WebLogo), and residues involved with binding FliM and CheA (Figure 2). The CheY from the related species *C. jejuni* contains the same conserved sites (Yao et al., 1997); and preliminary BLASTx analysis has identified potential FliM (NCBI #WP_002945945.1) and CheA (WP_002943755.1) homologs from *C. rectus*. The structure of Cr-CheY protein was modeled using the I-TASSER server. As shown in Figure 3, the predicted tertiary structure of Cr-CheY shares significant homology with the known structure of a CheY protein from the related epsilon-proteobacterium *Helicobacter pylori* (protein database #3H1F, <http://www.rcsb.org/pdb/home/home.do>); including a conserved arrangement and number of alpha helices and beta strands.

Figure 2

WebLogo of Cr-CheY and related CheY proteins. The identity and position of conserved amino acids within Cr-CheY are indicated. A score of 4.0 bits indicates 100% conservation of the indicated amino acid among the 24 CheY proteins originally aligned with T-Coffee (Materials and Methods). The circled aspartate (D) is phosphorylated by CheA in related CheY proteins. The residues surrounded by rectangles are involved with binding FliM. Residues involved with binding CheA are surrounded by triangles.

Figure 4

Coomassie-stained SDS-PAGE gel (15%) showing expression of recombinant Cr-CheY (from *C. rectus*) in *E. coli*. Equal amounts of cell lysate were loaded per well. The arrow indicates 17 Kilodaltons and the arrowheads denote IPTG-induced Cr-CheY at the expected molecular weight (16 Kilodaltons). Ladder = EZ-Run protein standard (Fisher Scientific, USA); WT = wild type *E. coli*; Δ cheY = isogenic mutant *E. coli*.

The effect of Cr-CheY in WT and Δ cheY *E. coli* was assessed by observing changes in swarming motility that requires CheY functionality, and thus requires chemotaxis. While WT *E. coli* display swarming motility, Δ cheY *E. coli* lack swarming motility after 24 hours of growth (Figure 5, panels A, B, C, D). Cr-CheY expressed in Δ cheY *E. coli* did not restore swarming, and therefore does not directly replace the function of *E. coli* cheY (Figure 5, panel F). However, the expression of Cr-CheY in WT *E. coli* inhibited the swarming ability of WT cells (Figure 5, panel E). The mean radii swarming distance decreased from 40.0 mm to 6.3 mm (Figure 5, compare panels C and E). The inhibition of swarming in WT *E. coli* was not due to differences in growth rates. More specifically, tracking the optical density (600 nanometer readings) of WT *E. coli* carrying Cr-CheY in the presence or absence of 10 μ M IPTG showed no difference in growth rate. After 24 hours of incubation, the mean optical densities for samples minus or plus IPTG were 1.81 and 1.80, respectively. As expected, a mutated version of Cr-CheY; in which a key aspartate residue (D53) has been changed to an alanine, restores 100% of swarming motility when expressed in WT *E. coli* (Figure 5, panel G). The expression of mutated Cr-CheY in WT *E. coli* was confirmed by SDS-PAGE (data not shown).

Figure 5

DISCUSSION

In this study a cheY gene from *C. rectus*, an understudied periodontal pathogen, was identified and characterized. Degenerate PCR, RT-PCR and bioinformatics (Figures 1, 2, and 3) supported the notion that *C. rectus*, which is motile and capable of chemotaxis (Paster and Gibbons, 1986), expresses a gene (Cr-CheY) that shares significant homology with known CheY proteins. *E. coli* strains without CheY lack swarming motility, which is dependent on chemotaxis (Bourett et al., 1990). To begin to characterize the function of Cr-CheY, a recombinant version of the protein was expressed in an *E. coli* strain lacking CheY. Although Cr-CheY did not complement the function of *E. coli* (Figure 5, panel F), the expression of Cr-CheY in wild-type *E. coli* inhibited swarming motility (Figure 5, panel E). The lack of complementation may be due to insufficient functional similarity. In *E. coli*, chemotaxis is dependent on CheY being phosphorylated by a histidine kinase, CheA (Bourett et al., 1990). The mechanism by which Cr-CheY inhibits *E. coli* swarming is unknown. However, the same effect has been observed in *E. coli* complementation experiments using CheY proteins from *Leptospira interrogans* and *Helicobacter pylori* (Li et al., 2006; Foynes et al., 2000). Interestingly, a mutated version of Cr-CheY expressed in wild-type *E. coli*, which lacks an aspartate residue that is phosphorylated during *E. coli* chemotaxis, does not inhibit swarming (Figure 5, panel G). This observation supports speculation that Cr-CheY expressed in wild-type *E. coli* competes with *E. coli* CheY for phosphates from CheA; and thus inhibits swarming

(Foynes et al., 2000; Pittman et al., 2001). Overall, the ability of Cr-CheY to inhibit swarming of *E. coli* suggests that Cr-CheY likely plays a role in *C. rectus* chemotaxis.

Chemotaxis, or directed movement in response to chemical stimulants, has been shown to play a role in the disease-causing capabilities of many bacterial pathogens including *C. jejuni*, *Helicobacter pylori*, *Vibrio cholerae*, and *Listeria monocytogenes* (Yao et al., 1997; Foynes et al., 2000; Butler and Camili 2004; Dons et al., 2004). The protein network in model bacteria that function to direct chemotaxis includes methyl-accepting chemoreceptors (MCPs) that sense changes in extracellular chemical stimulant levels, and cytoplasmic proteins (e.g., cheY, cheA, and cheW) which transmit signals from the MCPs to the flagellum via a series of phosphotransfers (Baker et al., 2006). Although this study focused on CheY, preliminary analysis of the draft *C. rectus* genome suggests that *C. rectus* contains several genes, including MCPs and phosphotransfer proteins, known to play a role in chemotaxis

(https://img.jgi.doe.gov/cgi-bin/w/main.cgi?section=KeggMap&page=keggMap&map_id=map02030&gene_oid=644263128&myimg=0). In addition, this study is the first effort to characterize a cheY gene from any of the oral campylobacters.

In the related pathogen *C. jejuni*, cells lacking CheY do not swarm, colonize mice, and fail to induce disease in ferrets (Yao et al., 1997). *C. rectus* appears to be capable of moving within a host from one site to a distant site of infection; from the subcutaneous tissue to the placenta of pregnant mice (Offenbacher et al., 2005; Yeo et al., 2005). *C. rectus* is also capable of chemotaxis in vitro (Paster and Gibbons, 1986). Most significantly, *C. rectus* is attracted to a potential energy source, formate (Rams et al., 1993), which is also found in periodontal pockets (Paster and Gibbons, 1986). Hence, chemotaxis may contribute to both the systemic complications of *C. rectus*-related periodontitis and assist the bacterium in colonizing subgingival pockets during the development of periodontal disease. Additional studies of *C. rectus* chemotaxis have the potential to discover new approaches for disrupting this process.

This study was funded, in part, by NIH grant # K12GM000678.

References

Altschul S et al (1990) Basic local alignment search tool. *J Mol Biol* 215: 403-10.

Ausubel F et al (1990) Current protocols in molecular biology. New York: John Wiley and Sons.

Baker et al (2006) Signal transduction in bacterial chemotaxis. *BioEssays* 28: 9-22.

Bobetsis YA, Barros SP, and Offenbacher S (2006) Exploring the relationship between periodontal disease and pregnancy complications. *J Am Dent Assoc* 137: 7S-13S.

Bourret RB et al (1990) Conserved aspartate residues and phosphorylation in signal transduction by the chemotaxis protein CheY. *Proc Natl Acad Sci USA* 87: 41-45.

Braun M et al (1999) Cloning and characterization of two bistructural S-layer-RTX proteins from *Campylobacter rectus*. *J Bacteriol* 181: 2501-6.

Butler SM and Camili A (2004) Both chemotaxis and net motility greatly influence the infectivity of *Vibrio cholerae*. *Proc Natl Acad Sci USA* 101: 5018-5023.

Colombo AV et al (2006) Identification of oral bacteria associated with crevicular epithelial cells from chronic periodontitis lesions. *J Med Microbiol* 55: 609-15.

Crooks GE et al (2004) WebLogo: a sequence logo generator. *Genome Res* 14: 1188-1190.

Dons L et al (2004) Role of flagellin and the cheA/cheY system of *Listeria monocytogenes* in host cell invasion and virulence. *Infect Immun* 72: 3237-3244.

Dzink JL et al (1985) Gram negative species associated with active destructive periodontal lesions. *J Clin Periodontol* 5: 648-59.

Foynes S et al (2000) *Helicobacter pylori* possesses two cheY response regulators and a histidine kinase regulator, cheA, which are essential for chemotaxis and colonization of the gastric mucosa. *Infect Immun* 68: 2016-2023.

LaGier MJ, and Threadgill DS (2008) Identification of novel genes in the oral pathogen *Campylobacter rectus*. *Oral Microbiol Immunol* 2008 23: 406-12.

LaGier MJ, and Threadgill DS (2014) Identification and characterization of an invasion antigen B gene from the oral pathogen *Campylobacter rectus*. *Indian J Microbiol* 54: 33-40.

Lai CH et al (1992) *Wolinella recta* in adult gingivitis and periodontitis. *J Periodontal Res* 27:8-14.

Li ZH et al (2006) Characterization of the cheY genes from *Leptospira interrogans* and their effects on the behavior of *Escherichia coli*. *Biochem Biophys Res Commun* 345: 858-866.

Madianos PN et al (2001) Maternal periodontitis and prematurity Part II: Maternal infection and fetal exposure. *Ann Periodontol* 6: 175-82.

Marchant J, Wren B, and Ketley J (2002) Exploiting genome sequence: predictions for mechanisms of campylobacter chemotaxis. *Trends Microbiol* 10: 155-159.

Mucach P and Tanner AC (2000) *Campylobacter* species in health gingivitis and periodontitis. *J Dent Res* 79: 785-92.

Offenbacher S et al (1996) Periodontal infection as a possible risk factor for preterm low birth weight. J Periodontol 67: 1103-13.

Offenbacher S et al (2005) Effects of maternal *Campylobacter rectus* infection on murine placenta fetal and neonatal survival and brain development. J Periodontol 76: 2133-43.

Paster BJ and Gibbons RJ (1986) Chemotactic response to formate by *Campylobacter concisus* and its potential role in gingival colonization. Infect Immun 52: 378-382.

Pihlstrom BL, Michalowicz BS, and Johnson NW (2005) Periodontal diseases. Lancet 366: 1809-20.

Pittman MS et al (2001) Chemotaxis in the human gastric pathogen *Helicobacter pylori*: different roles for cheW and the three cheV paralogues, and evidence for cheV2 phosphorylation. Microbiol 147: 2493-2504.

Rams TE et al (1993) *Campylobacter rectus* in human periodontitis. Oral Microbiol Immunol 8: 230-235.

Thompson JD, Higgins DG, and Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting position-specific gap penalties and weight matrix choice. Nucleic Acids Res 22: 4673-80.

Teng YT et al (2002) Periodontal health and systemic disorders. J Can Dent Assoc 68: 188-92.

Wang B, Kraig E, and Kolodrubetz D (2000) Use of defined mutants to assess the role of the *Campylobacter rectus* S-layer in bacterium-epithelial cell interactions. Infect Immun 68: 1465-73.

Wren BW et al (1992) Degenerate PCR primers for the amplification of fragments from genes encoding response regulators from a range of pathogenic bacteria. FEMS Microbiol Lett 78: 287-91.

Yao R, Burr DH, and Guerry P (1997) CheY-modulation of *Campylobacter jejuni* virulence. Mol Microbiol 23: 1021-1031.

Yeo A et al (2005) *Campylobacter rectus* mediates growth restriction in pregnant mice. J Periodontol 76: 551-7.

Young KT, Davis, LM and Dirita VJ (2007) *Campylobacter jejuni*: molecular biology and pathogenesis. Nat Rev Microbiol 5: 665-79.

Zuatner AE et al (2012) Chemotaxis in *Campylobacter jejuni*. Eur J Microbiol (Bp) 2: 24-31.

Zhang Y (2008) I-TASSER server for protein 3D structure prediction. BMC Bioinformatics 9: 40.

Author Information

Michael J. LaGier

Department of Biology, Grand View University

Des Moines, IA, USA

mlagier@grandview.edu

Ihor Bilokopytov

Department of Biological Sciences, Florida Gulf Coast University

Fort Myers, FL, USA

Bradley Cockerill

Department of Biological Sciences, Florida Gulf Coast University

Fort Myers, FL, USA

Deborah S. Threadgill

Department of Veterinary Pathology, Texas A&M University

College Station, TX, USA