Promoting Homeostasis To Avoid Rebound Weight Gain In Yo-Yo Dieters

L Outland, M Stoner-Smith

Citation

L Outland, M Stoner-Smith. *Promoting Homeostasis To Avoid Rebound Weight Gain In Yo-Yo Dieters*. The Internet Journal of Advanced Nursing Practice. 2013 Volume 12 Number 1.

Abstract

Rebound weight gain is more often the rule than the exception and the reason may not be lack of willpower. Recent findings in the study of weight homeostasis have uncovered a host of orexigenic and anorexigenic hormones working to keep a stable healthy weight. When energy stores are depleted through dieting, overcompensation to replenish them often occurs. Overeating is one of the homeostatic responses. However, there are other involuntary responses to hunger that lower metabolism, reduce thyroid hormones, and produce fat cells. Traditional weight control advice to cut down on caloric intake may inadvertently trigger the compensatory mechanism that lead to rebound weight and yo-yo dieting. Tuning into homeostatic cues to prevent excessive hunger without getting too full is the hallmark of intuitive eating. Tips are given on how to eat intuitively in order to promote homeostasis. Supporting homeostasis will help keep weight stable and prevent weight cycling and rebound weight gain that often accompanies dieting.

WEIGHT CYCLING THE RESULT OF DIETING

Cyclic weight gain and loss (yo-yo dieting) is a common problem with potentially more negative health effects than overt obesity.1, 2, 3,4 The phenomenon of weight gain after weight loss is more common than keeping the weight off after a weight loss plan.5,6 The weight regain that often occurs may in part be responsible for the excess weight plaguing much of the developed and developing world.7,8 The dual existence of record numbers of dieters, and overweight individuals may not be the paradox it appears.9,10,11,12 Physiologic research is now uncovering reasons why the more most of these individuals diet the heavier they become. As health care providers concerned with the health of patients, advanced practice nurses (APN) might want to examine the standard for patient education around weight control.13 In light of two decades of discoveries in the field of weight homeostasis, the upshot appears to be that cutting down and eating less may be at the root of much yo-yo dieting and the excess weight seen in chronic dieters.14 This paper will examine the wisdom of this advice in light of the effects hunger has on the complex system of weight homeostasis. Finally, a step-by-step alternative that providers can give patients will be provided.

PATIENT

A 42-year female patient presented to the clinic for a routine

physical. After her blood pressure and weight were taken, before beginning the history, the APN noted that the patient's weight had increased 43 pounds since her last appointment 2 years ago, putting her at a body mass index (BMI) of 30. In the 8 years that this patient had been coming to the clinic, she had gained and lost the same 40–50 pounds three times, and in fact, she was almost 20 pounds heavier than she was before her first appointment. The provider was unsure how to proceed. This patient always vowed to lose weight, often succeeded, only to gain the weight back again months or years later.

WEIGHT HOMEOSTASIS RESISTS WEIGHT LOSS

Not Willpower

When patients present appointment after appointment with excess weight despite health care advice, both patients and providers can be frustrated. This frustration is often intensified when the patient has other comorbid disorders like diabetes and hypertension. However, for patients who are becoming yo-yo dieters, it may not be a "loss of will power" that is keeping them heavy. Recent physiologic studies indicate that rebound weight gain may have little to do with weak resolve and more to do with homeostasis of energy (weight homeostasis).15, 16, 17, 18 This rebound weight gain what happens with energy homeostasis and dieting. The human body is confronted with a loss of energy (dieting). That loss of energy is perceived by homeostatic processes as a disturbance which is then responded to as a threat. Perhaps because loss of weight can be deadly faster that weight gain, homeostasis tends to err on the side of resisting "starvation" over the promotion of weight loss.19

Weight Homeostasis 101

Homeostasis is a well-known topic for health care providers. When pH balance rises, respirations automatically increase to rid the body of bicarbonate ions. However, the restoration to a normal level could also occur voluntarily. For example, in the critical care unit when pH drops, a patient is sometimes given bicarbonate. If a patient is cold, shivering may ensue involuntarily. At the same time, warmed blankets can be provided. In each of these cases automatic and voluntary ways are available to restore homeostasis. As an example of energy homeostasis, when calories are withheld to the point of hunger, the body becomes flooded with hunger hormones.20 The hungry individual is spurred to eat, and thus eats alot.21 Eating in response to hunger is a voluntary way to replenish lost energy stores. However, the body has a built in way to involuntary stop further loss of energy. The complex cascade of homeostatic biochemicals produces fat cells22, decreases resting energy metabolism23, decreases thermogenesis, and decreases thyroid secretion.24 These processes are all automatic ways to restore lost energy.

Many biochemicals are involved in energy homeostasis. Some are secreted when hunger arises and are termed orexigenic, and some work to create a feeling of fullness and are thus termed anorexigenic. The most common orexigenic hormones are the gut hormone ghrelin 25 and neuropeptide Y26. Peptide YY (PYY), a gut hormone, and leptin, a hormone produced by fat cells are anorexigenic hormones.17, 27 These substances are just some of the peptides, proteins, and hormones that will reduce the loss of energy automatically, despite the desire of the dieting individual.

Weight Homeostasis Errs on the Side of Preventing Weight Loss

To understand the uniqueness of weight homeostasis, it is important to note that unlike pH and temperature, where either too high or too low levels are equally dangerous, sudden or rapid weight loss is more immediately dangerous than weight gain. Throughout human history, survival depended on maintaining adequate weight, which sometimes meant the body adapted to changing conditions through erring on the side of preventing weight loss when extreme energy loss is threatened 28.

Simply starting an infusion of an anorexic agent like PYY to create a feeling of satiety and avoid overeating is not an option. Due to the principles of weight homeostasis that prioritize preventing starvation, an infusion of PYY (the peptide that turns off the desire to eat) only has a short term effect. To prevent weight loss, the body develops a tolerance to the peptide over time.27 Likewise, leptin, the long term satiety hormone might cause a decrease in appetite in the beginning of weight gain, but resistance soon develops and appetite returns to pre–weight gain levels.29, 30

Supporting the bias in favor of maintaining weight for survival, no such tolerance is found with the hunger hormones.31, 32 As a result, high levels of neuropeptide Y and ghrelin continue to stimulate feelings of hunger indefinitely.26 To further reduce the loss of energy, high ghrelin levels not only create fat, increase appetite and food intake, as mentioned previously, they prevent the loss of energy by decreasing thyroid function and decreasing thermogenesis.33,34,35

A POSSIBLE SOLUTION

Both providers and patients who are yo-yo dieters would do well to focus on promoting homeostasis rather than trying to overcome it. Despite the human body being hardwired to resist starvation, it is also programmed to keep body weight within a healthy range. Evidence can be found in the fact that anorexigenic gut hormones outnumber orexigenic hormones 5 to 1.36 Eating 3 meals a day in response to metabolic hunger, supports homeostasis.37, 38 Skipping a meal and not eating for a long period of time increases ghrelin levels and leads to extra weight.39 The other extreme eating behavior, regular overeating, will also increase weight and create the same tolerance that a constant infusion of peptide YY would have.40 Eventually, despite extra weight, the individual will eat as much as before the weight was put on.41 The key is to prevent threatening the homeostatic system by getting too hungry or depleting energy stores. As long as homeostasis is not profoundly or constantly disturbed, by either ignoring fullness and overeating, or ignoring hunger and under-eating, an even, healthy weight will be the result. The way to achieve a healthy weight is to eat when hungry and stop eating when full.

How to Tune into Metabolic Hunger and Fullness

One of the problems facing chronic dieters is that they do not know when they are ideally hungry. Hunger has been the enemy for so long that they have "turned off" the hunger signal. If a chronic dieter has this problem they may be helped by following tips in Table 1.

Table 1a

Tips to Help Your Patients Eat Healthier

- Say YES to giving your body the amount and type of food to be healthy
 The benefit of eating intuitively allows people to eat with a positive attitude. Instead of
 musing, "Oh no, I shouldn't eat because I'm so fat", the approach is positive, "I'm
 hungry, I should eat until I'm satisfied". When full tell yourself "I have had enough".
 Intuitive or homeostatic eating entails several key practices.
- · Key practices of ideal/intuitive eating
- Begin a meal or snack when IDEALLY hungry. Ideally meaning nice and hungry without hunger pangs or a rumblings.
- Eat until IDEALLY satisfied. Ideally satisfied means you will not be hungry again for 4-6 hours, and have not gotten too full, or "stuffed".
- Choosing what your body is craving from your favorite foods that are both delicious and healthy
- · How to know when you are hungry

The cold water test

- > 4.5 to 5 hours after eating a meal drink a cold glass of water
- > If you feel it coat the bottom of your stomach you are ideally hungry

Table 1b

Tips to Help Your Patients Eat Healthier

How to know when you are full

Taste experiment

- When you become ideally hungry, provide yourself with a fist sized portion of delicious, healthy food
- > Close your eyes and notice flavors as you eat your portion
- > Stop and wait 10 minutes
- Put another bite of the same food in your mouth. Notice the flavors again if the food does not taste as good this time as it did when you began, you are probably full
- Get some help
 - There are many books and programs out there to help people eat intuitively, from Geneen Roth⁴² to the Gabriel Method⁴³ reference, to the new iPhone App, 'Leat' (IWC, for Intuitive Weight Control). These resources explain the reasons to eat intuitively, what to expect when you go from dieting to eating intuitively, and give you tools to help you nail every meal eating exactly what your body needs; no more, no less.
 - Instead of counting calories or keeping a food journal, keep a hunger/ fullness log, noting how hungry you are when you begin each meal or snack and how full you get when you stop.
 - > Find a group to support you in your life style change and an activity you enjoy doing

HOW INTUITIVE EATING HELPED THE PATIENT

Finding Out What She Was Doing Wrong

After several months of attempting to eat intuitively and keeping a log, the patient realized she ate all of her daytime meals extremely hungry and without getting full enough. She would eat a very small breakfast and lunch, then nothing until dinner. The sparse eating pattern put her body into starvation mode and resulted in such fierce hunger signals that she was famished by dinner. Her hunger level impelled her to eat too much at dinner and continue eating throughout the evening. In reviewing her food log with a nutritional consultant, the patient saw her issue was not a lack of willpower that led to over eating, but was her body's cascade of chemicals helping her not starve to death. In her very empty stomach throughout the day, the hunger hormone ghrelin was being pumped out by gastric cells and causing levels to soar. The desire to eat was chemical44, something she could not control but could prevent.

Resetting the Body's Signals

Once the patient started eating a healthy breakfast when she got hungry, and doing the same at lunch, over eating at dinner and throughout the evening started to be less of a problem. Because she now responds to her hunger hormones and eats proportionately, her total food consumption is less.

CONCLUSION - SIMPLE BUT NOT EASY

Intuitive eating is simpler than counting calories but may be difficult for the chronic dieter.44 A person who starts practicing intuitive eating is encouraged to tune in to hunger feelings instead of tuning them out.45 Intuitive eating helps chronic dieters change their tuning out behavior and notice when they are ideally hungry. It is healthy to eat when hungry.46 The complex mechanism of weight homeostasis is preprogrammed to ensure our bodies have a certain amount of weight.28 Eating just enough to be satisfied is the antidote to hunger and excess weight. The key is to stop eating when ideally full. The exact amount of food and the frequency and timing of meals will vary from person to person. Giving patients permission to eat when hungry and stop eating when full will not only break the cycle of yo-yo dieting, but will help patients make peace with themselves, and help providers make peace with their patients.47 Π

References

1. Gaesser GA. Weight, weight loss, and health: a look at the evidence. Healthy Weight J. 2003;(17):8-12.

2. Bacon L, Aphramor L. Weight science: evaluating the evidence for a paradigm shift. Nutr J. 2011;(10):69. 3. Rzehak P, Meisinger C, Woelke G, et al. Weight change, weight cycling and mortality in the ERFORT male cohort study. Eur J Epidem. 2007; 2210(10): 665-673. 4. Graci S, Izzo G, Savino S, et al. Weight cycling and cardiovascular risk factors in obesity. Int J Obes.

2004;28(1):65-71. 5. Korkeila M, Rissanen A, Kaprio J, et al. Weight-loss attempts and risk of major weight gain: a prospective study in Finnish adults. Am J Clin Nutr. 1999;70(6):965-975. 6. Mann T, Tomiyama AJ, Westling E, et al. Medicare's search for effective obesity treatments: diets are not the

answer. Am Psychol. 2007;62(3):220-233. 7. Jeffery RW, Adler SA, Forster JL. Prevalence of dieting among working men and women: the healthy worker project. Health Psychol. 1991;10(4):274-81.

8. Heini A, Weinsier R. Divergent trends in obesity and fat intake patterns: the American paradox. Am J Med. 1997;102(3):259-264.

9. Field AE, Austin SB, Taylor CB, et al. Relation between dieting and weight change among preadolescents and adolescents. Peds. 2003;112(4):900-906.

10. Neumark-Sztainer M, Wall M, Haines J, Story M, Eisenberg M. Why does dieting predict weight gain in adolescents? Findings from Project EAT-II: a 5-year longitudinal study. J Am Diet Assoc. 2007;107(3): 448-455. 11. Schur EA, Cummings DE, Callahan HS, Foster-Schubert KE. The association of cognitive restraint with ghrelin, leptin, and insulin levels in subjects who are not weight –reduced. Phys Behav. 2008;93(4-5):706-712.

12. Crujeiras AB, Goyenechea È, Abete I, Lage M, Carreira MC. Weight regain after a diet-induced loss is predicted by higher baseline leptin and lower ghrelin plasma levels. J Clin Endocrinol Metab. 2010;95(11):5037-44. 13. US Preventive Services Task Force: Behavioral

counseling in primary care to promote a healthy diet: recommendations and rationale Am J Prev Med. 2003;(24):93-100.

14. Appelhans, BM, Whited MC, Schneider K L, et al. Time to abandon the notion of personal choice in dietary counseling for obesity? J Am Diet Assoc. 2011;111(8): 1130-1136.

15. Myers MG Jr, Leibel RL, Seeley RJ, et al. Obesity and leptin resistance: distinguishing cause from effect Trends Endocrin Metab. 2010;21(11):643-651.

16. Wren AM, Small CJ, Ward HL, et al. The novel hypothalamic peptide ghrelin stimulates food intake and growth hormone secretion. Endocrinol. 2000;(141):4325-4328.

17. Friedman JM, Halaas JL. Leptin and the regulation of body weight in mammals. Nature. 1998; 395;(6704):763-770.

18. French SA, Jeffery RW. Consequences of dieting to lose weight: effects on physical and mental health. Health Psychol. 1994; 13(3):195-212.

19. Yamada T, Katagiri H. Avenues of communication between the brain and tissues/organs involved in energy homeostasis. J Endocrinol. 2007(54):497-505.

20. Gil-Campos M, Aguilera CM, Cañete R, et al. Ghrelin: A hormone regulating food intake and energy homeostasis. Br J Nutr. 2006;(96): 201-226.

21. Druce MR, Wren AM, Park AJ, et al. Ghrelin increases food intake in obese as well as lean subjects. Int J Obes (London). 2005;(29):1130–1136.

22. Tschop M, Smiley DL, Heiman ML. Ghrelin induces adiposity in rodents. Nature. 2000;407 (6806):908-913. 23. Marzullo P, Verti B, Savia G, et al. The relationship between active ghrelin levels and human obesity involves alterations in resting energy expenditure. J Clin Endocrinol Metab. 2004;89(2):936-939.

24. Castañeda TR, Tong J, Datta R, et al. Ghrelin in the regulation of body weight and metabolism . Front

Neuroendocrinol. 2010;(31):44-60. 25. Kojima M, Hosoda H, Date Y, et al: Ghrelin is a growthhormone-releasing acylated peptide from stomach. Nature. 1999;402(6762):656-660.

26. Chen HY, Trumbauer ME, Chen AS, et al. Orexigenic action of peripheral ghrelin is mediated by neuropeptide Y and agouti-related protein. Endocrinol. 2004;145 (6):2607-2612

27. Renshaw D, Batterham RL. Peptide YY: a potential therapy for obesity. Curr Drug Targets; 2005;(6):171-179. 28. Outland L. Bringing homeostasis back into weight control. J Obes Wt Loss Ther. 2012;(2):115. doi:10.4172/2165-7904.1000115

29. Belgardt BE, Brüning JC. CNS leptin and insulin action in the control of energy homeostasis. Ann N Y Acad Sci. 2010;1212:97-113.

30. Schur EA, Cummings DE, Callahan HS, Foster-Schubert KE. The association of cognitive restraint with ghrelin, leptin, and insulin levels in subjects who are not weight -reduced. Physiol Behav. 2008; 93(4-5):706-712.

31. Unniappan S, Kieffer TJ. Leptin extends the anorectic effects of chronic PYY(3-36) administration in ad libitumfed rats. Am J Physiol Regul Integr Comp Physiol. 2008;295:R51-58.

32. Cummings DE. Ghrelin and the short- and long-term regulation of appetite and body weight. Physiol Behav. 2006;89(1):71-84.

33. Dulloo AG, Calokatisa R. Adaptation to low calorie intake in obese mice: contribution of a metabolic component to diminished energy expenditures during and after weight loss. Int J Obes. 1991;15(1):7-16.

34. Meier U, Gressner AM. Endocrine regulation of energy metabolism: review of pathobiochemical and clinical chemical aspects of leptin, ghrelin, adiponectin, and resistin; Clin Chem. 2004;50(9): 1511-1525.

35. Nogueiras R, Tschöp MH, Zigman JM. Central nervous system regulation of energy metabolism: ghrelin versus leptin. Ann N Y Acad Sci. 1126:14-19.

36. Chaudhri O, Small C, Bloom S. Gastrointestinal

hormones regulating appetite. Philos Trans R Soc Lond B Biol Sci. 2006; 361;1471:1187-1209.

37. Ciampolini M, Borselli L, Giannellini V. Attention to metabolic hunger and its effects on Heliocobacter pylori infection. Physiol Behav. 2000;70:287-296.

38. Bellisle F, McDevitt R, Prentice AM. Meal frequency

and energy balance. Br J Nutr. 1997;77: 57-70. 39. MacLean PS, Higgins JA, Jackman MR, et al. Peripheral

metabolic responses to prolonged weight reduction that promote rapid, efficient regain in obesity-prone rats. Am J Physiol Regul Integr Comp Physiol. 2006;290

(6):R1577-1588.

40. Ryan KK, Woods SC, Seeley RJ. Central nervous system mechanisms linking the consumption of palatable high-fat diets to the defense of greater adiposity. Cell Metab. 2012;(8);152:137-149.

41. Moss C, Dhillo WS, Frost G, et al. Gastrointestinal hormones: the regulation of appetite and the anorexia of ageing. J Hum Nutr Diet. 2012;25(1):3-15.

42. Roth G. Emotional Eating. http://www.geneenroth.com/ 43. Gabriel J. The Gabriel Method. NewYork , NY: Atria Books, Simon-Schuster; 2008.

44. Outland L. Intuitive eating: a holistic approach to weight control. Holist Nurs Pract. 2010; 24(1):35-43.

45. Tribole E, Resch E. Intuitive Eating. New York, NY: St Martin's Press; 2003.

46. Ciampolini M. Requested meals versus scheduled meals. Int J Gen Med. 2012;5:345–353. Published online April 13, 2012.

47. Bacon, L. End the war on obesity: make peace with your patients. Medscape Gen Med. 2011; 8(4):40.

Author Information

Lauren Outland, DrPH, MSN, APRN Assistant Professor of Nursing California State University Dominguez Hills Carson, CA, USA

Marilyn Stoner-Smith, PhD, RN Professor of Nursing

California State University, San Bernardino San Bernardino, CA, USA