

# Using the Lagree Megaformer TM for Rehabilitation in Patient with Severe Neuromuscular Dysfunction and Deconditioning: A Case Report

M Greenberg, A Brzenski

---

## Citation

M Greenberg, A Brzenski. *Using the Lagree Megaformer TM for Rehabilitation in Patient with Severe Neuromuscular Dysfunction and Deconditioning: A Case Report*. The Internet Journal of Neurology. 2016 Volume 19 Number 1.

DOI: [10.5580/IJCH.43569](https://doi.org/10.5580/IJCH.43569)

## Abstract

The Lagree Megaformer, (with the most current version, the M3S and Supraformer) is a spring based resistance trainer, which can be used as a therapy device, to train patients with neuromuscular deconditioning. Due to the mechanical workings of the machine, the springs, which normally confer the resistance, can also be used to support the patient during exercise. Once the patient has regained function, support can be gradually removed and resistance added until the patient is fully recovered and beyond. We recommend physical therapists, chiropractors, sports medicine physicians and other practitioners treating debilitated patients with neuromuscular disorders consider using this device as part of a therapeutic treatment plan.

## INTRODUCTION

Since its introduction into the fitness world by creator Sebastien Lagree, the Lagree Fitness (LF) method, with its spring-based trainer the Megaformer, has been a state of the art tool for body sculpting and improving dynamic fitness. Its utility is apparent in subjects new to exercise, as well as, experienced athletes. Through several re-designs, the machine has evolved to devices (Megaformer M3S, Supraformer) that can be used to work any part of the body in all planes, especially the core musculature, (Figures 1&2). While there are many differences compared to the traditional Pilates Reformer, a unique feature of the Megaformer is the ability for the subjects to stand on the surface, training balance and strength. While doing LF training, the subject is taught to move the target muscle very slowly through the full range of motion, sequentially recruiting muscle fibers until muscular failure occurs. Once the muscle has been worked to exhaustion, the subject's myocytes will hypertrophy, increasing their capacity for work. The net effect in athletic subjects is increased muscle strength and definition. LF training also promotes the neuro-muscular connection needed to train stability musculature often neglected in standard weight training. When using the Megaformer in patients with severe debilitation, or neurologic damage, modifications are needed to accommodate the disability. The aim of this case report is to

illustrate the outcome of a patient treated with LF training and describe some of the modifications to the typical LF training session that are needed when treating deconditioned and disabled patients.

## CASE REPORT

The patient is a 38 year-old male with severe atrophy and weakness, primarily in his lower extremities, below the knee (Figure 3). The patient had normal growth and development, participating in cross-country running until age 17, after which he began to notice increasing weakness, especially in his lower legs. Over the next few years he became progressively weaker in his legs until he was no longer able to ambulate, using a motorized wheel chair and walker to get around. While he did experience some weakness in his hands, his cognition and motor function above the knees were preserved. His diagnostic evaluation was extensive, including MRI, EMG, nerve conduction studies, and blood tests. These tests were all non-diagnostic. Genetic testing including exon analysis was done, but did not lead to a diagnosis. He had mildly elevated CPK at baseline (125mg/dl), but his other labs were normal. EMG and nerve conduction were consistent with atrophy and slow neural transmission. On physical exam he had essentially no function and extreme wasting of his gastrocnemius and tibialis anterior. He had 2/5 strength in hamstrings,

quadriceps and gluteus maximus muscles, bilaterally. His hip flexors and abdominal musculature function, while deconditioned, was intact. He had 4/5 strength in his upper extremities, except for the intrinsic hand muscles, which had mild weakness and contractures. On presentation he was able to perform abdominal exercises, but was unable to lunge or squat to any degree. He had extreme difficulty using a walker.

After 20 years, this man still had no medical diagnosis and his strength showed no signs of spontaneous recovery. He had tried traditional physical therapy but he felt he was not improving. The patient made the decision to try exercise therapy of his own choice. His rehabilitation program included daily Bikram hot yoga and two 30 minute, LF sessions per week. Initially he was unable to stand or do any standing exercises without assistance. The initial LF sessions focused on strengthening his core muscles in all 3 planes, in both flexion and extension, while gently working the upper legs, and arms. The ultimate goal was self-ambulation.

Due to his severely weak lower legs, the initial target of the therapy was his upper legs and hips. In contrast to standard LF workouts where a single side of a target muscle group is worked to exhaustion, in this patient, we limited the workout to one exercise per side at a time. This was done to allow more recovery time so he could get more total workload. A sample exercise set is displayed below (Figure 4). Over the course of 6 months he was gradually able to increase the load and duration of the sets. His form during the exercise sessions significantly improved, as he was very unstable at the beginning of the program. We were able to add squats and lunges, first assisted then unassisted. To add lunges, the patient was started with the ‘side lunge’ exercise and the springs were adjusted to give maximal assistance (3-4 yellow springs). As he progressed he was able to tolerate removal of springs due to increased strength in quads, glutes and hamstrings. The most remarkable improvement was his ability to control the movement and position of his legs. This was accomplished using visual and tactile cues from the therapist while repeating simple movements over many repetitions. Over time he was able to regain control his movements, which showed shaking and poor coordination at the start of the therapy

During the later portion of his rehabilitation, muscular endurance improved to the point he was able to swim laps, bike 20 miles, and walk over 1 mile without assistance. To provide motivation, we made a goal to compete in an

endurance race at his one-year anniversary. Recently he was able to achieve that goal and finish a mini sprint triathlon after 12 months, of continuous training.

## **DISCUSSION**

Physical therapy has been the mainstay of treatment for neuromuscular diseases that result in peripheral muscle weakness, such as cerebral palsy and multiple sclerosis. However, in the 20th century controversy existed with the use of strengthening exercises in these patients, with concern that increased muscle mass might lead to increased muscular spasticity with little improvement in overall function. Recent reviews have demonstrated that dynamic exercises, such as swimming (1), as well as progressive resistance exercise (2), increase overall strength(3,4) while not leading to increased spasticity (5,6). The long-term effects, including increased societal participation, are much more difficult to study and current studies have not specifically used this as an endpoint. However, this initial positive data coupled with the patient’s desire to increase motor strength and function, led us to undertake a rehabilitation program with the Lagree fitness method.

Due to the innovative design, the Lagree fitness method using the Megaformer is optimally suited to rehabilitation type exercise. While known primarily for its use in the fitness industry, the Megaformer has several mechanical features that make it ideal for use in rehabilitation. First, the Megaformer has a very solid and stable platform. Patients of all weight, including morbidly obese patients, will be able to use the machine safely. The platform allows multiple positions to be used such that patient can be sitting, standing, lying down supine, or prone, which is helpful for disabled patients. Further, with over 300 defined exercises and countless user described modifications, there is a very large selection of ways to train each muscle group. The springs provide many different resistances, allowing the resistance to be adjusted in even severely weak patients, as in this case. Counter intuitively, exercises done on the front of the machine, get easier as resistance is added. This is ideal for the weak patient, since many will only be able to output a small amount of force. While doing exercises on the front of the machine (like ‘side lunge’), the therapist can add springs which support the patient while allowing them to experience work at the target muscles. This is accomplished by increasing the number of yellow springs (the lightest tension) on the machine. This is in contrast with most gym equipment, which the patient is required to do all of the

work, creating frustration during therapy.

There are several other differences in the LF regiment between training fitness clients and rehabilitation patients (Figure 5). Setting up the sequence of exercises is especially important with the more disabled patients. Its important not to push the patient past his physical limits to ensure they can get through the session without severe discomfort or injury. Unlike the typical fitness client, the rehabilitation patient will need more recovery between sets. Initially it may be helpful to perform the exercise on the left then right, instead of working one side to exhaustion, as is normally done for fitness training. Another difference is setting up the patient to promote stability. This enables the patient to focus on the exercise without the likelihood of falling over or injury. This is in direct contrast to fitness clients. While training to maximize fitness gains, clients are purposely placing in positions of instability, forcing the client to use stabilizing muscles. This instability is unique compared to standard fitness equipment, like a universal gym machine, providing a definite benefit of LF training for athletics. However, given weakness in many rehabilitation patients, this unique aspect should be avoided initially, preventing additional injury during the rehabilitation program.

The rehabilitation patient will need more supervision and assistance than typical fitness clients. Initially, these patients should not be treated in a class environment. Individual supervision will improve safety and decrease frustration. Advanced planning and evaluation of the patient's injury, or current physical status, will ensure the patient receives customized exercise, based on their specific disability. As the patient improves, they should require decreasing supervision, enabling possible transition to group classes.

Establishing expectations prior to the initiation of a rehabilitation regiment is essential to success in these patients. Some disabled patients may be able to do very little work during the initial sessions due to extreme deconditioning, requiring patience on the part of the therapist/instructor. Unlike what is popularized on television, complete reversal of the disease through miracle cures will not be likely. However, with consistent training and optimization of the patient's nutritional intake, slow steady gains can occur. Independent of the improvement in muscular strength, the patient will make neuro-muscular connections that will mature over time, through re-wiring. However, it is essential to be careful to avoid overtraining and injury. Initially, 1-2 session per week is sufficient for

most patients to allow time for adequate recovery. This can be increased to 3-4 sessions as the patients' muscular strength and endurance improve. Optimizing the patients' nutritional intake will further maximize the muscular gains, which occur between training sessions.

Rehabilitation through physical therapy can be extremely challenging for patients with advanced neurologic disease. It takes dedication to any program if the patient is to receive benefit. The patients may be experiencing other coexisting medical disease, such as cardiac failure, COPD, sleep apnea, diabetes mellitus in addition to their neurologic dysfunction. Medical evaluation and clearance will be necessary before initiating any rehabilitation program. In addition to other systemic co-morbidities, a diagnosis of depression may also be present. The mind-body connection has been shown to have a significant impact on overall health (7,8,9) and recovery (10). Untreated depression can slow the rehabilitation process and should be treated via a multimodal approach. While physical and occupational therapy is the standard of care for rehabilitation, other modalities can play a positive role in successful recovery. Yoga, endurance and strength training all will benefit patients who are motivated to recover and are able to tolerate the effort needed to perform these activities.

## **CONCLUSION**

Rehabilitation of deconditioned muscles is a difficult and complex process. In many cases neurological and psychological dysfunction is ongoing. The Lagree Fitness training with the Megaformer can be a valuable tool in this patient population. Disabled patients can improve muscular strength, endurance, coordination, and balance using a modified version of LF training. For appropriate patients in a well-monitored and controlled environment, LF training can be used to reverse the debilitating effects of many neurologic disorders. Medical clearance should always be sought before starting any rehabilitation program.


**Figure 1**

Lagree Fitness Megaformer



**Figure 2**

Lagree Fitness Supraformer



**Figure 3**

Patient getting therapy on the Lagree Fitness Megaformer M3S



**Figure 4**

Sample routine (30 minutes)

Wheelbarrow  
Pushups  
Superman chest pre  
Plank hold

Side Lunge L leg  
Side Lunge R leg  
Bungee Press L leg  
Bungee Press R leg  
Side leg press L leg  
Side Leg press R leg

Leg circles > reverse direction  
Double leg strap press  
Glute raises

Megaformer crunches

Sexy back  
Triceps press  
Biceps curls

**Figure 5**

The differences between fitness clients and rehabilitation patients while on the Megaformer

	<b>Fitness client</b>	<b>Rehabilitation patient</b>
Supervision	Group dynamic improves performance. Assistance infrequently required.	Private training improves safety and ensures customized workouts tailored to the patient's disability. Assistance frequently required.
Muscle activation strategy	Work one muscle group, on each side. Avoid going back and forth between sides	Perform each exercise on the right and left side before moving to next exercise
Recovery	Work muscle group to failure. No recovery between sets on the same muscle group	Allow ample recovery time
Resistance	Apply enough resistance to allow muscle groups to work towards failure	Use springs to assist patients until enough muscle strength improves
Spring	3-4 Yellow springs adequate	4-6 Yellow springs increases levels of low resistance or assistance
Sequence	Any combination acceptable.	Working disabled area first will eliminate excessive fatigue preventing the patient from working target areas.
Frequency	3-4 sessions per week	1-2 sessions per week
Stability	Promote instability, increases fitness gains	Avoid instability, allows patient to focus on target muscles

## References

1. Gehlsen GM, Grigsby SA, Winant DM. Effects of an Aquatic Fitness Program on the Muscular Strength and Endurance of Patients with Multiple Sclerosis. *Physical Therapy*. 1984; 64(5): 653-7.
2. Taylor NF, Dodd KJ, Damiano DL. Progressive Resistance Exercise in Physical Therapy: A Summary of Systemic Reviews. *Physical Therapy*. 2005; 85(11): 1208-23.
3. Darrah J, Fan JS, Chen LC, et al. Review of the effects of progressive resisted muscle strengthening in children with cerebral palsy: a clinical consensus exercise. *Pediatric Physical Therapy*. 1997; 9: 12-7.
4. Dodd KJ, Taylor NF, Damiano DL. A systemic review of the effectiveness of strengthening for people with cerebral palsy. *Arch Phys Med Rehabil*. 2002; 83:1157-64.
5. MacPhail H, Kramer J. Effect of isokinetic strength on functional ability and walking efficiency in adolescents with cerebral palsy. *Dev Med Child Neurol*. 1995; 37: 763-775.
6. Tweedy S. Evaluation of Strength and Flexibility Training for Adolescent Athletes With Cerebral Palsy: Full Report. Belconnen, Australia: Australian Sports Commission; 1997.
7. Unutzer J, Patrick D, Marmom T, Simon G, Katon W. Depressive Symptoms and mortality in a prospective study of 2,558 older adults. *Am J Geriatr Psychiatry*. 2002; 10(5): 521-30.
8. Glassman A, Shapiro P. Depression and the course of coronary artery disease. *Am J Psychiatry*. 1998; 155: 4-11.
9. De Groot M, Anderson R, Freedland K, Clouse R, Lustman P. Association of depression and diabetes complications: a meta-analysis. *Psychosom Med*. 2001; 63: 619-30.
10. Gordon JL, Ditto B, Lavoie KL, Pelletier R, et al. The effect of major depression on postexercise cardiovascular recovery. *Psychophysiology*. 2011; 48(11): 1605-10.

**Author Information**

**Mark Greenberg, M.D.**

Director Pediatric Anesthesiology Professor of Anesthesiology and Pediatrics, University of California, San Diego  
San Diego, CA, USA

**Alyssa Brzenski, M.D.**

Associate Professor of Anesthesiology, University of California, San Diego  
San Diego, CA, USA