

Relationship of Hand wrist and panoramic radiographs

B Rai, S Anand

Citation

B Rai, S Anand. *Relationship of Hand wrist and panoramic radiographs*. The Internet Journal of Forensic Science. 2007 Volume 3 Number 1.

Abstract

The purpose of this study was to investigate the relationship between the stage of calcification of various teeth and skeletal maturity stage among Indians individuals. The study subjects consisted of 23 male subjects and 22 female subjects ranging in age from 8 to 20 years. A total of 85 hand-wrist, panoramic radiographs were obtained and analyzed. Skeletal age and skeletal maturity stages were determined from hand-wrist radiographs by using the method outlined in the atlas of Fishman's system. The tooth development of mandibular canines, first and second premolars and second molars were assessed according to the Demirjian system. The second molar was the tooth showing the highest correlation. This suggested that tooth calcification stages from panoramic radiographs might be clinically useful as a maturity indicator after pubertal growth period.

INTRODUCTION

Hand wrist radiographs have been used for calculation of skeletal age for many years. The ossification and development of the carpal bones of the wrist, the metacarpals of the hands, and the phalanges of the fingers form a chronology of skeletal development. A satisfactory hand wrist radiograph and a dental x-ray or the cephalometric x-ray source. In use, the overall pattern observed in the hand-wrist film is compared with age standards in a reference atlas¹ to obtain a skeletal age for the patient. In addition, the status of certain specific landmarks such as the ulnar sesamoid or hamate bones can be used to obtain an estimate of the timing of the adolescent growth spurt.² Various areas of the skeleton have been used: the foot, the ankle, the hip, the elbow, the hand wrist, and the cervical vertebrae.³

MATERIALS AND METHODS

The study sample consisted of 45 patients (M:F :: 23:22) selected from Department of Orthodontics, Government Dental College and Hospital associated with Pt. B.D. Sharma Postgraduate Institute of Medical Science, Rohtak (Haryana). Hand wrist panoramic radiographs and cephalogram of 45 patients were taken.

Selection criteria included:

1. The subject had no previous history of trauma or injury to the face and hand and wrist region.
2. The subjects had normal dental conditions.

3. The subjects had undergone neither previous orthodontic treatment nor extraction of any permanent teeth.

Assessment of parameters:

A. To evaluate the stage of skeletal maturation of each Hand-wrist radiograph according to the method described by Fishman.⁵

The following selected ossification events were determined:

MP₃ : The middle phalanx of the third finger, the epiphysis equals its diaphysis.

S Stage : The first mineralization of the ulnar sesamoid bone.

MP_{3Cap} : The middle phalanx of the third finger, the epiphysis caps its diaphysis.

MP_{5Cap} : Capping of epiphysis seen in middle phalanx of fifth finger.

DP_{3u} : The distal phalanx of third finger, complete epiphyseal union.

MP_{3u} : The middle phalanx of third finger, complete epiphyseal union.

Tooth calcification from panoramic radiographs was rated according to the method described by Demirjian et al.⁶ In which 8 stages of calcification.

Left mandibular teeth in panoramic radiographs were examined. In case of any missing left mandibular teeth, the right teeth corresponding to the missing teeth were substituted. Mandibular incisor and first molars were not rated because apical closure had already take place.

All of the assessments were made simultaneously on an illuminated viewing box in dark room. The interpretations of hand wrist, panorgamic and cephalometric radiographs were discussed until agreement was reached. Exact chronological ages were verified by reference to the patient's birth date. The entire data collected was subjected to statistical an analysis by using SPSS (SPSS Inc., Chicago, Illinois).

RESULTS

The sample consisted of 45 patients (M:F :: 23:22) subjects ranging in age from 8 years to 20 years.

Figure 1

Table 1 : Mean and standard deviation of chronological and skeletal age by Fishman method (Hand wrist radiographs) and by Hassel and Farman method (Cephalogram).

Skeletal maturity stages		Chronological age (in years)		According to Fishman method (in years)	
		Mean + S.D.		Mean + S.D.	
Fishman	Hassel	Male	Female	Male	Female
MP ₃	Stage 1	11.4±1.2	10.1±1.3	11.68±1.06	10.58±.88
S	Stage 2	11.7±1.4	10.9±1.2	12.33±1.09	11.22±1.11
MP _{5Op}	Stage 3	13.5±1.5	11.8±1.5	13.75±1.06	12.06±0.96
MP _{3Op}	Stage 4	14.2±1.4	12.1±1.4	14.38±1.08	12.34±0.90
DP _{3u}	Stage 5	14.9±1.3	13.6±1.2	15.11±1.03	13.10±0.87
MP _{3u}	Stage 6	15.8±1.2	14.5±1.2	16.40±1.02	14.77±0.96

Figure 2

Table 2 : Percentage distribution of calcification stages of canine at MP, S, MP MP, DP, and MP stage

Stages	MP ₃		S		MP _{5Op}		MP _{3Op}		DP _{3u}		MP _{3u}	
	F	M	F	M	F	M	F	M	F	M	F	M
D	2.2	-	-	-	-	-	-	-	-	-	-	-
E	14.6	5.2	5.1	4.3	2.3	1.3	-	-	-	-	-	-
F	65.2	51.1	47.7	32.4	25.6	4.2	5.6	1.2	4.8	0	1.8	0
G	12.5	20.9	30.4	27.3	24.4	14.9	12.8	10.5	18.8	11.8	4.8	6.8
H	7.5	17.9	17.1	28.0	47.3	75.8	70.0	86.5	76.4	88.2	94.5	93.5
I	1.0	4.7	1.0	8.0	1.4	2.8	10.6	1.8	0	0	2.5	0

The canine stage H in male and female at MP_{3u} showed highest percent distribution.

Figure 3

Table 3 : Percentage distribution of calcification stages of second molar at MP, S, MP MP, DP, and MP stage

Stages	MP ₃		S		MP _{5Op}		MP _{3Op}		DP _{3u}		MP _{3u}	
	F	M	F	M	F	M	F	M	F	M	F	M
D	25.3	21.2	10.3	8.3	3.4	2.1	1.4	0	0	0	0	0
E	53.5	49.5	52.5	49.2	27.1	26.9	29.1	30.1	7.6	6.3	2.8	0
F	11.7	11.8	23.8	21.3	38.5	37.9	48.5	58.2	21.5	39.5	52.5	51.7
G	9.5	10.8	15.4	14.9	21.5	20.9	12.5	11.3	55.8	59.3	44.7	48.3
H	0	0	0	5.9	10.2	9.8	9.2	0	13.8	0	0	0
I	0	0	0	-	0	0	0	0	2.3	0	0	0

Second molar stage G in males and females at DP_{3u} showed highest percent distribution of calcification.

Figure 4

Table 4 : Percentage distribution of calcification stages of first pre-molar at MP, S, MP MP, DP, and MP stage

Stages	MP ₃		S		MP _{5Op}		MP _{3Op}		DP _{3u}		MP _{3u}	
	F	M	F	M	F	M	F	M	F	M	F	M
D	4.1	0	2.7	0	1.3	0	0	0	0	0	0	0
E	44.4	17.3	26.5	6.5	30.5	9.1	12.6	9.3	1.2	0	0	0
F	33.5	43.7	38.3	17.5	42.4	9.1	27.1	9.3	3.9	0	1.3	0
G	12.4	21.6	11.4	49.7	15.4	81.8	29.4	81.5	5.3	2.4	8.1	0
H	5.1	17.4	9.1	26.3	4.1	0	32.7	0	84.6	97.6	85.6	100
I	0	0	0	0	0	0	0	0	0	0	0	0

The first molar stage H in males and females at MP_{3u} showed highest percentage distribution of calcification.

Figure 5

Table 5 : Percentage distribution of calcification stages of second pre-molars at MP, S, MP MP, DP, and MP stage

Stages	MP ₃		S		MP ₃ Op		MP ₃ Op		DP _{3u}		MP _{3u}	
	F	M	F	M	F	M	F	M	F	M	F	M
D	0	0	3.7	0	0	0	0	0	0	0	0	0
E	53.3	26.5	23.9	5.3	17.5	0	18.1	0	0	0	1.2	0
F	31.2	47.6	55.2	38.0	35.1	12.0	34.3	11.2	21.5	0	2.1	21.3
G	15.5	16.7	11.5	44.7	27.8	44.3	29.1	45.1	44.2	11.3	15.5	75.4
H	0	9.1	5.7	11.0	19.6	43.7	17.4	43.7	33.3	88.7	81.3	3.3
I	0	0	0	0	0	0	0	0	0	0	0	0

Second premolar stage H in males at DP_{3u} and females at MP_{3u} showed highest percent distribution of calcification.

Figure 6

Table 6 : Correlation coefficient between skeletal and dental development stages in male and female subjects.

Tooth	Correlation coefficient (r value)	
	Males	Females
Canine	0.58	0.52
First premolar	0.69	0.58
Second premolar	0.42	0.43
Second molar	0.73	0.69

p<0.01.

Highest correlation coefficient showed by second premolar in male and female.

DISCUSSION

From the current study, maturation patterns of tooth development have shown that males tend to be more advanced as compared with females in relation to skeletal maturity stages as previous study. ⁷ The correlation coefficient between skeletal maturity and calcification stages of the teeth, were quite high, ranging from 0.42 to 0.73 and were statistically significantly significant (p<0.01). Chartkow et al. ⁸ and Coutinho et al. ⁹ have suggested a high relationship between calcification of mandibular canine and skeletal maturity indicators while in this study mandibular second molar in male as well as female. By the time the skeletal age assessment was performed, hand wrist radiographs and cephalograms from male subjects clearly differed from the standard plates more frequently than those of the female subjects, particularly in carpal bone area, which always showed less maturity compared with the other bones as by Acheson et al. ¹⁰ and Carpenter and Lester. ¹¹

The skeletal correlation with orthopedic appliances such as frankel appliance, Herbst appliance and cervical pull head could be initiated during the accelerating growth period and continued through the peak velocity period, depending on the severity of skeletal malrelationship.

From the present study, the relationship between the tooth calcification stages and the skeletal maturity indicators probably allows the clinician to more easily identify the stages the pubertal growth period from the panoramic radiographs. The second molar was the tooth showing the highest correlation.

CONCLUSION

Second molar was the tooth showing the highest correlation. If a strong association exist between skeletal maturity and dental calcification stages, the stages of dental calcification might be used as a first level diagnostic tool to estimate the timing of the pubertal growth spurt. The ease of recognizing dental developmental stage, together with availability of intraoral or panoramic radiographs in most orthodontic or pediatric dental practice, are practical reasons for attempting to assess physiologic maturity without resorting to hand-wrist radiographs. Hence further study recommended in large sample size and further study should be address development of second molars.

CORRESPONDENCE TO

Dr. Balwant Rai S/o Sh. Ramsawroop Village – Bhangu Distt. Sirsa Post Office – Sahuwala I. E-mail : drbalwantraissct@rediffmail.com Mobile No. : 091-9812185855

References

1. Greulich WW, Pyle SL : Radiographic atlas of skeletal development of the hand and wrist, 2nd, Palo Alto, Calif; 1959, Stanford University Press.
2. Grave KC, Brown T : Skeletal ossification and the adolescent growth spurt. Am J Orthod. 1976; 69 : 611-19.
3. Hassel B, Farman AG. Skeletal maturation evaluation using cervical vertebrae. Am J Orthod Dent Ofac Orthop 1995; 107 : 58-66.
4. Anderson DL, Thompson GW, Popovich F. Interrelationship of dental maturity, skeletal maturity, height and weight from age to 14 years. Growth Am J Orthod 1975; 39 : 453-462.
5. Fishman LS. Radiographic evaluation of skeletal maturation. Angle Orthod 1982; 52 : 88-112.
6. Demirjian A, Goldstein H, Tanner J.M. A new system of dental age assessment. Hum Biol 1973; 45 : 211-227.
7. Chertkow S. Tooth mineralization as an indication of the pubertal growth spurt. Am J Orthod 1980; 77 : 79-91.
8. Chertkow S, Fatti P. The relationship between tooth mineralization and early evidence of the ulnar sesamoid. Angle Orthod 1979; 49 : 282-288.
9. Coutinho S, Buschang PH, Miranda F. Relationship

between mandibular canine calcification stages and skeletal maturity. Am J Orthod 1993; 104 : 262-268.

10. Achenon RM, Vicinus JH, Fowler GB. Studies in the reliability of assessing skeletal maturity from x-rays. Part III.

Greulich-Pyle atlas and Tanner - Whitehouse Method contrasted. Hum Biol. 1966; 38 : 204-218.

11. Carpenter CT, Lester EL. Skeletal age determination in young children : analysis of three regions of hand wrist film. J Pediatrorthop 1993; 13 : 76-79.

Author Information

Balwant Rai, B.D.S.

Government Dental College, Pt. Bhagwat Dayal Sharma, Post Graduate Institute of Medical Science

S.C. Anand, (M.D.S.) Oral & Maxillofacial Surgery and Orthodontics

Government Dental College, Pt. Bhagwat Dayal Sharma, Post Graduate Institute of Medical Science