Large Gastric Perforation In A Neonate

G Parmar, M Khoradiya, B Ambasana

Citation

G Parmar, M Khoradiya, B Ambasana. *Large Gastric Perforation In A Neonate*. The Internet Journal of Surgery. 2009 Volume 24 Number 2.

Abstract

Large gastric perforation in a neonate is very rare condition. We are reporting such a condition in a two-day-old male child. This condition requires urgent surgical intervention. We did emergency laparotomy with suturing of large gastric perforation in two layers.

INTRODUCTION

Large spontanous gastric perforation is a very rare condition with only about 200 cases reported in the literature (1). The condition entails very high mortality. Its pathogenesis is greatly debated.

Congenital absence of gastric wall musculature (2), stress ulceration secondary to neurogenic difficulties (3), and ischaemia of the gastric wall secondary to vascular shunting (4) have been proposed as aetiological factors.

We report a case of large gastric perforation in a two-day-old child seen by us.

CASE REPORT

A two-day-old child was admitted in the paediatric ward with complaints of dyspnoea and abdominal distention. There was no history of vomiting and the baby had passed meconium. He had had a full-term normal vaginal delivery with a birth weight of 2.75kg. There was birth asphyxia when he was born.

After 5-6 hours, the patient developed complaints of abdominal distention. When he came to hospital, he was in respiratory failure and in shock with a respiratory rate of 52 per minute and a pulse rate of 160 per minute. Immediate endotracheal intubation was done and he was kept on ventilatory support and management of shock with vasopressors and plasma expanders. The abdomen was markedly distended with absent bowel sounds and liver dullness was masked. The rest of the examination was normal.

On investigation, haemoglobin was 15 g/dl and TLC 12000/cumm (P75, L20, M3, E2). Abdominal x-ray

(standing) showed free gas under both domes of the diaphragm. The viscera were displaced medially producing the typical "saddle bag" appearance (fig.1). Chest x-ray, serum creatinine, blood urea and serum electrolytes were within normal limits.

Figure 1

Fig. 1: Abdominal x-ray (erect) showing free air under the diaphragm. The abdominal viscera are displaced medially producing the "saddlebag" appearance.



Urgent exploratory laparotomy was performed by upper transverse incision. This revealed a large gastric perforation of about 3x1cm with ragged edges on the anterior wall. The surrounding stomach wall was ischemic (fig. 2).

Figure 2

Fig. 2: Large gastric perforation in the anterior wall of the stomach.



Excision of the ischemic edges was done and the perforation was closed with silk 3-0 in two layers.

Postoperatively, the patient was kept on ventilatory support and management of shock with vasopressors and plasma expanders. Nasogastric aspiration was done two-hourly. Unfortunately, we lost our patient on the 8th postoperative day due to respiratory failure and shock.

DISCUSSION

Spontaneous gastric perforation as reported earlier is more common in the preterm baby, most commonly occurring between the 2^{nd} and 7^{th} day (5,6). The highest reported incidence of rupture is on the 3^{rd} day of life (6). Our baby was a full term baby and the perforation had occurred on the 2^{nd} day of life. Sudden abdominal distension and respiratory distress have been reported as predominant symptoms (1,5). Those commonly observed include prematurity, respiratory distress at birth, asphyxia and resuscitation, premature rupture of membranes and breech, cesarean or twin delivery, in decreasing order of frequency.

Most commonly the perforations have been linear tears seen on the greater curvature, usually high and measuring between 0.5 and 8cm (1,5). In the present case the perforation was about 3x1 cm in size. It was located on the anterior wall of the stomach.

Prompt surgical intervention with repair of gastric tear is the recommended management. Any delay in surgery will result in a higher mortality (8,9).

The pathogenesis of spontaneous gastric perforation is much debated (2-4,10-12). Gastric tissue ischemia secondary to hypoxia is one plausible explanation (4,11). During severe hypoxic stress there is selective shunting of blood away from the splanchnic vascular bed leading to vascular damage and perforation. This was the probable sequence of events that operated in our patient.

References

1. Houck WS, Griffin JA: Spontaneous linear tears of the stomach in the newborn infant. Ann Surg; 1981; 193: 763-767.

2. Herbut PA: Congenital defect in the musculature of the stomach with rupture in a newborn infant. Arch Patho1; 1943; 36: 91-94.

3. Kiesewetter WB: Spontaneous rupture of the stomach in the newborn. Am J Dis Child; 1956; 91: 162-167.

4. Touloukian RJ: Gastric ischemia: The primary factor in neonatal perforation. Clin Pediatr; 1973; 12: 219-225.5. Holgersen LO: The aetiology of spontaneous gastric

perforation of newborn: A re-evaluation. J Pediatr Surg; 1981; 16: 608-613.

6. Hood JH: Clinical consideration of intestinal gas. Ann Surg; 1966; 163: 359-366.

7. Prabhakar G, Agrawal LD, Shukla A, et al.: Spontaneous gastointestinal perforation in neonate. Indian Pediatr; 1991; 28: 1277-1280.

8. Saracli J, Mann M, Frend DM, et al.: Rupture of the stomach in the newborn infant: Report of three cases and review of the world literature. Clin Pediatr; 1967; 6: 583-588.

9. Inouye WY, Evans G: Neonatal gastric perforation. Arch Surg; 1964; 88: 471-485.

10. Kneiszi F: Some data on the etiology of gastric rupture in the newborn, Biol Neonate; 1962; 4: 201-222.

11. Lloyd JR: The etiology of gastrointestinal perforations in the newborn. J Pediatr Surg; 1969; 4: 77-84.

12. Nagaraj HS, Sandhu AS, Cook LN, Buchino JJ, Groff DB: Gastrointestinal perforation following indomethacin therapy in very low birth weight infants. J Pediatr Surg; 1981, 16: 1003-1007.

Author Information

Girish Parmar, M.B.B.S.,M.S.

Assistant Professor,, Department of Surgery,, Shri M.P.Shah Medical College, Jamnagar-361 001, Gujarat, India.

Mahesh Khoradiya, M.B.B.S.

Junior Resident,, Department of Surgery,, Shri M.P.Shah Medical College, Jamnagar-361 001, Gujarat, India.

B.C. Ambasana, M.B.B.S., M.S., PGDHHM, DNHE, CDM

Professor of Surgery,, Department of Surgery,, Shri M.P.Shah Medical College, Jamnagar-361 001, Gujarat, India.