

Phytochemical and Biological Investigations of *Sida Rhomboidea* Linn.

A Chowdhury, M Ashraful Alam, M Rahman, M Rashid

Citation

A Chowdhury, M Ashraful Alam, M Rahman, M Rashid. *Phytochemical and Biological Investigations of Sida Rhomboidea Linn.*. The Internet Journal of Alternative Medicine. 2008 Volume 7 Number 2.

Abstract

Daucosterol (1) was isolated from the n-hexane soluble fraction of a methanol extract of the stems of *Sida rhombifolia* Linn. (Syn. *S. rhomboidea* Linn.). The n-hexane, carbon tetrachloride and dichloromethane soluble fractions of this methanol extract were subjected to antimicrobial and antioxidant screening as well as brine shrimp lethality bioassay. The dichloromethane and carbon tetrachloride soluble fractions showed moderate inhibitory activity to microbial growth while the n-hexane fraction showed highest cytotoxicity having LC_{50} 0.84 μ g/ml. The dichloromethane soluble fraction also revealed potent antioxidant activity with IC_{50} 20 μ g/ml.

INTRODUCTION

Sida rhombifolia Linn. (Syn. *S. rhomboidea* Linn., Family-Malvaceae, Bengali Name-Lal Berela) is a small erect undershrub, which grows all over Bangladesh ¹. The stems are used to treat rheumatism and tuberculosis ². It is also reputed as anthelmintic and sedative. The roots are used in epilepsy, neurological problem, and fever and heart diseases ³. Fruits are as known to be antibacterial and antifungal. The whole plant in combination with other drugs is prescribed as an antidote to snake and scorpion venom ⁴. Previous phytochemical investigations of *Sida* species resulted in the isolation of stigmasterol, stigmatenol and β -sitosterol. We, herein, report the preliminary antimicrobial, cytotoxic and antioxidant activities of the organic extractives as well as isolation of daucosterol from the n-hexane soluble fraction of *S. rhomboidea* for the first time.

MATERIALS AND METHODS

GENERAL EXPERIMENTAL PROCEDURE

The ¹H NMR spectra was recorded using a Bruker AMX-400 (400-MHz) instrument. For NMR studies deuterated chloroform was used and the values for ¹H spectrum was referenced to the residual nondeuterated solvent signal.

PLANT MATERIAL

Stems of *S. rhomboidea* were collected from Azimpur Govt. Quarter Area, Dhaka in the month of September 2007. A voucher specimen for this collection has been deposited in

Bangladesh National Herbarium, Dhaka (accession no. -32897).

EXTRACTION AND ISOLATION

The powdered stems (600 g) of *S. rhomboidea* was soaked in 1.5 L methanol for 7 days with occasional shaking and stirring and filtered through a cotton plug followed by Whatman filter paper number-1. The extract was then concentrated by using a rotary evaporator at reduced temperature and pressure. A portion (5 g) of the concentrated methanol extract was fractionated by the modified Kupchan partitioning method ⁵ into n-hexane (1.0 g), carbon tetrachloride (0.7 g), dichloromethane (0.75 g) and aqueous soluble materials (2.0 g). The n-hexane soluble material was fractionated by column chromatography (CC) over silica gel (60-120 mesh) using n-hexane, ethyl acetate and methanol mixtures of increasing polarities to give 30 fractions each 5 ml. Preparative thin layer chromatography (stationary phase-silica gel F₂₅₄, mobile phase 20% ethyl acetate in toluene, thickness of plates 0.5 mm) of fraction n-hexane eluted with 20-30% ethyl acetate in n-hexane afforded compound 1.

BIOASSAYS

The antimicrobial activity of the extractives was determined by the disc diffusion method ⁶. The samples were dissolved separately in chloroform or methanol and applied to sterile filter paper discs at a concentration of 400 μ g/disc. Kanamycin disc (30 μ g/disc) was used as standard.

For cytotoxicity assay DMSO solutions of the plant extracts


were applied against *Artemia salina* in a 1-day in vivo assay, the experimental details of which could be found elsewhere ⁷. For the experiment 4 mg of each of the Kupchan fractions was dissolved in DMSO and test solutions of varying concentrations such as 400, 200, 100, 50, 25, 12.50, 6.25, 3.125, 1.563 and 0.781 µg/ml were obtained by serial dilution technique. The median lethal concentration LC₅₀ of the test samples after 24 hours of exposure was obtained by a plot of percentage of the shrimps killed against logarithm of the sample concentration. Here vincristine sulphate was used as the standard.

The free radical scavenging activity of the extractives on the stable radical 1, 1-diphenyl-2-picrylhydrazyl (DPPH) were estimated by the method of Brand-Willimams ⁸. Here 2.0 ml of a methanol solution of the samples at different concentration were mixed with 3.0 ml of DPPH methanol solution (20µg/ml). The antioxidant potential was assayed from the bleaching of purple colored methanol solution of DPPH radical by the plant extract as compared to that produced by the standard antioxidant agents of tert-butyl hydroxytoluene (BHT) and ascorbic acid (ASA) by UV spectrophotometer. Free radical scavenging activities of different partitionates of methanolic extract and standard tested by DPPH method. IC₅₀ value of the standard BHT obtained 10 µg/ml.

RESULTS AND DISCUSSION

Compound 1 was isolated from the n-hexane soluble fraction of methanolic extract of the stems of *S. rhomboidea* by repeated chromatographic separation and purification over silica gel. The structure of the isolated compound was determined by comparison of ¹H NMR data with previously reported values ⁹, as well as co- TLC with authentic sample. Although it has previously been reported from many plants, this is the first report of its isolation from *S. rhomboidea*.

Figure 1



Daucosterol (1)

In the antimicrobial screening, the extractives as *S. rhomboidea* exhibited mild to moderate antimicrobial activity. The zone of inhibition produced by the carbon tetrachloride and dichloromethane soluble fractions of methanolic extract ranged from 8-13 and 8-12 mm respectively (Table -1). The n-hexane and aqueous soluble fractions did not show inhibit the growth of microorganisms.

Figure 2

Table 1. Antimicrobial activity of extractives at 400 µg/disc.

Test microorganisms	Diameter of zone of inhibition (mm)		
	CT	DCM	KAN
Gram positive bacteria			
<i>Bacillus cereus</i>	12	10	35
<i>B. megaterium</i>	9	10	36
<i>B. subtilis</i>	9	8	34
<i>Staphylococcus aureus</i>	8	10	34
<i>Sarcina lutea</i>	11	9	35
Gram negative bacteria			
<i>Escherichia coli</i>	10	8	36
<i>Pseudomonas aeruginosa</i>	8	9	36
<i>Salmonella paratyphi</i>	11	12	35
<i>S. typhi</i>	13	9	35
<i>Shigella dysenteriae</i>	8	10	32
<i>S. boydii</i>	8	9	33
<i>Vibrio mimicus</i>	10	9	37
<i>V. parahemolipicus</i>	11	8	31
Fungi			
<i>Candida albicans</i>	10	10	32
<i>Aspergillus niger</i>	10	10	32
<i>Sacharomyces cerevacae</i>	10	9	31

CT: carbon tetrachloride soluble fraction of the methanolic extract; DCM: dichloromethane soluble fraction of the methanolic extract; KAN: standard kanamycin disc (30µg/disc).

Following the procedure of Meyer ⁷, the lethality n-hexane (HX), carbon tetrachloride (CT), dichloromethane (DCM)

and aqueous (AQ) soluble fractions of the methanolic extract to brine shrimp was determined on. Table 2 shows the results of the brine shrimp lethality testing after 24 hours of exposure to the samples and the positive control, vincristine sulphate. The LC_{50} obtained from the best fit line slope were found to be 0.84, 5.89, 7.6 and 2.7 $\mu\text{g/ml}$ for n-hexane, carbon tetrachloride, dichloromethane and aqueous soluble materials, respectively (Table -2). In comparison with the positive control (vincristine sulphate, LC_{50} 0.46 $\mu\text{g/ml}$) the cytotoxicity exhibited by the n-hexane, methanolic aqueous, carbon tetrachloride fraction of methanolic extract was significant.

Figure 3

Table 2. LC data of test samples of

Sample	LC_{50} ($\mu\text{g/ml}$)
VS	0.46
HX	0.84
CT	5.8
DCM	7.6
AQ	2.7

VS: vincristine sulphate (std); HX: n-hexane soluble fraction of methanolic extract; CT: carbon tetrachloride soluble fraction of methanolic extract; DCM: dichloromethane soluble fraction of methanolic extract; AQ: aqueous soluble fraction of methanolic extract.

The antioxidant activity of various fractions of the methanolic extract was also determined. Table-3 shows antioxidant activity of the test samples. Here, tert butyl-1-hydroxy toluene (BHT) was used as positive control. The IC_{50} values for n-hexane, carbon tetrachloride, dichloromethane and methanolic aqueous were found to be 316, 217, 20 and 32 $\mu\text{g/ml}$, respectively. In comparison with positive control (BHT) the antioxidant activity exhibited by the dichloromethane and aqueous soluble fractions of methanolic extract was significant. The pure compound 1 was also tested for antioxidant activity, but no significant inhibitory capacity was found (data not shown in table-3).

Figure 4

Table 3. IC data of test samples of

Sample	IC_{50} ($\mu\text{g/ml}$)
BHT	9
HX	316
CT	217
DCM	20
AQ	32

BHT: tert butyl-1-hydroxy toluene (Std); HX: n-hexane soluble fraction of methanolic extract; CT: carbon tetrachloride soluble fraction of methanolic extract; DCM: dichloromethane soluble fraction of methanolic extract; AQ: aqueous soluble fraction of methanolic extract.

The results of antimicrobial, cytotoxic and antioxidant activities support the folk uses of *S. rhomboidea* in various diseases.

ACKNOWLEDGEMENT

The authors wish to thank Centre for Biomedical Research, University of Dhaka, Dhaka-1000, Bangladesh for providing some laboratory facilities for carrying out the research.

References

1. Rahman, M. M., Khan, M. Salar and Hossain, S. Flora of Bangladesh, Malvaceae – Ministry of Environment and Forest Govt. of the People Republic of Bangladesh, Pp.47, 2003.
2. Islam, M.E., Haque M.E. and Mosaddik, M.A. Cytotoxicity and Antibacterial Activity of *Sida rhombifolia* (Malvaceae) Grown in Bangladesh. Phytotherapy Research, Phytother. Res. 17: Pp.973-975, 2003.
3. Choudhuri, A. B. Forest plants of Eastern India. Publishing house 8/81 Punjabi Bagh, New Delhi- 110026. Pp.136, 1993.
4. Kirtikar, K.R. and Basu, B.D. Indian medicinal plants, B. Singh and M.P. Sing publishers, India, Vol.1, 2nd ed. Pp.310, 1980.
5. Vanwagenen, B.C., Larsen, R., Cardellina, J.H., Ranzazzo, D., Lidert, Z.C. and Swithenbank, C. Ulosantoin, a potent insecticide from the sponge *Ulosa ruetzleri*. J. Org. Chem. 58: Pp.335-337, 1993.
6. Bauer, A.W., Kirby, W.M.M., Sherris, J.C. and Truck, M. Antibiotic susceptibility testing by standard single disc diffusion method. Am. J. Clin. Pathol. 45: Pp.426-493, 1966.
7. Meyer, B.N., Ferringni, N.R., Puam, J.E., Lacobsen, L.B., Nichols, D.E. and McLaughlin, J.L. Brine shrimp: a convenient general bioassay for active constituents. Planta Med. 45: Pp.31-32, 1982.
8. Choi HY, Jhun EJ, Lim BO, Chung IM, Kyung SH, Park DK. Application of flow injection-chemiluminescence to the study of radical scavenging activity in plants. Phytother. Res. 14: Pp.250-253, 2000.
9. Moghddam F. M., Farimani M.M., Salahvarzi S. and Amin G., eCAM Advance Access published September 26: Pp.1, 2006.

10. Dictionary of Natural Products, Published by Chapman and Hall, 2002.

Author Information

Akhtaruzaman Chowdhury, M.Sc.

Department of Chemistry Rajshahi University of Engineering and Technology Rajshahi Bangladesh

Md. Ashraful Alam, Ph.D.

Department of Chemistry Rajshahi University of Engineering and Technology Rajshahi Bangladesh

Mohammad Sharifur Rahman, M.Pharm

Department of Pharmaceutical Chemistry Faculty of Pharmacy University of Dhaka Bangladesh

Mohammad Abdur Rashid, Ph.D

Department of Pharmaceutical Chemistry Faculty of Pharmacy University of Dhaka Bangladesh