

# Case of successful Extracorporeal Life Support (ECLS) in a patient with refractory asthma – Importance of increasing awareness of the role of ECLS in the UK

A Tewari, T Ali, R Baba

---

## Citation

A Tewari, T Ali, R Baba. *Case of successful Extracorporeal Life Support (ECLS) in a patient with refractory asthma – Importance of increasing awareness of the role of ECLS in the UK*. The Internet Journal of Healthcare Administration. 2008 Volume 6 Number 2.

## Abstract

Life threatening asthma is acute, severe and often unexpected. 1 Ventilation is not end stage management. Extracorporeal Life Support (ECLS) which traditionally has provided significant support for acute respiratory distress in infants and children and recently adults 2 is also enabling improved prognosis in those patients with severe refractory asthma .3 We describe the use of ECLS in a patient with acute severe asthma where despite all aggressive medical management and mechanical ventilation, a fatal outcome appeared likely. Following initiation of ECLS the patient showed a considerable clinical improvement within 24 hours. Thus we highlight the importance of ECLS in patients with severe refractory asthma and ask that clinicians be increasingly aware of this facility in the UK.

## INTRODUCTION

In the UK asthma is an important cause of impaired quality of life, hospital admission and mortality across the whole spectrum of age, sex, ethnicity, socioeconomic status and geographical location.<sup>4</sup> The prevalence of asthma is increasing <sup>5</sup> and in the UK it currently affects 5.1 million adults (approximately 13%).<sup>6</sup> Interestingly the number of deaths attributed to asthma is decreasing; data on asthma deaths in the UK show that on average 1,400 people die from asthma each year, which equates to almost four people every day.<sup>6</sup>

Acute asthma is characterized by spasm of bronchial smooth muscle, mucosal oedema and increased bronchial mucous secretion.<sup>7</sup> Treatment options include inhaled bronchodilators for rapid control of bronchospasm, theophylline to relax smooth muscle by increasing intracellular cAMP, corticosteroids to decrease bronchial inflammation and magnesium which inhibits calcium influx by blocking voltage-dependent calcium channels thus relaxing smooth muscle airways.<sup>8</sup> Endotracheal intubation and positive pressure ventilation is reserved for patients with severe hypoxaemia or hypoventilation despite these measures.

We describe the use of Extracorporeal Life Support (ECLS)

and positive outcome in a patient with severe asthma that was refractory to mechanical ventilation.

## CASE REPORT

A 20 year old male with a medical background of bronchial asthma and no previous ITU admissions presented to accident and emergency with shortness of breath and wheeze over the last 12 hours. He showed no improvement with his regular medication (becotide, salbutamol). 2 weeks prior to admission he had developed a sore throat and cough productive of sputum.

On initial examination his GCS 15 but he was tachypnoeic with a respiratory rate of 28 breaths/minute, blood pressure was 128/90mmHg and pulse 85 beats per minute. He was talking in broken sentences and there were bilateral inspiratory crackles to midzone and bilateral expiratory wheezes on auscultation. He was treated for acute infective exacerbation of asthma with nebulised bronchodilators, supplemental oxygen, intravenous magnesium sulphate and methylprednisolone. He was also started on tazocin 4.5g three times a day and gentamicin 300mg once a day. However, his clinical condition worsened and pulse oximetry showed SpO<sub>2</sub> of 95% on 15L and 84% on room air. The arterial blood gas on air showed a pH of 7.38 pCO<sub>2</sub> 5.00kPa, pO<sub>2</sub> 7.15kPa. ECG showed sinus tachycardia and

initial full blood count, electrolytes and liver function tests were normal. He was breathing with a respiratory rate of 50 breaths/minute and his saturations were deteriorating to 90% on 15 litres oxygen. He was starting to tire and the decision was made for intubation and further management on the intensive care unit.

He was induced with propofol 200mg, fentanyl 100mcg and suxamethonium 100mg and after intubation and ventilation he was maintained on propofol, and alfentanil infusions. At peak pressure of 40cm H<sub>2</sub>O (PEEP at 0cmH<sub>2</sub>O) he has tidal volumes of 200-300mls and autopeep 15cm H<sub>2</sub>O. I:E ratio was 1:4 and respiratory rate 10/minute. However he was proving very difficult to ventilate. Despite addition of a vecuronium infusion peak pressures were still reaching 45cm H<sub>2</sub>O though tidal volumes temporarily increased to 300-350mls/ breath. Arterial blood gas showed a PCO<sub>2</sub> of >20kPa, pH 7.1, pO<sub>2</sub> 12kPa on FiO<sub>2</sub> .80. Despite adequate sedation and paralysis, airflow obstruction and dynamic hyperinflation worsened. Portable chest radiographs demonstrated evidence of a left sided pneumothorax and a chest drain was inserted immediately. 2 hours later the arterial blood gas showed pH of 6.82, pCO<sub>2</sub> still >20kPa, pO<sub>2</sub> 10kPa and BE incalculable. The patient also developed atrial fibrillation with a ventricular rate of 110 beats/minute.

At this stage his case was discussed with a tertiary respiratory centre. The patient was accepted for treatment for acute lung injury via veno-venous extra corporeal life support (ECLS). At this point, peak inspiratory pressures were ranging between 45-50 cm H<sub>2</sub>O, with an auto peep of 12-14cm H<sub>2</sub>O. Prior to cannulation for ECLS a right sided apical chest drain was inserted. ECLS was initiated. Less than 24 hours later, whilst receiving ECLS, the patient was ventilating with a peak pressure of 21cm H<sub>2</sub>O. On an FiO<sub>2</sub> 0.40, he had a pH of 7.43, pCO<sub>2</sub> 5.5kPa, pO<sub>2</sub> 10.8kPa. He was transferred back to our hospital 48 hours later where he was weaned off respiratory support and extubated approximately 36-48 hours later. Recovery and discharge following this was uneventful.

## DISCUSSION

Extracorporeal life support (ECLS) has traditionally been used in infants and children with acute respiratory distress syndrome due to reversible pulmonary disease such as meconium aspiration, idiopathic respiratory distress syndrome, sepsis with pneumonia and persistent pulmonary hypertension of the newborn.<sup>2</sup> It also provides significant support for acute respiratory distress in adults.<sup>9</sup> The first

successful use of ECLS was described by Hill and colleagues in 1972<sup>10</sup> in a 24 year old male who developed post traumatic acute respiratory distress syndrome. By 1986, Gattinoni and colleagues had demonstrated that use of extracorporeal carbon dioxide removal in patients with ARDS was safe and associated with a survival advantage.<sup>11</sup>

The earliest reports of the successful application of ECLS to asthma were described in 1981<sup>13</sup> and 1993.<sup>14</sup> The therapeutic principle in asthma is to ensure adequate oxygenation by partially or totally replacing the lungs' function in gas exchange, giving time for the acute lung injury to heal and also allowing minimization of mechanical ventilation and its risks of O<sub>2</sub> toxicity as well as barotraumas and volutrauma. ECLS is indicated for patients with respiratory failure of a reversible aetiology and who are not responding to standard medical management. Patients in whom ECLS is initiated within five days of mechanical ventilation tend to be most likely to respond whereas those in whom ECLS is begun after 10 days do very poorly.<sup>12</sup>

Our patient had acute severe asthma which did not improve despite maximum medical management and ventilatory support. This patient continued to deteriorate and developed severe dynamic hyperinflation with peak pressures ranging between 45-50mm H<sub>2</sub>O with resultant bilateral pneumothoraces, and severe respiratory acidosis. However following initiation of ECLS as an adjunct to medical and mechanical ventilation, his respiratory function normalised within 24 hours of therapy. ECLS is uniquely suited to the management of severe asthma attacks that are refractory to control with conventional treatment by virtue of the membrane lung's extremely efficient clearance of CO<sub>2</sub>. Rapid normalisation of PCO<sub>2</sub> and correction of respiratory acidosis enables lower airway pressures to be used and lower concentrations of oxygen which helped to minimise oxygen toxicity and effects of volutrauma and barotraumas.<sup>12</sup> In this case the short burst of this treatment helped ensure a rapid recovery from respiratory support and then a steady medical recovery leading to discharge. There is currently no long term data on the statistics for repeat requirement of ECLS for respiratory failure due to refractory asthma.

There are other case reports<sup>131415161718</sup> and one case series<sup>19</sup> in the literature procuring findings that ECLS is a useful method for preventing death in patients with near-fatal status asthmaticus.

The indications for ECLS in the case reports included

cardiovascular compromise and/or cardiac arrest<sup>131517</sup> and severe hypercapnic acidosis.<sup>141618</sup> The indications for ECLS in the 3 patients in the case series<sup>19</sup> included severe, refractory hypercapnia and acidosis, hypoxemia and eventual shock, and dynamic hyperinflation demonstrated by peak inspiratory pressure up to 50–60 cm H<sub>2</sub>O. All patients benefited from 21 to 86 hours of ECLS. In our case the patient was referred for ECLS because of increasing hypoxaemia and hypercapnia during mechanical ventilation, recurrent pneumothoraces and acidosis. He benefited from 24 hours of ECLS.

In their case series, Kukita and colleagues<sup>19</sup> provided an algorithmic approach to mechanical ventilation in patients with status asthmaticus. We ask that clinicians consider ECLS if despite institution of hypoventilation and while tolerating increased peak inspiratory pressure, pH is still less than 7.2, PaCO<sub>2</sub> is greater than 100 mm Hg, or life-threatening conditions of hypoxaemia, hypotension, or barotrauma develop.

In conclusion, this case illustrates the effective use of ECLS for acute, severe, reversible respiratory failure due to status asthmaticus. We ask that clinicians be increasingly aware of this facility in the UK as we believe that ECLS should be considered early in the treatment of patients with status asthmaticus whose gas exchange cannot be satisfactorily maintained by conventional aggressive medical therapy and mechanical ventilation.

## CORRESPONDENCE TO

angela.tewari@yahoo.co.uk, Intensive Care Unit, Mayday Hospital, 503 London Road, Croydon CR7 4YE

## References

- Holgate ST. Pathogenesis of asthma. *Clin Exp Allergy*. 2008 Jun;38(6):872–97. Review.
- Kolla S, Awad SS, Rich PB, Schreiner RJ, Hirschl RB, Bartlett RH. Extracorporeal life support for 100 adult patients with severe respiratory failure. *Ann Surg*. 1997 Oct;226(4):544–64; discussion 565–6
- Mikkelsen M, Meredith EP, et al Emergency Extracorporeal Life Support for Asphyxic Status Asthmaticus Case Reports *Respir Care* 2007; 52 (11): 1525–1529
- Ross et al 50 years of asthma: UK trends from 1955 to 2004 *Thorax* 2007;62;85–90
- Chronic Conditions: A Challenge for the 21st Century,” National Academy on an Aging Society, 2000
- Fact sheet : Trends in asthma mortality in Great Britain Lung and asthma information agency
- Borish L, Culp JA Asthma: a syndrome composed of heterogeneous diseases. *Ann Allergy Asthma Immunol*. 2008 Jul;101(1):1–8; quiz 8–11, 50.
- Skobeloff, EM et al. Intravenous magnesium for the treatment of acute asthma in the emergency department. *JAMA* 1989; 262: 1210–1213
- Schuerer DJ, Kolovos NS, Boyd KV, Coopersmith CM. Extracorporeal membrane oxygenation: current clinical practice, coding, and reimbursement. *Chest*. 2008 Jul;134(1):179–84.
- Hill JD, O'Brien TG, Murray JJ, et al. Prolonged extracorporeal oxygenation for acute post-traumatic respiratory failure (shock-lung syndrome). Use of the Bramson membrane lung. *N Engl J Med* 1972; 286: 629–34.
- Gattinoni L, Pesenti A et al Low frequency positive pressure ventilation with extracorporeal CO<sub>2</sub> removal in severe acute respiratory failure *JAMA* 1986;256:881–85
- Wasserman J, Mazer CD ECMO in adults: what is its role *Canadian Journal of Anaesthesia* 1998 45: 11 p 1044–1048
- MacDonnell KF, Moon HS, Sekar TS, Ahluwalia MP. Extracorporeal membrane oxygenator support in a case of severe status asthmaticus. *Ann Thorac Surg* 1981;31(2):171–175
- Shapiro MB, Kleaveland AC, Bartlett RH. Extracorporeal life support for status asthmaticus. *Chest* 1993;103(6):1651–1654.
- King D, Smales C, Arnold A, Jones OG. Extracorporeal membrane oxygenation as emergency treatment for life threatening acute severe asthma. *Postgrad Med J* 1986;62(731):855–857.
- Tajimi K, Kasai T, Nakatani T, Kobayashi K. Extracorporeal lung assist for patient with hypercapnia due to status asthmaticus. *Intensive Care Med* 1988;14(5):588–589.
- Mabuchi N, Takasu H, Ito S, Yamada T, Arakawa M, Hatta M, Katsuya H. Successful extracorporeal lung assist (ECLA) for a patient with severe asthma and cardiac arrest. *Clin Intensive Care* 1991;2(5):292–294.
- Sakai M, Ohteki H, Doi K, Narita Y. Clinical use of extracorporeal lung assist for a patient in status asthmaticus. *Ann Thorac Surg* 1996;62(3):885–887.
- Kukita I, Okamoto K, Sato T, Shibata Y, Taki K, Kurose M, et al. Emergency extracorporeal life support for patients with near-fatal status asthmaticus. *Am J Emerg Med* 1997;15(6):566–569.

**Author Information**

**A. Tewari**

Specialist Registrar, Mayday Hospital

**T. Ali**

Specialist Registrar, Mayday Hospital

**R. Baba**

Consultant Intensivist, Mayday Hospital