Genital Trichomoniasis Among Women At A University Teaching Hospital: Findings From A Study On 8,443 Clinical Urogenital Samples In Jos, North Central Nigeria.

G Jombo, S Opajobi

Citation

G Jombo, S Opajobi. Genital Trichomoniasis Among Women At A University Teaching Hospital: Findings From A Study On 8,443 Clinical Urogenital Samples In Jos, North Central Nigeria.. The Internet Journal of Parasitic Diseases. 2006 Volume 2 Number 1.

Abstract

Aim: The study was set to ascertain the incidence of Trichomonas vaginalis among women in Jos.

Methods: The study was retrospective in nature. Data generated by the Microbiology laboratory of the Jos University Teaching Hospital (JUTH) on high vaginal swab/endocervical swab (HVS/ECS) samples was compiled for a period of five years (July 2000-June 2005). Samples were collected, transported, stored and processed using standard laboratory procedures. Additional information was obtained from patients' records. Results were analysed using Epi Info 6 statistical software, while P values <0.05 were considered significant.

Results: The incidence of T. vaginalis among women was found to 1.2% in Jos. The peak age of infection was 20-29 years, and there was no significant age difference (P> 0.05). Clinical features associated with T. vaginalis infection include: Nil 6.7%, Itching + Rashes 10.6%, Discharge only 35.6%, Rashes only 8.6%, Itching + Discharge 27.9%, and Itching only 4.8%.

Conclusion:

T. vaginalis is not a rare STD among women in Jos. Proper sex education especially for the youths and adolescents should be intensified so as to ensure a population fully aware of the Dos and Don'ts of STDs and hence reduce the spread and rate of infection with T. vaginalis and other STDs.

INTRODUCTION

Trichomonas vaginalis is a trophozoite; oval in shape as well as flagellated. Five flagella arise near the cytosome; four of these immediately extend outside the cell together, while the fifth flagellum wraps backwards along the surface of the organism_{1,2}. T. vaginalis is an established cause of sexually transmitted diseases (STDs), and currently affects at least170 million people worldwide; more than gonorrhea, syphilis and Chlamydia combined_{3,4,5}. The organism has been reported to cause pneumonia, bronchitis and oral lesions_{6,7}, while a greenish-yellow frothy vaginal secretion and itching are its commonest presentations_{8,9,10,11}. Infertility, premature rupture of membranes, preterm labor, and abortions have all been associated with the organism_{7,8,9}.

T. vaginalis, though generally believed to be a female agent has been detected in between 11% to 17% of men attending

STD clinic in which it was the sole organism detected in the urethra_{12>13}. In North Carolina USA, T. vaginalis was detected in 29% of women attending an STD clinic using polymerase chain reaction (PCR)₁₄; while in Dhaka, Bangladesh, T. vaginalis was detected in 45.5% of the 269 street-based female sex workers₁₅. Also in Cotonou, Benin, T. vaginalis was detected in 10.5% of female commercial sex workers₁₆.

T. vaginalis has been reported to induce cervical intraepithelial neoplasia (CIN) (a premalignant condition) faster than the hitherto strongly associated organisms_{17,18}. In India it was found that, CIN detected by cytology was detected in 7.04%, 4.89% and 4.89% of women infected with T. vaginalis, herpes simplex virus, and human papilloma virus respectively₁₉. T. vaginalis has also been shown to induce epithelial monolayer disruption, creating a microenvironment conducive for HIV-I replication₂₀, a phenomenon which could encourage the spread of HIV infection_{21,22,23,24,25}.

Recent findings have shown a gradual built up of metrinodazole resistant T. vaginalis $_{26,27}$, the drug of choice for its treatment. This has been attributed to a reduction in transcription of the ferrodoxin gene thereby decreasing the ability of the cell to activate the drug₂₈. This has constituted a potential challenge towards the future management of trichomoniasis_{27,28}.

In Nigeria, Nwosu, et al reported T. vaginalis prevalence of 12.5% among females in a cross-sectional study₂₉, while Akerele, et al, in Benin-city₃₀, reported about 15% detection of the organism among pregnant women, Papin, et al from a study on 659 men with urethral discharge reported T. vaginalis to account for 13.8% of the cases across West Africa₃₁.

Among the agents of STDs, T. vaginalis, though fairly researched, appears to attract less enthusiasm among researchers compared to the other agents of $STD_{32,33,34}$. Also knowledge of the depth of the infection in this locality needs to be known_{35,36}. Such information would be a useful guide to the health personnel involved in the management of similar and related patients as well as a good projection of the sequelae that might arise from such ailments, hence the need for this study.

MATERIALS AND METHODS

Setting The study was carried out at the Microbiology laboratory of the Jos University Teaching Hospital (JUTH), Jos Plateau state.

Procedure The study was retrospective in nature. Data generated by the Microbiology laboratory of JUTH endocervical/high vaginal swab (ECS/HVS) samples was compiled for a period of five years (July 2000- June 2005). Samples were collected, transported, stored and processed using standard laboratory procedures, while T. vaginalis was detected by wet preparation of the genital samples. Additional information was obtained from the patients' records.

Analysis of Results: Results obtained were analysed using Epi Info statistical software, while P values < 0.05 were considered significant.

RESULTS

The incidence of T. vaginalis infection among the women studied was 1.2% (n= 104) (Fig. 1).

Table II shows the age distribution pattern of T. vaginalis infection in Jos: no isolate was recovered from those aged less than 10 years; the peak age of infection was 20- 29 and 30-39 years 0.3% (n= 29) each. The rate of infection among other ages was: 10-19 (0.2%; n= 14); 40-49 (0.2%; n= 16); 50-59 (0.1%; n= 9), and 60 years and above, (0.05%; n= 5). Incomplete data was recorded in two (0.02%) cases. Overall, there was no significant age difference in infection (P>0.05).

Figure 2 shows the clinical features associated with isolation of T. vaginalis: Nil, 6.7% (n= 7); Itching + Rashes, 10.6% (n= 11); Discharge only, 35.6% (n= 37); Rashes only, 8.6% (n= 9); Itching + Discharge, 27.9% (n= 29); Itching only, 4.8% (n= 5); 5.8% (n= 6) had inconclusive information.

Figure 1

Figure 1: Rate of isolation of from female genital specimens in Jos, Plateau state.



Figure 2

Table 1: Age distribution and rate of isolation of from female genital specimens in Jos, Plateau state.

Age (Years)	T. vaginalis Present (%)	T. vaginalis Absent (%)	Total (%)
1-9	0(0.0)	29(0.3)	29 (0.3)
10-19	14 (0.2)	564 (6.7)	578 (6.9)
20-29	29 (0.3)	2201 (26.1)	2230 (26.4)
30-39	29 (0.3)	2989 (35.4)	3018 (35.7)
40-49	16 (0.2)	1299 (15.4)	1315 (15.6)
50-59	9 (0.1)	705 (8.4)	714 (8.5)
60 & Above	5 (0.05)	442 (5.2)	447 (5.3)
Unclassified	2 (0.02)	110(1.3)	112(1.3)
Total	104 (1.2)	8339 (98.8)	8443 (100)

X²= 2.38; OR= 1.92; RR= 1.9; P= 0.12

Figure 3

Figure 2: Clinical Features associated with isolation of from female genital specimens in Jos, Plateau state.



DISCUSSION

The incidence of T. vaginalis infection among women in Jos

was found to be 1.2%. This finding is in line with that of Uneke , et al in Abakaliki, Nigeria₃₇ who obtained an incidence of 2.8% among antenatal women studied. The strong association of T. vaginalis infection with several obstetric and gynaecological problems_{38,39,40} makes its detection among women at any rate of utmost clinical importance, so as to safeguard against them.

Special measures aimed at preventing the transmission of T. vaginalis as well as other agents of STD should be put in place. These include: proper sex education among the adolescents, the youthful girl and boy child by parents in order to imbibe a decent sexual habit ahead of adult life; community based organizations and religious groups as part of their social responsibility, could inculcate sex education especially for the youths as part of their community remodeling measures. These would eventually entrench safe and disciplined sexual habits and culture including the use of protective devices such as condoms and diaphragms. The overall impact would be a reduction in the spread of STDs including T. vaginalis 41:42:43.

The finding of T. vaginalis incidence of 1.2% in the present study is much lower than that of: Adad, et al, in Sao Paulo, Brazil₄₄, where 10%, 19.8%, and 15.9% incidence was reported in different decades on Obstetrics and Gynaecology outpatients using papanicolaou test; Rahman, et al, in Dhaka Bangladesh_{15,45} where incidence of 45.5% and 4.3% was respectively reported among street based and hotel female sex workers; and over 23% prevalence among African-American women in the general population₄₅ using culture and genetic methods of diagnosis. Also, Nwosu, et al 29, and Akerele, et al 30, respectively reported 12.5% and 15% in Nigeria; though higher than the present study with similar methodology: this may be due to variation in the degree of sexual exposure of the two groups as well as differences in protective measures among the two. The much higher 21.9% prevalence recorded by Bakare, et al in Ibadan₄₆ was understandably strictly among commercial sex workers sampled from 18 brothels.

The low incidence of T. vaginalis recorded in the present study may partly be incidental arising from a pool of lightly infected population; it may also have to do with the method of detection of the organism. The wet preparation method which was used in the present study has been shown to be about 50% as sensitive the culture and PCR methods which were methods of choice in most of other studies₄₄,47,48, these are however lacking in this study centre. Improvement in the methods of detection could have probably recorded higher incidence rates.

The peak age of infection with T. vaginalis was found to be 20-29 years. This finding is in consonance with that of Uneke, et al in Abakaliki, Nigeria₃₇ where the peak age of infection was found to be 20-25 years old; and Nessa, et al, in Bangladesh₄₅, where the peak of T. vaginalis infection was detected 18-25 years old range. Besides being a very sexually active, the adventurous disposition of this age bracket and the desire to experiment could as well contribute to these findings₄₉.

The absence of clinical features in some subjects with T. vaginalis infection underscores the need to search for their proper treatment in other cases of STDs most especially in the high risk groups such as commercial sex workers₅₀.

More reliable diagnostic methods should be introduced in Trichomonas research in the country such as: culture methods, PCR, and P16 Immunocytochemistry₅₁. Since these have all been proven to be more sensitive (>98%) and specific (> 97.8%) compared to the current wet preparation method commonly in use in the nations health centres.

In conclusion, T. vaginalis is not an uncommon STD among women in Jos; proper sex education of adolescents and young adults should be carried out by parents, religious organizations, community based organizations, and other relevant character modeling agencies. These, if well articulated and sustained will entrench a culture of restraint built on proper sex education and prevention of STDs. The overall impact being a decrease in the spread and transmission of STDs including T. vaginalis with its avalanche of sequelae.

CORRESPONDENCE TO

JOMBO G T A Department of Medical Microbiology & Parasitology, College of Medical Sciences, University of Calabar, P M B 1115 Calabar, Nigeria. E.mailjombogodwin@yahoo.com Tel-08039726398

References

 Wikipedia: Trichomonas vaginalis. http://en.wikipedia.org/wiki/image:pop_test_trichomonas.JP G Accessed 17th February 2007.
Schwebke J, Burgess D. Trichomoniasis. Clin Microbiol Rev. 2004; 17(4): 794-803.
Gulmezoglu A M, Garner P. Trichomoniasis treatment in

women: a systemic review. Trop Med Int Health 1998; 3: 553-558. 4. Crowell A L, Sanders-Lewis K A, Secor W E. Invitro metronidazole and Tinidazole activities against metronidazole-resistant strains of Trichomonas vaginalis. Antimicrob Agents Chemother 2003; 47(4): 1407-1409. 5. Quan M. Vaginitis, meeting the clinical challenge. Clin Cornerstone 2000; 3: 36-47. 6. Cudmore S, Delgaty K, Hayward-McClelland S, Petrin D, Garber G. Treatment of infections caused by metronidazole resistant Trichomonas vaginalis. Clin Microbiol Rev 2004; 17(4): 783-789 7. Soper D. Trichomoniasis: under control or under controlled?. Am J Obstet Gynaecol 2004; 190(1): 281-290. 8. Meri T, Jokiranta T S, Suhanen L, Meria S. Resistance of Trichomonas vaginalis to metronidazole: Report of the first three cases from Finland and optimization of in vitro susceptibility testing under various oxygen concentrations. J Clin Microbiol 2000; 38(2): 763-767. 9. Engbring J A, O'Brien J L, Alderete J F. Trichomonas vaginalis adhesion proteins display molecular mimicry to metabolic enzymes. Adv Exp Med Biol 1998; 408: 207-223 10. Prasad J H, Abraham S, Kurz K M, George V, Lalitha M K. John R. et al. Reproductive tract infections among young married women in Tamil Nadu, India. Intl Fam Plann Perspectives 2005; 31(2): 73-82. 11. Szarka K, Temesvari P, Kerekes A, Tage A, Repkeny A. Neonatal pneumonia caused by Trichomonas vaginalis. Acta Microbiol Immunol Lung 2002; 49: 15-19. 12. Rughooputh S, Greenwell P. Trichomonas vaginalis: paradigm of a successful sexually transmitted organism. http://www.redorbit.com Accessed 20th March 2007 13. Schwebke J R, Lawing L F. Improved detection by DNA amplification of Trichomonas vaginalis in males. J Clin Microbiol 2002; 40(10): 3681-3683. 14. Cohan M S, Hobbs M, Miller W. Trichomonas vaginalis polymerase chain reaction compared with standard diagnostic and therapeutic protocols for detection and treatment of vaginal trichomoniasis. Clin Infect Dis 2002; 35: 576-586. 15. Rahman M, Alam A, Nessa K, Hossaim A, Nahar S, Datta D, et al. Aetiology of sexually transmitted infections among street-based female sex workers in Dhaka, Bangladesh. J Clin Microbiol 2000; 38(3): 1244-1246. 16. Lowndes C M, Alary M, Meda H, Gnintoungbe C A B, Mukange-Tshibaka L, Adjovi C, et al. Role of core and bridging groups in the transmission dynamics of HIV and STIs in Cotonou, Benin, West Africa. Sex Transm Inf 2002; 78: 169-177 17. Crouch M L, Alderete J F. Trichomonas vaginalis interactions with fibronectin and laminin. Microbiol 1999; 145: 2835-2843. 18. Wendel K A, Rompalo A M, Erbelding E J, Chang T H, Alderete J F. Double stranded RNA viral infection of Trichomonas vaginalis

infecting patients attending a sexually transmitted diseases clinic. J I D 2002; 186: 558-561. 19. Madhu J, Charu G, Mahan K. Sexually transmitted diseases and carcinogenesis. J Obstet Gynaecol Ind 2004; 54(1): 73-76. 20. Guenthner P C, Secor W E, Dezzutti C S. Trichomonas vaginalis induced epithelial monolayer disruption and human immunodeficiency virus type 1 (HIV-1) replication: Implications for the sexual transmission of HIV-I. AIDS 2001; 15: 105-110. 21. McClelland R S, Wang C C, Mandeliya K, Overbaugh J, Reimer M T, Penteleeff D D, et al. Treatment of cervicitis is associated with decreased cervical shedding of HIV-I. AIDS 2001; 15: 105-110. 22. Gilbert R O, Elia G, Beach D H, Klaessig S, Singh B N. Cytopathogenic effect of Trichomonas vaginalis on human vaginal epithelial cells cultured in vitro. Infect Immun 2000; 68: 4200-4206. 23. Rendon-Maldonado J, Espinosa-Cantellano M, Soler C, Toorres J V, Martinez-Palomo A. Trichomonas vaginalis: in vitro attachment and internalization of HIV-I and HIV-I-infected lymphocytes. J Eukaaryat Microbiol 2003; 50: 43-48. 24. Limpakarnjanarat K, Mastro T D, Saisorn S, Uthaivoravit W, Kaewkungwal J, Korattana S, et al. HIV-I and other sexually transmitted infections in a cohort of 313-324. female sex workers in Chiang Rai, Thailand. Sex Transm Infect 1999; 75: 30-35. 25. de Vincenzi I, European Study Group on Heterosexual Transmission of HIV. A longitudinal study of human immunodeficiency virus transmission by heterosexual partners. N Engl J Med 1994; 331: 341-346. 26. Quon D V, d'Oliveiri C E, Johnson P J. Reduced transmission of the ferrodoxin gene in metronidazole-resistant Trichomonas vaginalis. Proc Natl Acad Sci 1992; 89: 4402-4406. 27. Petrin D, Delgaty K, Bhatt R, Garber G. Clinical and Microbiological aspects of 443-450. Trichomonas vaginalis. Clin Microbiol Rev. 1998; 11(2): 300-317 28. Meri T, Jokiranta T S, Suhanen L, Meri S. Resistance of Trichomonas vaginalis to metronidazole: report of the first three cases from Finland and optimization of invitro susceptibility testing under various oxygen concentrations. J Clin Microbiol. 2000; 38(2): 763-767 29. Nwosu M C, Nwosu M N, Mba I E K, Opara C, Nwajuaku C. Genital ulcers and s117. sexual transmitted disease in rural Nigeria. Jnl of Medical Investigation and Practice. 2001; 2: 28-33. 30. Akerele J, Abhulimen P, Okonofua F. Prevalence of asymptomatic genital infection among pregnant women in Benin-city, Nigeria. Afr J Reprod Health 2002; 6(3): 93-97. 31. Pepin J, Sobela F, Deslandes S, Alary M, Wegner K, Khonde N, Kintim F,

Kamuragiye A, Sylla M, et al. Aetiology of urethral discharge in West Africa: the role of Mycoplasma genitalium and Trichomonas vaginalis. Bull World Health Organ 2001; 79(2): 118-126. 32. Buve A, Weiss H A, Laga M, Van Dyek E, Musonda R, Zekeng L, Kahindo M, et al. The epidemiology of trichomoniasis in women in four African cities. AIDS 2001; 15(Suppl. 4): s89-s96. 33. Irvine J W, North M J, Coombs G H. Use of inhibitors to identify essential cysteine proteinases of Trichomonas vaginalis. FEMS Microbiol. Lett. 1997: 149: 45-50. 34. Kwodler L A, Edwards M R, Schofield P J. The intracellular amino acid pools of Giardia intestinalis, Trichomonas vaginalis, and Crithidia luciliae. Exp. Parasitol. 1994; 79: 117-125. 35. Zhang Z F, Begg C B. Is Trichomonas vaginalis a cause of cervical neoplasia? Results from a combined analysis of 24 studies. Int J Epidemiol 1994; 23: 682-690. 36. Shafir S C, Sorvillo F J. Viability of Trichomonas vaginalis in urine: Epidemiology and clinical implications. J Clin Microbiol 2006; 44: 3787-3789 37. Uneke C J, Ugwuoru C D C, Ali M. Trichomonas vaginalis infection among pregnant women in South-Eastern Nigeria. Internet J of Gynaecology & Obstetrics 2006; 6(1): 38. Burja I T, Shurbaji M S. Clinical impact of identifying Trichomonas vaginalis on cervico-vaginal (papanicolaou) smears. Diagn Cytopathol 2001; 24: 195-199. 39. Moodley P, Wilkinson D, Conolly C, Moodley J, Sturn A W. Trichomonas vaginalis is associated with pelvic inflammatory disease in women infected with human immunodeficiency virus. C I D 2002; 34: 519-522. 40. Wang P D, Lin R S. Epidemiologic differences between candidal and trichomonal infections as determined in cytologic smears in Taiwan. Public Health 1995; 109: 41. D'Costa L J, Plummer F A, Bowmer I, Fransen L, Piot P, Ronald A R, Nsanze H. Prostitutes are a major reservoir of sexually transmitted diseases in Nairobi, Kenya. Sex Transm Dis 1985; 12: 64-67. 42. Ndoye L, Mboup S, De Schryver A, Van Dyok E, Moran J, Samb N D, Sakho M L, Thior I, et al. Diagnosis of sexually transmitted infections in female prostitutes in Dakar, Senegal. Sex Transm Dis 1998; 74(Suppl. 1): s112-43. Grosskurth H, Rangaiyan G. Management and control of sexually-transmitted infections and their implications for AIDS control in South-East Asia. J Health Management.2003; 5: 261-276. 44. Adad S J, Vaz de Lima R, Sawan Z T E, Silva M L G, de Souza M A H, Saldonha J C, et al. Frequency of Trichomonas vaginalis, Candida species and Gardnerella vaginalis in cervical-vaginal smears in four different

decades. Sao Paulo Med J 2001; Gaydos C A. Diagnosis of 119(6): 2516-2527. Trichomonas vaginalis infection by PCR using vaginal swab 45. Nessa K, Warris S A, Sultan Z, Monira S, Hossain M, samples. J Clin Nanar S, Habibur R, et al. Microbiol 1998; 36(11): 3205-3210. Epidemiology and Aetiology of sexually transmitted 49. Garbase A C, Rowley J T, Heyman D H L, Berkley S F infections among hotel-based B, Piot P. Global prevalence sex workers in Dhaka, Bangladesh. J Clin Microbiol 2004; and incidence estimates of selected curable STDs. Sex Transm Inf. 1998; 74(Suppl. 42(2): 618-621. 46. Bakare R A, Oni A A, Umar U S, Fayemiwo S A, Fasina 1): s12-s16. 50. Dellabetta G A, Gerbase A C, Holmes K K. Problems, N A, Adewole I F, Shokunbi N U. Prevalence of Trichomonas vaginalis solutions and challenges in amongst commercial sex syndromic management of sexually transmitted diseases. workers (CSWs) in Ibadan, Nigeria. Af J of Clinical and Sex Transm Inf 1998; Experimental Microbiology 74(Suppl, 1): s1-s11. 51. Pantanowitz L, Cao Q J, Goulart R A, Otis C N. 2002; 3(2): 72-77. 47. Emerging Infectious Diseases. Trichomonas vaginalis, Diagnostic utility of P16 HIV, and African-Americans. Immunocytochemistry for Trichomonas in urine cytology. http://www.medscape.com/viewpublication/80 Accessed 20th Cyto Journal 2005; 2(11): March 2007. 2-11.

48. Madico G, Quinn T C, Rompalo A, McKee T J Jr,

Author Information

G. T.A. Jombo, FMCPath

Department of Medical Microbiology & Parasitology, College of Medical Sciences, University of Calabar

S. O. Opajobi, M.Sc.

Department of Medical microbiology, Jos University Teaching Hospital