

Effect of Saliva on pH and volume of gastric contents while sampling from stomach with two different techniques of orogastric intubation

A Hussain, A Hasan Samarkandi, S Shahid Habib, A Al-Omani

Citation

A Hussain, A Hasan Samarkandi, S Shahid Habib, A Al-Omani. *Effect of Saliva on pH and volume of gastric contents while sampling from stomach with two different techniques of orogastric intubation*. The Internet Journal of Anesthesiology. 2006 Volume 13 Number 1.

Abstract

Objectives: To explore the effect of saliva on pH and volume of gastric aspirate by using two different techniques of blind gastric aspiration of gastric contents.

Materials And Methods: This prospective and randomized clinical trial was conducted in the Department of Anaesthesia at King Khalid University Hospital, Al-Riyadh, Saudi Arabia from August to December, 2006 on 140 adult inpatients of either sex, aged 15-70 years and American Society of Anesthesiologists physical status I-II. An orogastric tube was passed by conventional method in Group A and through an endotracheal tube placed in oesophagus in Group B. Gastric contents were aspirated with a large bore, multi-orifices gastric tube after tracheal intubation and analyzed for the presence of bile salts, pH and volume.

Results: Thirty nine (28.57 %) samples were contaminated with duodenal contents and one with blood. Six patients have no gastric contents in Group A due to failed orogastric intubation and none in Group B ($p = 0.0280$). Saliva, by conventional method of orogastric intubation, significantly affected both the pH (A-2 versus B-2: $p < 0.0001$) and volume (A-2 versus B-2: $p = 0.0045$) of gastric contents. Duodenogastric refluxate significantly affected both the pH (A-1 versus A-2: $p = 0.0236$), B-1 versus B-2: $p = 0.0019$) and volume (A-1 versus A-2: $p = 0.0343$), B-1 versus B-2: $p = 0.0005$) of gastric contents.

Conclusion: Saliva significantly affected both the pH and volume of gastric contents when Sampled by conventional method of orogastric intubation. Duodenogastric refluxate significantly affected both the pH and volume of gastric contents.

INTRODUCTION

Pulmonary aspiration of gastric contents is the inhalation of gastric contents into the larynx and lower respiratory tract. Its severity depends upon the nature (pH) and amount (volume) of the aspirated material, and the host's factors that predispose the patient to aspirate ¹.

The importance of pH and volume of gastric contents as a determinant of risk for gastric aspiration has long been discussed ². Since Robert and Shirley's study³ published in 1974, many authors have quoted pH < 2.5 and volume greater than 0.4 ml/kg, or 25 ml of gastric contents for most adult patients, as a risk factor for serious sequelae of pulmonary aspiration of gastric contents. Many studies have been done to examine the effectiveness of various preoperative

medications such as antacids, H_2 – receptor antagonists, proton pump inhibitors and prokinetic drugs alone or in combination to reduce the pH and volume of gastric contents. The most common method employed in these studies for the aspiration of gastric contents remained the blind aspiration with gastric tube.

Saliva can mix up with gastric contents at the level of pharynx while sampling from stomach. Duodenogastric reflux can affect the pH and volume of gastric contents at the stomach level that have already been mixed with gastric contents. Our primary aim of the study was to explore impact of saliva on gastric aspirate by using the conventional method of orogastric intubation versus orogastric tube intubation through an endotracheal tube placed in

oesophagus. Although, the later technique is old and basically described by inserting gastric tube through a naso-oesophageally placed endotracheal tube^{4, 5}. We modified this technique and passed it orally to prevent epistaxis but no body has utilized this technique in previous studies. To see the impact of Saliva on gastric aspirate, we have to exclude those samples contaminated with duodenogastric refluxate. The secondary aim of study was to see the impact of duodenogastric refluxate on gastric contents. In other words, whether these two potential sources of contamination i.e. saliva and duodenogastric refluxate of gastric contents at two levels i.e. hypopharynx and stomach are significant in clinical practice or not? This issue has never been considered important in any previous study while evaluating the effectiveness of drugs used for the prophylaxis of acid aspiration syndrome.

PATIENTS AND METHODS

The study was approved by the College of Medicine Research Centre (CMRC) and College Ethics Committee. Written informed consent was obtained from all the patients.

PATIENTS AND GROUP ASSIGNMENT

We explore the effect of duodenogastric refluxate on the pH and volume of gastric contents and then compared the effect of saliva on the pH and volume of gastric contents by using two methods of orogastric intubation in the Department of Anaesthesia at King Khalid University Hospital, Al-Riyadh, Saudi Arabia on 140 inpatients aged 15-70 years of either sex and American Society of Anaesthesiologists (ASA) physical status I-II, to be intubated with cuffed endotracheal tube.

Patients suffering from the disorders of salivary glands or upper gastrointestinal tract or past history of operations on the salivary glands and upper gastrointestinal tract, drugs known to alter the secretory function of salivary gland, i.e. parasympathomimetics e.g. pyridostigmine, parasympatholytics, e.g. hyosine, antihistamines, e.g. chlorpheniramine, etc., receiving medications known to affect the secretory and /or motor functions of the stomach e.g. ranitidine, omeprazole. etc, Body Mass Index (BMI) more than 40 kg/m², Mallampati class V and /or mouth opening less than 5 centimetres and /or thyromental distance less than 6.5 centimetres and/or history of difficult intubation, parturients and intestinal obstruction were excluded from the study. Gastric contents mixed with blood in the gastric tube were also not included in the statistical analysis while

analyzing pH and volume of gastric contents.

We prepared two sets of envelopes of the same size, colour and shape and packed one set with pieces of papers written on them "conventional method of orogastric intubation A" while other set containing pieces of papers written "orogastric intubation through endotracheal tube B". These envelopes were mixed together. On the pre-operative anaesthesia visit, a day before surgery, the nature and purpose of the study was explained to each patient. We asked each patient to pick up only one envelope from the envelopes. Thus, the patients were allocated either to Group A (conventional method of orogastric intubation) or Group B (orogastric intubation through an endotracheal tube placed in oesophagus) randomly by sealed envelope method. Age, sex, weight, height, BMI, ASA physical status were recorded for each patient. All patients were premedicated with oral diazepam 10 mg at 9.00 p.m. According to the Hospital policy, all patients were fasted from 12 midnight and Dextrose water 5% + 0.45 % NaCl + KCL 20 mmol/L started intravenously from 6.00 a.m. onwards at the rate of 2-3 ml/kg/hour.

COLLECTION AND ANALYSIS OF GASTRIC CONTENTS

In the operating room, routine monitors were attached to the patients and turned on. After pre-oxygenation with 100 % O₂ by face mask using four breaths vital capacity method, anaesthesia was induced with injection fentanyl 1-3 µg/kg, propofol 2-3 mg/kg and rocuronium 0.6-0.9 mg/kg. The lungs were ventilated taking care not to inflate the stomach. Maintaining cricoid pressure, trachea was intubated with cuffed endotracheal tube. Placement and position of endotracheal tube was confirmed with EtCO₂ monitor and then secured properly.

After establishing stable anaesthesia, a predetermined length marked with adhesive tape (Xiphoid process to ear lobules-from ear lobules to nasal tip) of stomach tube₆ (Jamjoom Medical Industries, Jeddah, Saudi Arabia) sized 18 F was passed orally in the stomach in Group A while in Group B gastric tube of the same size was passed through an endotracheal tube sized 8.5 mm internal diameter, coated internally with paraffin liquid, placed in oesophagus with anterior displacement of larynx.. Placement of gastric tube within the stomach by either method was verified by auscultation over the epigastrium during insufflation of 10-15 ml of air. Gastric contents were gently aspirated manually with 60 ml of syringe. Applying manual pressure

over the epigastrium while the patient was in supine and then left and right lateral positions, gastric tube was then manipulated to ensure maximum emptying of gastric contents. Stomach tube was removed in Group A while in Group B; stomach tube was removed followed by esophageally placed endotracheal tube. Time was noted with stop watch to complete the procedure of orogastric intubation in both groups. Any problem encountered during inserting or removing the oro-esophageally placed endotracheal tube or gastric tube was also recorded. The volume of gastric contents was measured with graduated syringe and pH with pH meter (Model 215 version 3.4, Denver Instrument Company, United States). The pH meter was calibrated using standard buffers at pH values of 4, 7 and 9.20. This pH meter has a precision of 0.01 units over the entire pH range. A minimum of one-millilitre volume of gastric contents was sufficient for pH determination with this pH meter. In case of very little amount of gastric contents, we cut the stomach tube and aspirated gastric material with disposable plastic pipette. Samples less than one- millilitre were considered as no gastric contents because a minimum volume of one- millilitre of gastric contents was sufficient for pH- metery. Using bile salts as a marker for bile, we applied qualitative Hay's Sulphur test for the presence of bile salts. A minimum volume of one millilitre of gastric contents was adequate to perform Hay's Sulphur test. In this test finely powered Sulphur is sprinkled upon the surface of cool (17 °C or below) liquid. If bile salts are present Sulphur sinks down, sooner or later, in accordance with their percentage in the fluid.

(a) If bile salts are present in from 1:5000(0.02 % or 200 µg/ml) to 1:10,000(0.01 % or 100 µg/ml) Sulphur at once begins to sink and all precipitate in two or three minutes; even in a dilution of 1:120,000 (0.0008 % or 8.33 µg/ml) precipitation occurs ₇ .

(b) If Sulphur remains floating upon the surface of liquid, the bile salts are absent.

Anaesthesia was maintained with Air, O₂ and sevoflurane. The patients also received incremental doses of fentanyl and rocuronium as required. At the end of surgery, injection atropine and neostigmine were given to antagonize the residual effect of rocuronium. All patients were extubated in lateral position and then transferred to recovery room.

Time since premedication, time since Nil per Os. (NPO), pH, volume of gastric contents and result of Hay's Sulphur

test were also recorded for each patient. On the basis of Hay's Sulphur test, we further divided Group A into Group A-1 (including samples contaminated with duodenogastric refluxate) and Group A-2 (samples non-contaminated with duodenogastric refluxate) and Group B into Group B-1 (including samples contaminated with duodenogastric refluxate) and Group B-2 (samples non-contaminated with duodenogastric refluxate) to evaluate the effect of duodenogastric refluxate on pH and volume of gastric contents. After subtracting these contaminated samples we, then, compared Group-A-2 with Group B-2 to explore the effect of saliva on gastric pH and volume by utilizing two different techniques of orogastric intubation.

STATISTICAL ANALYSIS

Statistical tests were performed using GraphPad Software, Inc., San Diego, United States, and results are expressed as absolute values (percentage) or mean ± standard deviation (SD).

Statistical comparisons between the two Groups were carried out using two-tailed Student's (unpaired) t test for age, weight, height, BMI, time since premedication, time since NPO, time to complete the procedure of orogastric intubation, pH and volume. Two- tailed Fisher's exact test was applied for sex, ASA physical status and failure rate of orogastric intubation.

A p- value of less than 0.05 was considered statistically significant.

RESULTS

One hundred and forty (140) adult inpatients of either sex undergoing elective General (n=81), Orthopaedic (n=30), Gynaecological (n=14), Urology (n=6), Thoracic (n=4) and Plastic (n=4) and Vascular (n= 1) Surgery were studied. Physical characteristics of patients and timings of events are shown in Table 1. There was no statistically significant difference between Group A and B regarding age, sex, ASA physical status, weight, height, BMI, time since premedication and time since NPO. There was a statistically significant difference between Group A and B (p <0.0001) regarding the time to complete the procedure of orogastric intubation.

We obtained gastric contents of 133 patients. Hay's test was performed on all these samples and was positive in 39 patients (29.32 %). One sample was mixed with blood in Group B. Six patients patient have no gastric contents due to

failed orogastric intubation in Group A. Failure rate of orogastric intubation was higher in Group A compared with Group B (p- value) The average (range) pH and volume of contaminated cases with duodenal contents 4.88 (1.35-7.05) and 45.88 (23.0-87.0) ml and with blood 7.13 and 3.0 ml. These cases were considered as contaminated and not included in statistical analysis while analyzing pH and volume of gastric contents.

Saliva significantly affected both the pH and volume of gastric contents when sampled by conventional method of orogastric intubation. There was a statistically significant difference between Group A-2 and Group B-2 regarding pH ($p < 0.0001$) and volume ($p = 0.0045$) of gastric contents. Duodenogastric refluxate statistically significantly affected both the pH and volume of gastric contents. There was a statistically significant difference between Group A-1 and Group A-2 regarding pH ($p = 0.0236$) and volume ($p = 0.0343$) of gastric contents. The same was true between Group B-1 and B-2 as regard pH ($p = 0.0019$) and volume ($p = 0.0005$) of gastric contents. The detail is shown in table 3.

Figure 1

Table 1: Physical characteristics of patients and timings of events. Values are expressed either as mean \pm SD or numbers (percentage).

Physical characteristics of patients	Group A n = 70	Group B n = 70	p- value
Age (years)	34.86 \pm 12.28	34.07 \pm 12.56	0.7088
Sex			1.0000
Male	35 (50%)	35 (50%)	
Female	35 (50%)	35 (50%)	
ASA physical status			0.3566
Class – I	46 (65.71%)	52 (74.28 %)	
Class – II	24 (34.28 %)	18 (25.71%)	
Weight (kilograms)	69.84 \pm 15.86	74.56 \pm 14.67	0.0695
Height (centimetres)	161.44 \pm 7.79	161.80 \pm 7.51	0.8109
Body Mass Index (kilograms/ meter ²)	26.90 \pm 5.88	26.60 \pm 5.48	0.0793
Timings of events			
Time since premedication (minutes)	786.23 \pm 98.31	791.30 \pm 95.22	0.7570
Time since NPO (minutes)	606.31 \pm 127.02	611.39 \pm 95.20	0.7566
Time to complete orogastric Intubation (seconds)	300.97 \pm 5.88	242.30 \pm 62.95	<0.0001

Figure 2

Table 2: Facts and figures about gastric aspirate.

	Group A	Group B	Total
No. of cases	70	70	140
Male	35	35	70
Female	35	37	70
Failed orogastric intubation *	6	0	6
Samples mixed with blood	0	1	1
Samples available for Hay's Sulphur test	64	69	133
Samples mixed with duodenal contents**	18	21	39
Male ***	7	9	16
Female	11	12	23

Note:

*p-value (0.0280)

**total incidence 29.32%

*** Incidence between male and female ($p = 1.0000$)

Figure 3

Table 3: pH and volume of gastric contents. Values are expressed as mean \pm SD.

Variables	Group A n = 70		Group B n = 70	
	Group A-1 n = 64	Group A-2 n = 46	Group B-1 n = 69	Group B-2 n = 48
pH	3.76 \pm 1.75	3.05 \pm 1.28	2.85 \pm 1.69	2.21 \pm 0.42
Volume (millilitres)	32.64 \pm 19.33	25.17 \pm 16	25.28 \pm 13.14	18.15 \pm 4.77

Note:

(a) Patients with failed orogastric intubation six (6) in Group A and one (1) patient with blood mixed sample in Group B are excluded from the above analysis.

(b) Where

Group A-1 and Group B-1 represent Groups including contaminated samples with duodenogastric refluxate.

Group A-2 and Group B-2 represent Groups excluding contaminated samples with duodenogastric refluxate.

Comparisons between the different Groups

pH & volume between Group A-1 and Group A-2 (p- value 0.0236 & 0.0343).

pH & volume between Group B-1 and Group B-2 (p- value 0.0019 & 0.0005).

pH & volume between Group A-2 and Group B-2 (p-value <0.0001 & 0.0045).

DISCUSSION

Aspiration of gastric contents (Mendelson's syndrome) was first described by Mendelson CL in 1946 in obstetrical cases⁸. Since then a lot of work has been done and published in the form of brief reports, forums, original papers, editorials and review articles in anaesthesia literature. In all the previous studies conducted importance of saliva and duodenogastric reflux (DGR), as possible and potential factors that can affect both the pH and volume of gastric contents, has never been addressed.

While evaluating the effectiveness of pharmacological agents used to decrease the acidity and volume of gastric contents, the most commonly used method is blind aspiration with gastric tube. We may come across four types of secretions from our alimentary tract. These are the salivary secretion, oesophageal secretion, gastric secretion and duodenal fluid.

Saliva is produced by the parotid, submandibular, sublingual and many small buccal glands. The daily secretion of saliva normally ranges between 800 and 1500 millilitres at the basal rate of 0.5 millilitres/minute. Saliva has a pH between 6.0-7.0 with the concentration of bicarbonate 50-70 mEq/L; about two to three times that of plasma. Salivary glands are controlled mainly by parasympathetic nervous signals from the superior and inferior salivatory nuclei in the brain stem. The salivatory nuclei are excited by both taste and tactile stimuli from the tongue and other areas of the mouth and pharynx. Smooth objects in the mouth markedly increase salivation, whereas rough objects cause less salivation and occasionally even inhibit salivation. Salivation can also be stimulated or inhibited by nervous signals arriving in the salivatory nuclei from higher centres of the central nervous system. Sympathetic stimulation can also increase salivation a moderate amount, but much less so than does parasympathetic stimulation⁹.

In this current study, we passed gastric tube through an endotracheal tube (Group B) passed blindly in the oesophagus. We avoided possibility of contamination of gastric contents with pooled saliva in pharynx during inserting, manipulating or particularly removing gastric contents with suction applied to proximal end of the gastric tube either with syringe or wall suction. In the awake states, as already noted above, the basal rate of saliva production is about 0.5 ml/minute, but this may increase to 5 ml/minute with intense stimulation¹⁰.

Firstly, insertion of oropharyngeal airway, act of

laryngoscopy and laryngoscope itself and tracheal tube insertion are the stimulants that increase the production rate of saliva. Secondly, saliva pools due to the lack of swallowing reflex in pharynx. Thirdly, in an intubated patient, the oesophagus may be occluded by inflated endotracheal tube cuff. It is difficult to pass or remove stomach tube without the entry of saliva through the side holes into the tube because the stomach tubes do not have obturator as we use in tracheotomy tubes. If we do not apply suction, with the loss of suction effect we will lose gastric content from the gastric tube. We obtained other number of advantages with this technique. Firstly, under general anaesthesia as noted above, swallowing reflex is depressed and in an intubated patient, the oesophagus may be occluded by inflated endotracheal tube cuff and can interfere with stomach tube insertion. Secondly, this technique also avoids finding the upper oesophageal opening and coiling of the tube in the mouth even after successfully passing the distal end of tube into stomach so giving us higher success rate compared with conventional method. Thirdly, manipulation of gastric tube in and out during different positions is very easy causing no or very minimal trauma to the patient. In previous studies, no investigator has mentioned that pharynx was suctioned before inserting or removing the gastric tube. We did follow the same policy in group A.

Oesophageal secretions are entirely mucoid in character and principally provide lubrication for swallowing. The mucus secreted by the compound mucus glands in the upper oesophagus prevent mucosal excoriation by newly entering food in the upper oesophagus where as the compound mucus gland located near the esophagogastric junction protect the oesophageal wall from digestion by acidic gastric juice that often reflux from the stomach back into the lower oesophagus. We could not find exact pH and volume of oesophageal secretions in the literature. Probably oesophageal secretions do not play any role to contaminate the gastric aspirate.

Gastric juice is a mixture of secretions from gastric glands and epithelial cells. Other components are salt, water, HCl, pepsin, intrinsic factor, and mucus. All of these constituents increase after a meal. The greater the secretory rate of gastric juice, the higher the hydrogen and chloride ion concentrations. The daily secretion of gastric juice is 1500 ml with a pH of 0.8-3.5. At this pH, the hydrogen concentration is about 3 millions times that of the arterial blood. The cephalic phase of gastric secretion occurs even

before food enters the stomach, especially while it is eaten. The cephalic phase of gastric acid secretion is elicited by the sight, smell, and taste of food. This phase of gastric secretion normally accounts for about 20 % of total gastric secretion. The gastric phase is initiated by the presence of food in the stomach and accounts for about 70 % of total gastric secretion. The intestinal phase of gastric secretion starts when food enters the upper portion of small intestine particularly the duodenum. It accounts about 10 % of total gastric secretion. The stomach secretes a few millimetres of gastric juice each hour during the interdigestive period when little or no digestion is occurring anywhere in the gut. However, emotional stimuli frequently increase highly acid and peptic interdigestive secretion to 50 ml/hour or more. This point is very important in anaesthesia practice because all patients undergoing general anaesthesia remain NPO.

Duodenogastric reflux, the trans-pyloric retrograde flow of duodenal contents into the stomach, is well known, well established clinical entity^{11, 12, 13} with variable incidence. Mild to moderate duodenogastric reflux occurs in approximately one third (33%) of normal subjects, and in one third (33%) of patients with non-ulcer dyspepsia as shown by the radiological tests of Keet¹⁴ and Huges et al¹⁵, in other words, the pylorus is normally not competent in a significant percentage of normal subjects and approximately the same percentage of patients with non-ulcer dyspepsia. Raved et al¹⁶ a study based on the results of 4256 primary endoscopies of the stomach and duodenum: 3673 in non-operated patients and 583 in patients having undergone vagotomy and resection. Duodenogastric reflux was judged on the basis of yellow coloration of the gastric juice and regurgitation of bile at the time of examination. Reflux was positive in 330(8.98 %) in non-operated patients and 98 (16.81%) in operated patients. Wolverson et al¹⁷ studied the incidence of duodenogastric reflux in peptic ulcer disease using 99mTc Hydroxy Iminodiacetic acid (HIDA) scan, with a gamma camera in the supine position in control patients and patients with active duodenal ulceration. Cholecystokinin was injected intravenously during the test to contract the gall bladder. Patients with benign gastric ulcers, and a group of age matched controls, were investigated for duodenogastric bile reflux in the sitting position by a nasogastric aspiration technique after 10 % dextrose meal. Of 60 patients with duodenal ulceration 32(53%) were reflux positive, and of 13 control patients 6 (46%) were positive. Of 30 patients with gastric ulceration 17 (53%) were reflux positive, and of 8 out of 15 (53%) control subjects were

positive. The incidence of duodenogastric reflux assessed supine in the fasting state, and seated after a liquid meal, was similar in patients with peptic ulceration and in normal controls. In healthy subjects, duodenogastric reflux occurs sporadically in the interdigestive states. Its underlying mechanisms are poorly understood¹⁸. Our reported incidence 28.57 % is slightly less than the previously reported above mentioned studies.

Duodenal fluid (contents) consists of bile (volume 1000 ml/day; pH 7.8), pancreatic juice (volume 1000 ml/day; pH 9.0-8.3), small intestine secretion (volume 1800 ml/day; pH 7.5-8.0) and Brunner's gland (volume 200 ml/day; pH 8.0-8.9). All these secretions are, of course, alkaline in nature due to HCO_3^- ions. When duodenal contents flow in retrograde fashion, then mix with acid and Pepsin¹⁹ in the stomach and bring the pH towards less acidity thus affecting pH and at the same time increase the volume of gastric contents similar to oral ingestion of sodium bicarbonate. The change in intragastric pH related to variations of the amount of different components in the regurgitated duodenal fluid²⁰. Fushs et al²⁰ studied the variability in the composition of physiological duodenogastric reflux and found pancreatic enzyme aspirate was significantly more often associated with a rise in pH in comparison to bile reflux. To overcome this problem, firstly, we aspirated gastric contents in optimal position of the patient as described by Niinai et al²¹. Secondly; we passed a predetermined length of stomach tube so that it should not go beyond pyloric sphincter. Thirdly, we excluded those samples that were positive for Hay's Sulphur test while analyzing pH and volume of gastric contents. Lastly, the average (range) volume of contaminated cases with duodenal contents was 45.88 (23.0-87.0) ml that can only be aspirated from a storage organ like stomach. It is very amazing that so many rather overflow of studies have been conducted and published in anaesthesia literature but no body has reported even a single case of golden yellow, yellow, dark green or light green colour of gastric aspirate from that one may suspect this is bile or intestinal or Brunner gland secretions.

The Bilitec™ 2000 ambulatory bile reflux recorder is currently the only commercially available device that is proven effective in measuring bile reflux. Using Bilirubin as a marker for bile, the Bilitec 2000 recorder captures the frequency and duration of bile exposure either in the stomach or oesophagus over a 24-hour period. This method was not feasible for us we applied Hay's Sulphur test to

detect bile salts in the gastric contents. This simple, sensitive and fairly reliable test²² depends on the principal that bile salts have the property of reducing the surface tension of fluids in which they are contained²³, was devised in 1886 by Matthew Hay (1855-1932).

After excluding those samples contaminated with duodenogastric refluxate, hypopharynx is the only place where pooled saliva can contaminate the gastric aspirate while inserting or removing the gastric tube. Orogastric intubation through an endotracheal tube placed in oesophagus significantly prevented contamination at this point. This technique provided more accurate results compared with conventional method of Orogastric intubation. In this current study we found that saliva and duodenogastric refluxate significantly affected both the pH and volume of gastric aspirate on the basis of significant p-value. If these values, even, not statistically significant, then, is contamination with either saliva or duodenogastric reflux not important? We believe these factors are still important because we will not get true results regarding pH and volume of gastric contents. These results will be of mixed gastric contents either with duodenogastric refluxate or saliva rather than pure gastric contents.

The common techniques to aspirate the residual volume of gastric contents are Fiberoptic gastroscopy, Indicator dilution technique and Blind aspiration via gastric tube.

In this current study, total gastric volume may have been underestimated by the blind aspiration via gastric tube in each patient due to the functional divisions of the stomach into antral and fundal sacs. A similar error would occur in all patients of both groups and inter-group comparisons are, therefore, valid. This method is simple, inexpensive, and easy to perform and has been widely used in the similar studies. As the effect of a drug on intragastric volume reduction is difficult to demonstrate using blind aspiration via gastric tube via gastric tube, the pH values seem preferable, therefore, for comparisons of results in the literature.

CONCLUSION

Saliva significantly affected both the pH and volume of gastric contents when sampled by conventional method of orogastric intubation. Samples obtained with this method of orogastric intubation are not true gastric contents rather gastric contents mixed with saliva. Aspiration of gastric contents done with orogastric intubation through an

endotracheal tube prevented significantly contamination from saliva. This method possesses high success rate, minimal or no trauma to the patients and is time saving. Duodenogastric refluxate also significantly affected both pH and volume of gastric contents. We used pH as a marker of saliva; more work is needed to identify the specific marker for saliva (like bilirubin, bile salts for bile reflux) in the gastric aspirate to see its impact on gastric contents.

CORRESPONDENCE TO

Dr. Altaf Hussain Senior Registrar, Department of Anaesthesiology (41), King Khalid University Hospital, Post Box No. 7805, Al-Riyadh 11472, Saudi Arabia. Phone: 00966-1-4671597 & 4671599 (Department) Phone: 00966-1-4682963 (Home) Fax: 00966-1-4679364 (Department) E-mail: altafdoa@hotmail.com

References

1. Paul E. Marik. Aspiration Pneumonitis and Aspiration Pneumonia. *N Eng J Med.* March 1, 2001; 344 (9):665-671.
2. James CF, Model JH, Gibbs CP, Kuck EJ, Ruiz BC. Pulmonary aspiration- effect of pH and volume in rat. *Anesth Analg.* 1984; 63:665-8.
3. Roberts RB, Shirley MA. Reducing the risk of acid aspiration during caesarean section. *Anesth Analg.* 1974; 53:859-68.
4. Siegel IB, Kahn RC. Insertion of difficult nasogastric tube through a naso-esophageally endotracheal tube. *Crit Care Med.* 1987; 15:876-877.
5. Jonathan L. Benumof. *Clinical procedures in Anesthesia and Intensive Care.* Philadelphia New York London Hagerstown. J.B. Lippincott Company. 1992:829.
6. McConnell EA. Ten problems with nasogastric tubes and how to solve them. *Kango Gijutsu.* 1980 Nov; 26(15):2071-6.
7. John Dixon Mann. *Physiology and Pathology of Urine.* Griffin. 1913:227.
8. Mendelson CL. The aspiration of stomach contents into the lungs during obstetric anaesthesia. *Am J Obst and Gynecol.* 1946; 52:191-205.
9. Guyton AC. *Textbook of Medical Physiology* 10th Edition. Philadelphia: W.B. Saunders, Inc. 2000; 738-753.
10. Ian Power and Peter Kam. *Principles of Physiology for the Anesthetist.* London. Arnold, 2001; 169.
11. Stein HJ, Kauer WK, Feussner H, Siewert JR. Bile acids as components of the duodenogastric refluxate: detection, relationship to bilirubin, mechanism of injury, and clinical relevance. *Hepatogastroenterology* Jan-Feb 1999; 46(25): 66-73.
12. Schidlbeck NE, Heinrich C, Stellaard F, Paumgartner G, Muller-Lissner SA. Healthy controls have as much bile reflux as gastric ulcer patients. *Gut* 1987; 88:1577-1583.
13. Joel E, Richter. Duodenogastric reflux -induced (alkaline) esophagitis. *Curr Treat Options Gastroenterol.* 2004; 7:53-58.
14. Keet AD. A new tubeless radiological test for duodenogastric reflux. *S Afr Med J.* 1982; 61: 78-81.
15. Hughes K, Robertson DAR, James WB. Duodenogastric reflux in normal and dyspeptic patients. *Clin Radiol.* 1982; 33:461-466.

16. Raved D, Kotsev I, Panaiotov P, Karageorgiev L, Georgiev M, Bozhanin D, and Dimov D. The incidence of duodenogastric reflux and its relation to stomach and duodenal diseases. *Khirurgiia (Sofia)*.1990;43(4):61-64.
17. Wolverson RL, Sorgi M, Mosimann F, Donovan IA, Harding LK, Alexander-Williams J. The incidence of duodenogastric reflux in peptic ulcer disease. *Scand J Gastroenterol Suppl.* 1984; 92:149-50.
18. G.H. Koek, R.Vos, D. Sifrim, R. Cuomo, J. Janseens and J.Tack. Mechanisms underlying duodeno-gastric reflux in man. *Neurogastroenterology and Motil.* April2005; Volume 17: 191.
19. Joel E, Richter. Duodenogastric reflux -induced (alkaline) esophagitis. *Curr Treat Options of Gastroenterol.* 2004; 7:53-58.
20. Fei Dai, Jun Gong, Ru Zhang, Jin-Yan Luo, You-Ling Zhu, Xue-Qin Wang. Assessment of duodenogastric reflux by combined continuous intragastric pH and bilirubin monitoring. *W J Gastroenterol.*2002; 8(2):382-384.
21. Niinai H, Nakao M, Nakatani K, Kawaguchi R, Takezaki T and Kobayashi N. Significance of patient's position in measuring gastric contents. *Masui* 1994; 43(11): 1665-7.
22. James Campbell Todd. *Clinical Diagnosis.* W.B. Saunders Company. Philadelphia. 1914; 155.
23. Joshi A. Rashmi. *A Textbook of Practical Biochemistry.* B. Jain Publishers. New Delhi. 2004; 51.

Author Information

Altaf Hussain, MBBS; DA; MCPS; FCPS

Specialist, Department of Anaesthesiology (41), King Khalid University Hospital

Abdul Hamid Hasan Samarkandi, MBBS; FFARCS

Professor & Chairman, Department of Anaesthesiology (41), College of Medicine & King Saud University Hospital

Syed Shahid Habib, MBBS; M.Phil; FCPS.

Assistant Professor, Department of Physiology, College of Medicine & King Saud University Hospitals

Antar Al-Omani, M.Sc. (Clinical Chemistry)

Laboratory Assistant Director, Clinical Chemistry Unit, King Khalid University Hospital