

Levels of Alkaline Phosphatase (ALP) In Saliva of Patients with Chronic Periodontitis; a Clinical and Biochemical Study

S Desai, H Shinde, J Mudda, V Patil

Citation

S Desai, H Shinde, J Mudda, V Patil. *Levels of Alkaline Phosphatase (ALP) In Saliva of Patients with Chronic Periodontitis; a Clinical and Biochemical Study*. The Internet Journal of Dental Science. 2008 Volume 8 Number 1.

Abstract

Objectives: The purpose of our study was to evaluate ALP levels in saliva of patients with chronic periodontitis. **Materials and methods:** Forty patients were assigned to each one of three groups C0, C3 and C4, based on their largest Community Periodontal Index of Treatment Needs CPITN code, totaling 120 participants. Unstimulated saliva was collected and analyzed with autoanalyzer (RA-XT Technicon® Auto Analyzer, USA) by using International Federation of Clinical Chemistry IFCC method in order to quantify their ALP concentration. **Results:** There were significant differences between levels of ALP from groups C0, C3 and C4. There was a significant positive correlation between clinical parameters and ALP concentration in saliva in each group. **Conclusion:** ALP levels in saliva increased with increase in the CPITN score. Group C0 had the least while group C4 had the highest ALP levels. Salivary ALP levels may be useful as a potential bone turnover marker to establish the diagnosis and prognosis of periodontal disease.

INTRODUCTION

Periodontal disease is one of the common inflammatory diseases with complex etiology and multifactorial in origin. Traditional periodontal diagnosis involves measures of probing depth, gingival recession, probing attachment level using graduated periodontal probe. These are indirect measures of bone loss. ¹

Various factors like probing force, presence of inflammation or long junctional epithelium can affect periodontal probing, leading to inaccuracies in recording the true pocket depth. ² The height of the alveolar bone margin and the shape and form of its outline are also examined with radiographs. Unfortunately, the sensitivity of radiographs in detecting an early osseous lesion is poor. ³ Biochemical markers can detect inflammatory changes in short period of time where as longer period is required to detect measurable changes in bone density using radiographs. ¹

Saliva is an oral fluid, and interest in it as a diagnostic medium has advanced exponentially in the last 10 years. ⁴ Saliva has been used as a diagnostic fluid in medicine and dentistry. ⁵ Salivary components for periodontal diagnosis include enzymes and immunoglobulins, hormones of host

origin, bacteria and bacterial products, ions, and volatile compounds. ⁶⁷

Intracellular enzymes are increasingly released from the damaged cells of periodontal tissues into the gingival crevicular fluid (GCF) and saliva. Several enzymes that are evaluated for the early diagnosis of periodontal disease are aspartate and alanine aminotransferase (AST, ALT), lactate dehydrogenase (LDH), creatine kinase (CK), alkaline and acid phosphatase (ALP, ACP), and gamma glutamyl transferase (GGT). ⁶⁸⁹¹⁰

The enzyme ALP plays a role in bone metabolism. It is a membrane-bound glycoprotein produced by many cells, such as polymorphonuclear leukocytes, osteoblasts, macrophages, and fibroblasts within the area of the periodontium and gingival crevice. ¹¹ Gao J et al. (1999) ¹² found that ALP activity was highest in osteoblasts, moderate in periodontal ligament PDL fibroblasts, and lowest in gingival fibroblasts. No activity was detected in cementoblasts. In the periodontium, ALP is very important enzyme as it is part of normal turnover of periodontal ligament, root cementum and maintenance, and bone homeostasis. Some forms of enzyme are also produced by plaque bacteria. ¹³

Some studies have shown a remarkably increased activity of ALP in the acute phase of periodontal disease, and after the periodontal therapy, the activity of these enzymes restored to the value as found with the healthy persons.¹⁴ ALP level has often been measured in GCF to examine the relationship between periodontal conditions and disease activity.¹¹¹⁵¹⁶ Plagnat et al. (2002)¹⁷ studied ALP in GCF from implants with and without periimplantitis and suggested that ALP could be a promising marker of bone loss around dental implants. Gilbert et al. (2003)¹⁸ studied ALP activity in serum from patients with chronic periodontitis and showed a relationship between loss of attachment in periodontal disease and ALP activity in serum. Perinetti et al. (2002)¹⁹ suggested that ALP activity in GCF reflects the biologic activity in the periodontium during orthodontic movement. Todorovic T et al. (2006)²⁰ examined the activity of CK, LDH, AST, ALT, GGT, ALP and ACP in saliva from patients with periodontal disease before and after periodontal treatment. They concluded that the activities of CK, LDH, AST, ALT, GGT, ALP and ACP enzymes were significantly increased in the saliva of patients with periodontal disease when compared to healthy subjects. Daltaban O et al. (2006)²¹ found no significant differences in the concentrations of ALP between periodontitis and control groups. Most of the studies are related to levels of ALP in GCF of patients with periodontal disease¹¹¹⁵¹⁶¹⁷¹⁸¹⁹²⁰²¹, very few studies have evaluated salivary ALP levels in patients with periodontitis.⁴²⁰²²

The purpose of this study was to evaluate ALP levels in saliva of patients with periodontitis as compared to healthy individuals.

MATERIALS AND METHODS

120 patients visiting out patient department of Periodontics, H.K.E Society's S. Nijalingappa institute of dental sciences and research, Gulbarga, Karnataka were screened and classified according to their CPITN scores, by a single professional, using a WHO 621 periodontal probe.²³²⁴ This index was primarily designed for rapid and practical assessments²⁵ of periodontal treatment needs rather than periodontal status, but it is based on various disease indicators such as gingival bleeding after probing, pocket depth, which are related to tissue destruction.

Patients of both genders between 20 to 65 years of age were included in the study. Individuals, who participated in the study, were assigned to one of the following three groups based on their highest CPITN score found among all the

examined sites:

- Group C0: No periodontal disease (Healthy periodontium).
- Group C3: Pathological pocket 4 to 5 mm in depth. Gingival margin situated on black band of the probe.
- Group C4: Pathological pocket 6mm or more in depth. Black band of the probe not visible.

Other clinical parameters recorded were Gingival index (Loe and Silness 1963) and Turesky - Gilmore - Glickman modification of Quigley - Hein plaque index (1970). Their medical and dental histories were obtained by interview. Patients with systemic diseases associated with an increase in ALP, patients with lesions of oral mucosa other than gingivitis and periodontitis, pregnant, lactating, post menopausal females, patients with removable prosthesis and/or caries, smokers, patients with disturbances in salivary flow related with systemic processes or therapeutic procedures, patients with history of any periodontal treatment in previous 6 months, patients with history of antibiotic therapy in previous 3 months were excluded from the study.

1 ml of whole saliva sample was collected by unstimulated passive drool in a sterile disposable plastic container. Patients were instructed not to brush or eat 8 hours before collection of the samples. No dental examination was carried out 48 hours prior to saliva collection. Samples were stored at 4°C and sent to the laboratory of Biochemistry. ALP activity remains stable at room temperature for four hours; frozen samples should be kept at room temperature for 12 hours to assure complete enzyme reactivation.²⁶ The level of ALP was estimated with an auto analyzer (RA-XT Technicon®) by using International Federation of Clinical Chemistry (IFCC) method.

For analysis, each saliva sample was centrifuged at 5000 rpm for 10 minutes. Reagents were added to about 10 µl of supernatant sample by the auto analyzer and the value of ALT estimated in U/L. The reagents used in estimation of saliva ALP are listed in Table (1).

Figure 1

Table-1: Reagents

ALP Reagent	
p-Nitrophenyl phosphate	16 mmol/L
Mg ²⁺	4 mmol
Tris/Carbonate buffer (pH 10.2 ± 0.2 at 25°C)	-
Aqua 4	
Double deionised 0.2 micron filtered particle free reagent grade water	


Aqua 4 is added to reagent bottle and swirled to dissolve. Vigorous shaking is avoided.

Methodology:

Adaptation by Wilkinson et al. ²⁷ of the Bessy, Lowrey et al. method. ²⁸

Principle:

Figure 2



At pH 10.3 alkaline phosphatase catalyses the hydrolysis of p-Nitrophenyl phosphate to yellow colored p-Nitrophenol and phosphate. The change in absorbance measured by autoanalyzer at 405 nm wavelength is proportional to ALP activity in the sample.

Data collected were compared using unpaired t test.

RESULTS

All patients (56 men and 64 women) selected for the study were 20–65 years old (mean 33 years). Mean and standard deviation of ALP concentration in saliva are reported in Table (2).

Figure 3

Table-2: ALP levels (U/L) in saliva from 120 patients

Group	Mean	Standard deviation
C0	20.20	± 6.17
C3	78.95	± 25.77
C4	136.55	± 77.08

The mean concentration of ALP in saliva for group C0 was 20.20 ± 6.17 U/L; for C3 was 78.95 ± 25.77 U/L; and for C4 was 136.55 ± 77.08 U/L. There was statistically significant difference in ALP concentrations in saliva for the groups C0, C3 and C4.. Comparing the salivary ALP levels of group C0 and group C3, the results were statistically highly significant (t = 9.95, P< 0.005). Comparing the ALP levels of group C0 and group C4, the results were statistically highly significant (t = 8.25, P< 0.001). ALP levels of group C4 were significantly greater than ALP levels of group C3 (t = 3.19, P< 0.01).

There was a significant positive correlation between clinical parameters and ALP concentration in saliva in each group. (Table-3. r = 0.14 to 0.36)

Figure 4

Table-3: Co-efficient correlation (r) between clinical parameters and ALP concentration

Group	Gingival Index	Plaque Index
C0	r = 0.24	r = 0.32
C3	r = 0.20	r = 0.16
C4	r = 0.36	r = 0.16

DISCUSSION

Periodontitis is a chronic destructive category of periodontal disease that progresses to the resorption of alveolar bone, which leads to progressive bone destruction and tooth loss. As a consequence of resorption, breakdown of products are released into periodontal tissues, migrating toward the gingival sulcus and gathering from the surrounding site in whole saliva, where several of them have been identified. ²⁹³⁰

Among the several host enzymes proposed as diagnostic

indicators of periodontal status, ALP was one of the first to be identified.³¹ ALP is released from polymorphonuclear cells (PMNs) during inflammation¹⁰ and from osteoblasts³² and periodontal ligament fibroblasts³³ during bone formation and periodontal regeneration respectively.

ALP activity in serum has been extensively studied, and it was suggested that ALP allows bone mineralization by releasing an organic phosphate that contributes to the deposition of calcium phosphate complexes into the osteoid matrix.^{34,35} ALP might also promote mineralization by hydrolyzing inorganic pyrophosphate, a potent inhibitor of hydroxyapatite crystal formation and dissolution, within the extracellular calcifying matrix vesicles.³⁵

CPITN has shown to be a rapid and practical method of periodontal status assessment and determining treatment needs.³⁶ The proposed method allowed establishing a quantitative relationship between AST levels in saliva and periodontal status assessed through CPITN recordings. This method simplified the clinical procedures involved in sample collection compared to the current diagnosis systems; only a 1.0 ml sample of the saliva must be deposited in a sterile disposable plastic container by the patient by unstimulated passive drool. Stimulated saliva collected by asking the patient to chew a gum base may get contaminated with blood as inflamed periodontal tissue may get damaged during chewing. This can cause alteration in the salivary enzyme levels.³⁷

Previous studies¹¹¹⁵¹⁶¹⁷¹⁸¹⁹²⁰²¹ mainly investigated the activities of this enzyme in gingival crevicular fluid, which is in a much closer contact with periodontal tissues and, due to this, it surely much better reflects the occurrences in them. However, the problem with the gingival crevicular fluid is in that the technique of collecting is rather complicated and that in a routine procedure, which possibly might be established, it would be hardly feasible in practice. Contrary to the gingival crevicular fluid, there is plenty of saliva, the procedure of its sampling is much easier and more bearable for the patient and, however, the same enzymes as those in the gingival crevicular fluid can be detected.²⁰ Because of the simple and non-invasive method of collection saliva samples were preferred in our study instead of GCF samples.

More sophisticated techniques to detect bone loss such as subtraction radiography and computer-assisted image analysis have been used as research tools, but at present, these have not found a place in routine clinical practice.³⁸

Changes in ALP in GCF, saliva and serum have been used as an inflammatory marker of periodontium as well as a marker of bone metabolism.¹¹¹⁸²¹³⁹ Various enzymes in serum have been routinely examined for the diagnosis of systemic diseases. Therefore, no specific laboratory devices are necessary.³⁷ We have used these conventional laboratory tests to estimate ALP levels in saliva sample.

Changes in the bone architecture take time to manifest, changes in the bone turnover can be ascertained relatively early by measuring increased levels of serum and urine biochemical markers of bone turnover,²⁹⁴⁰ such as collagen cross-links, ALP, and Osteocalcin.³⁴⁴¹ As ALP is a marker of bone remodeling, CPITN Code 3 (Pathological pocket 4 to 5 mm in depth) and Code 4 (Pathological pocket 6mm or more in depth) were included and Code 1 (Bleeding observed during or after probing) and Code 2 (Calculus or other plaque retentive factors either seen or felt during probing) were excluded from the study.

In our study, differences in ALP concentrations in saliva for the groups C0, C3 and C4 were statistically significant. Probing pocket depth greater than 6 mm (Group C4) was related to the highest ALP levels. The results showed statistically significant relationship between salivary ALP levels and pocket depth. Our results are in agreement with previous studies of ALP done in saliva.⁴²⁰²² The high levels of ALP in saliva in our study may be due to an increase in the inflammation and bone turnover rate as ALP is produced by PMNs, osteoblasts, macrophages, fibroblasts¹¹ and plaque bacteria¹³ within periodontal tissues or periodontal pocket.¹¹³¹ The increased activity of ALP is probably a consequence of destructive process in the alveolar bone in advanced stages of development of periodontal disease.

In our study, GI was positively correlated with increase in pocket depth, although not statistically significant. (Data is shown in table-3) Our results are in agreement with study done by Yoshie H et al. (2007)⁴² who reported statistically significant increase in GI with increase in pocket depth. The increase in salivary ALP levels might have come from PMNs during inflammation within the periodontal pocket. On the basis of results of this study it can be suggested that the activity of ALP enzyme was significantly increased in the saliva of patients with periodontal disease when compared to healthy subjects. It also established the correlation between the enzyme activity and the value of the gingival index. This is probably a consequence of pathological processes in periodontal tissues from where this

intracellular enzyme is increasingly released into saliva.

CONCLUSION

Salivary ALP levels may be useful as a potential bone turnover marker to establish the diagnosis and prognosis of periodontal disease. Diagnosis of periodontal disease by measuring salivary levels of ALP is economical, feasible, simple and convenient approach that does not require expert examiners. Further studies are warranted to confirm the reliability of saliva ALP levels for screening of periodontal disease.

References

1. McCauley LK, Nohutcu RM. Mediators of periodontal osseous destruction and remodeling: Principles and implications for diagnosis and therapy. *J Periodontol.* 2002; 73(11): 1377-91.
2. Liscgarten MA. Periodontal probing: What does it mean? *J Clin Periodontol.* 1980; 7:165-76.
3. Reddy MS. The use of periodontal probes and radiographs in clinical trials of diagnostic tests. *Ann Periodontol.* 1997; 2: 113-122.
4. Kinney JS, Ramseier CA, Giannobile WV. Oral fluid based biomarkers of alveolar bone loss in periodontitis. *Ann N Y Acad Sci.* 2007; 1098: 230-51.
5. Mandel ID. The diagnostic uses of saliva. *J Oral Pathol Med.* 1990; 19:119-25.
6. Kaufman E, Lamseier IB. Analysis of saliva for periodontal diagnosis- a review. *J Clin Periodontol.* 2000; 27:453-65.
7. Lamster IB, Kaufman E, Grbic JT, Winstone LJ, Singer RE. β -glucuronidase activity in saliva: relationship to clinical periodontal parameters. *J Periodontol.* 2003; 74:353-9.
8. Numabe Y, Hisano A, Kamoi K, Yoshie H, Ito K, Kurihara H. Periodontology Analysis of saliva for periodontal diagnosis and monitoring. *Dentistry in Japan* 2004; 40:115-9.
9. Ozomeric N. Advances in periodontal disease markers. *Clin Chim Acta.* 2004; 343(1-2):1-16.
10. McCulloch CA. Host enzymes in gingival crevicular fluid as diagnostic indicators of periodontitis. *J Clin Periodontol.* 1994; 21:497-506.
11. Chapple IL, Garner I, Saxby MS, Moscrop H, Matthews JB. Prediction and diagnosis of attachment loss by enhanced chemiluminescent assay of crevicular fluid alkaline phosphatase levels. *J Clin Periodontol.* 1999; 26:190-8.
12. Gao J, Symons AL, Haase H and Bartold PM. Should cementoblasts express alkaline phosphatase activity? Preliminary study of rat cementoblasts in vitro. *J Periodontol.* 1999; 70(9):951-9.
13. Shibata Y, Yamashita Y, Miyazaki H, Ueno S, Takehara T. Effective method of discriminating between oral bacterial and human alkaline phosphatase activity. *Oral Microbiol Immunol.* 1994; 9:35-9.
14. Yan F. Alkaline phosphatase level in gingival crevicular fluid of peri-odontitis before and after periodontal treatment. *Chung Hua Kou Chiang Hseueh Tsa Chin.* 1995; 30: 255-66.
15. Nakashima K, Roehrich N, Cimasoni G. Osteocalcin, prostaglandin E2 and alkaline phosphatase in gingival crevicular fluid: their relations to periodontal status. *J Clin Periodontol.* 1994; 21:327-33.
16. Binder TA, Goodson JM, Socransky SS. Gingival fluid levels of acid and alkaline phosphatase. *J Periodontal Res* 1987; 22:14-9.
17. Plagnat D, Giannopoulou C, Carrel A, Bernard JP, Mombelli A, Belser UC. Elastase, alpha2-macroglobulin and alkaline phosphatase in crevicular fluid from implants with and without periimplantitis. *Clin Oral Implants Res.* 2002; 13:227-33.
18. Gibert P, Tramini P, Sieso V, Piva MT. Alkaline phosphatase isozyme activity in serum from patients with chronic periodontitis. *J Periodontal Res.* 2003;38:362-5.
19. Perinetti G, Paolantonio M, D'Attilio M, et al. Alkaline phosphatase activity in gingival crevicular fluid during human orthodontic tooth movement. *Am J Orthod Dentofacial Orthop.* 2002; 122:548-56.
20. Todorovic T, Dozic I, Vicente-Barrero M, Ljuskovic B, Pejovic J, Marjanovic M, et al. Salivary enzymes and periodontal disease. *Med Oral Patol Oral Cir Bucal.* 2006; 11:E115-9.
21. Daltaban O, Saygun I, Bal B, Balos K, and Serdar M. Gingival Crevicular Fluid Alkaline Phosphatase Levels in Postmenopausal Women: Effects of Phase I Periodontal Treatment. *J Periodontol.* 2006;77:67-72.
22. Nakamura M, Slots J. Salivary enzymes: Origin and relationship to periodontal disease. *J Periodontal Res.* 1983; 18: 559-69.
23. WHO. Epidemiology, etiology and prevention of periodontal diseases. Report of WHO Scientific Group. World Health Organ Tech Rep Ser. 1978 ; (621):1-60.
24. Emslie RD. The 621 periodontal probe. *Int Dent J.* 1980; 30: 287-8.
25. Ainamo J, Barmes D, Beagrie G, Cutress T, Martin J, Sardo-Infirri J. Development of the World Health Organization (WHO) community periodontal index of treatment needs (CPITN). *Int Dent J.* 1982; 32: 281-91.
26. David H. Vroon, Zafar Israili. Aminotransferases. In: H. Kenneth Walker, W. Dallas Hall, J Willis Hurst, editors. *Clinical methods- The History, Physical and Laboratory Examinations.* 3rd ed. pp 492-3.
27. Wilkinson JH, Boutwell JH, Winsten S. Evaluation of a new system for kinetic measurement of serum alkaline phosphatase. *Clin Chem.* 1969; 15(6):487-95.
28. Bessy OA, Lowry OH, Brock MJ. *Biol Chem.* 1945; 159:399.
29. Riggs BL, Melton IJ 3rd. Involutional osteoporosis. *N Engl J Med.* 1986; 314:1676-86.
30. McGehee JW Jr, Johnson RB. Biomarkers of bone turnover can be assayed from human saliva. *J Gerontol A Biol Sci Med Sci.* 2004; 59:196-200.
31. Ishikawa I, Cimasoni G. Alkaline phosphatase in human gingival fluid and its relation to periodontitis. *Arch Oral Biol.* 1970; 15:1401-4.
32. Christenson RH. Biochemical markers of bone metabolism: an overview. *Clin Biochem.* 1997; 30:573-93.
33. Groeneveld MC, Van den Bos T, Everts V, Beertsen W. Cell-bound and extracellular matrix associated alkaline phosphatase activity in rat periodontal ligament. Experimental Oral Biology group. *J Periodontal Res.* 1996;31:73-9.
34. Taylor AK, Lueken SA, Libanati C, Baylink DJ. Biochemical markers of bone turnover for the clinical assessment of bone metabolism. *Rheum Dis Clin North Am.* 1994; 20:589-607.
35. Garnero P, Delmas PD. Biochemical markers of bone turnover. Applications for osteoporosis. *Endocrinol Metab Clin North Am.* 1998; 27: 303-23.
36. Cesco Rde T, Ito IY, de Albuquerque RF Jr. Levels of aspartate aminotransferase (AST) in saliva of patients with

different periodontal conditions. J Clin Periodontol. 2003; 30: 752-5.

37. Nomura Y, Tamaki Y, Tanaka T, Arakawa H, Tsurumoto A, Kirimura K, et al. Screening of periodontitis with salivary enzyme tests. J Oral Sci. 2006; 48 (4):177-83.
38. Eley BM, Cox SW. Advances in periodontal diagnosis.9. Potential markers of cell death and tissue degradation. Br Dent J. 1998; 184; 427-30.
39. Totan A, Greabu M, Totan C, Spinu T. Salivary aspartate aminotransferase, alanine aminotransferase and alkaline phosphatase: possible markers in periodontal diseases? Clin Chem Lab Med 2006; 44: 612-5.
40. Srivastava AK, Vliet EL, Lewiecki EM, Maricic M,

Abdelmalek A, Gluck O, et al. Clinical use of serum and urine bone markers in the management of osteoporosis. Curr Med Res Opin. 2005; 21:1015-26.

41. Khosla S, Atkinson EJ, Melton LJ 3rd, Riggs L. Effects of age and estrogen status on serum parathyroid hormone levels and biochemical markers of bone turnover in women: A population-based study. J Clin Endocrinol Metab. 1997; 82:1522-27.
42. Yoshie H, Tai H, Tetsuo K, Oda-Gou E, Nomura Y, Numabe Y, et al. Salivary enzyme levels after scaling and interleukin -1 genotypes in Japanese patients with chronic periodontitis. J Periodontol. 2007; 78(3): 498-503.

Author Information

Shrikar Desai, MDS

Associate Professor, Department of Periodontics, H.K.E. Society's S. Nijalingappa institute of dental sciences and research, Gulbarga, Karnataka.

Harshada Shinde, BDS

Post Graduate student, Department of Periodontics, H.K.E. Society's S. Nijalingappa institute of dental sciences and research, Gulbarga, Karnataka.

Jayashree Mudda, MDS

Professor, Department of Periodontics, H.K.E. Society's S. Nijalingappa institute of dental sciences and research, Gulbarga, Karnataka.

Veena Patil, MDS

Professor, Department of Periodontics, H.K.E. Society's S. Nijalingappa institute of dental sciences and research, Gulbarga, Karnataka.