

False-Positive Whole-Body I-131 Scan In Thyroid Carcinoma Caused By Gastroesophageal Reflux Disease

A Biyi, Y Oufroukhi, A Doudouh

Citation

A Biyi, Y Oufroukhi, A Doudouh. *False-Positive Whole-Body I-131 Scan In Thyroid Carcinoma Caused By Gastroesophageal Reflux Disease*. The Internet Journal of Nuclear Medicine. 2007 Volume 5 Number 1.

Abstract

Many false-positive findings on I-131 scans have been reported. Recognition of them may avoid unnecessary repeated therapeutic doses of radioactive iodine. The authors describe a false positive cervical and mediastinal radioiodine uptake due to gastro oesophageal reflux disease in a 63 yr-old man with papillary thyroid cancer. Through this case report, causes of such scintigraphic features are reviewed.

INTRODUCTION

Thyroid cancer is an hormone - dependent neoplasm, radio sensible in its differentiated shapes. After surgical ablation of the primitive tumor, radio iodine completes this treatment in case of cervical remnant or extra nodal metastases. Whole body ¹³¹I scintigraphy has aided the follow up of differentiated thyroid cancer for several decades. However, this strong tool based on the presence of the sodium iodide symporter (NIS) in the basolateral surface of thyroid follicular cells is not perfect. A wide spectrum of potentially misleading artefacts can arise in ¹³¹I whole body scans from various anatomical variants and physiological processes as well as several unrelated non-thyroidal disease processes [1]. Recognition of potential false-positive iodine-131 scans is critical to avoid the unnecessary exposure to further radiation from repeated therapeutic doses of radioactive iodine. Here, we describe a case of false positive whole body scan due to a gastro-oesophageal motility disorder.

Patient interrogation revealed heartburn and acid regurgitation. He was effectively treated for gastro oesophageal reflux disease.

A repeat scan after eating was negative. Serum thyroglobulin level was under 0,1 ng/ml. So we attributed the scintigraphic abnormalities to the gastroesophageal reflux disease.

CASE REPORT

A 63-yr-old man with a long past of type 2 diabetes (25 years) and more recently a Parkinson disease (5 years), received ten years ago 3, 7 GBq of ¹³¹I for post surgical ablation of residual cervical cells of a papillary carcinoma of the thyroid. Six months later, serum thyroglobulin test and whole body ¹³¹I scan were negative. The patient was considered free of disease and the same results were shown on serum tests and scintigraphic follow-up for the last decade. Recently, a whole body scan performed two days after oral administration of 167 MBq of ¹³¹I showed tree foci in the upper side of the neck and a linear mediastinal uptake.

Figure 1

Figure n°1: Anterior and posterior whole body I-131 scan 48 hours after oral administration of 167 MBq (4,5 mCi) of I-131. Besides physiological uptake in the gastrointestinal tract and the bladder, intense iodine uptake is shown in the anterior and upper side of the neck and mediastinum.



DISCUSSION

Whole-body radioiodine scan is an integral part of the follow-up algorithm for patients with differentiated thyroid carcinoma (DTC). The specificity of such exploration for detecting residual or recurrent local and metastatic disease is generally reported to be greater than 90% [2]. Recognition of normal and pathologic biodistribution of iodine is imperative for the nuclear medicine physician to avoid interpretation pitfalls and unnecessary repeated therapeutic doses [1]. A radioiodine scan showing abnormal uptake outside the thyroid bed must be studied carefully and alternative reasons for the finding must be considered. I-131 is excreted in gastric mucosa and can be seen in the oesophagus and pharynx after regurgitation or swallowing of saliva [3].

Oesophageal motility disorders are frequent in Parkinson

disease. In our case, the patient history, the knowledge of I-131 artefacts and the serum thyroglobulin level all served to identify the abnormal tracer uptake as a false-positive result. The images revealed typical activity in the digestive tract. The linear esophageal activity generally mandates simple additional images following both eating and drinking. In the majority of cases, the intensity and shape of the activity in the oral cavity, pharynx and esophagus will change and a correct diagnosis can be made. In addition, it has been reported that the incidence of false-positive scans increases with the dose used for diagnostic scanning [4]. Therefore, the increase in sensitivity (fewer false-negative scans) obtained with higher doses has to be balanced with the decrease in specificity.

The following table summaries physiopathologic classification of benign and malignant entities that can show a false-positive result on radioiodine scan.

Figure 2

Table n°1: classification of false-positive radioiodine scan causes

Localization related to NIS function	Localization in the thyroid at unusual sites: <ul style="list-style-type: none">• ectopic thyroid tissue [5,6], struma ovarii [7] Non-thyroidal tissues possessing NIS: <ul style="list-style-type: none">• Normal physiological uptake in salivary gland [8], lacrimal gland [9], nasopharynx [10] and lactating breast [11].• Neoplastic uptake: gastric adenocarcinoma [12], Warthin's tumour, adenocarcinoma of lungs [14], teratoma [15], meningioma[16], ovarian tumors[1].• Abnormally located gastric mucosa: Meckel's diverticulum [17], Barrett's oesophagus[18], hiatus hernia [19] and gastric pull up surgeries.
Localization unrelated to NIS function	<ul style="list-style-type: none">• Contamination by physiological secretions: saliva, urine, vomitus, nasal secretions[1].• Gastrointestinal anomalies: Zenker's diverticulum [20], oesophageal stricture, oesophageal motility disorders [3], asymmetrical salivary uptake.• Hepatic uptake by 131-I labelled thyroid hormone [21]• Pleural, peritoneal and pericardial effusions [1, 22], hydrocele, septated gallbladder, ovarian cyst, laryngocele [23], subdural hematoma [24]• Ectopic kidney, transplanted kidney, urinary tract divertcul, hydronephrosis [1]• Inflammatory conditions: cholecystitis [22], liver abscess [25], pulmonary tuberculosis [26], sebaceous cyst [22], rheumatoid arthritis associated lung disease, skin burn, psoriasis [1,22].
Uptake due to unknown mechanism	<ul style="list-style-type: none">• Thymic uptake [27], ectasia of carotid , pectus excavatum [9]. Recent myocardial infarct[1] , Intestinal scar [28].

References

1. Shapiro B, Rufini V, Jarwan A, et al. Artifacts, anatomic and physiological variants, and unrelated diseases that might cause false-positive whole-body I-131 scan in patients with thyroid cancer. Semin Nucl Med. 2000; 30:115–132.
2. McDougall IR. Whole-body scintigraphy with radioiodine-131: a comprehensive list of false-positives with some examples. Clin Nucl Med. 1995; 20:869–875
3. Bakheet S, Hammami MM: False-positive thyroid cancer metastasis on whole-body radioiodine scanning due to retained radioactivity in the oesophagus. Eur J Nucl Med 20:415, 1993
4. Waxman A. Ramanna L. Chapman N, et al. Significance of I-131 scan dose in patients with thyroid cancer:

determination of ablation (concise communication). *J Nucl-Med* 1981;22:861-865.

5. Strain J, Oates E, Nejad A: Unusual appearance of lingual thyroid in congenital hypothyroidism. *Clin Nucl Med* 23: 460,1998
6. Sironi M, Assi A, Andruccioli M, et al: Submandibular ectopic thyroid gland. *Clin Nucl Med* 21:585, 1996.
7. Joja I, Asakawa T, Mitsumori A, et al: 1-123 uptake in non-functional struma ovarii. *Clin Nucl Med* 23:10-12, 1998
8. Spitzweg C, Joba W, Eisenmenger W, et al: Analysis of human sodium iodide symporter gene expression in extrathyroidal tissues and cloning of its complementary deoxyribonucleic acids from salivary gland, mammary gland, and gastric mucosa. *J Clin Endocrinol Metab* 83:1746-1751, 1998
9. Bakheet SM, Hammami MM: False-positive whole- body scan in thyroid cancer patients due to unrelated pathology. *Clin Nucl Med* 19:325, 1994.
10. Park HM, Wellman HN: Hot nose after 1-131 Sodium Iodide thyroablation therapy. *Clin Nucl Med* 2:130, 1992
11. Duong RB, Fernandez-Ulloa M, Planitz MK, et al: I123 breast uptake in a young primipara with postpartum transient thyrotoxicosis. *Clin Nucl Med* 8:35-36,1983.
12. Wu S, Kollin J, Coodley E, et al. I131 total body scan: localisation of disseminated gastric adenocarcinoma. Case report and survey of the literature. *J Nucl Med* 1984; 25:1204-9
13. Burt RW. Accumulation of I 123 in a Warthin's tumour. *Clin Nucl Med* 1978; 3:155-6.
14. Fernandez-Ulloa M, Maxon HR, Mehta S, et al. Iodine 131 uptake by primary lung adenocarcinoma, misinterpretation of 131I scan. *JAMA* 1976; 236:857-8
15. Lakshmanan M, Reynolds JC, DelVecchio S, et al. Pelvic radioiodine uptake in a rectal wall teratoma after thyroidectomy for papillary carcinoma. *J Nucl Med* 1992; 33:1848-50.
16. Geatti O, Shapiro B, Orsolon PG, et al: An unusual false-positive scan in a patient with pericardial effusion. *Clin Nucl Med* 19:678, 1994
17. Caplan RH, Gundersen GA, Abellera M, et al. Uptake of

I131 by a Meckel's diverticulum mimicking metastatic thyroid cancer. *Clin Nucl Med* 1987;12:760-2.

18. Berquist TH, Nolan NG, Stephens DM, et al. Radioisotope scintigraphy in diagnosis of Barrett's esophagus. *AJR* 1971;123:401-11.
19. Willis LL, Cowan RJ. Mediastinal uptake of I131 in a hiatal hernia mimicking recurrence of papillary thyroid carcinoma. *Clin Nucl Med* 1993;18:961-3.
20. Boulahdour H, Meignan M, Melliére D, and al. False positive I-131 scan induced by Zenker's diverticulum. *Clin Nucl Med* 1992;17:243-4.
21. Rosenbaum RC, Johnston GS, Valente WA: Frequency of hepatic visualization during 131I imaging for metastatic thyroid carcinoma. *Clin Nucl Med* 13:657-660, 1988
22. Brucker-Davis F, Reynolds JC, Skarulis MC, and al. False positive iodine131 whole body scans due to cholecystitis and sebaceous cyst. *J Nucl Med* 1996; 37:1690-3.
23. Schmidt M, Dietlein M, Schroeder U, Schicha H. False-positive uptake of I-131 in a laryngocele mimicking thyroid remnant after thyroidectomy for papillary thyroid carcinoma. *Clin Nucl Med*. 2006; 31:716-7.
24. Ong SC, Eng DN, Sundram FX, Chan LL. A novel case of false-positive I-131 whole-body scan in thyroid carcinoma caused by subdural hematoma. *Clin Nucl Med*. 2004 Mar;29(3):164-6
25. Pena Pardo FJ, Crespo de la Jara A, Fernández Morejón FJ, Sureda González M, Forteza Vila J, Brugarolas Masllorens A. Solitary focus in the liver in a thyroid cancer patient after a whole body scan with 131 iodine. *Rev Esp Med Nucl*. 2007; 26: 294-6.
26. Bakheet SM, Hammami MM, Powe J, Bazarbashi M, Al Suhaibani H. Radioiodine uptake in inactive pulmonary tuberculosis. *Eur J Nucl Med*. 1999; 26:659-662.
27. Vermiglio F, Baudin E, Travagli JP, et al. Iodine concentration by the thymus in thyroid carcinoma. *J Nucl Med*. 1996; 37:1830-1831.
28. Sioka C, Dimakopoulos N, Kouraklis G, Kotsalou I, Zouboulidis. A False-positive whole-body scan after I-131 therapy in a patient with intestinal scar. *Clin Nucl Med* 2006; 31:232-3.

Author Information

Abdelhamid Biyi, MD

Department of nuclear medicine MILITARY HOSPITAL MOHAMMED V. HAY RIAD RABAT. MOROCCO

Yacir Oufroukhi, MD

Department of nuclear medicine MILITARY HOSPITAL MOHAMMED V. HAY RIAD RABAT. MOROCCO

Abderrahim Doudouh, MD

Department of nuclear medicine MILITARY HOSPITAL MOHAMMED V. HAY RIAD RABAT. MOROCCO