

Foreign Bodies of the Respiratory Tract

M Poirier

Citation

M Poirier. *Foreign Bodies of the Respiratory Tract*. The Internet Journal of Emergency Medicine. 2003 Volume 2 Number 1.

Abstract

Aspirated foreign bodies, whether in the upper airway, or the lower airway continue to present challenges to physicians who care for children in the acute setting. Appropriate maneuvers to relieve foreign body upper airway obstruction are age dependent. With children and infants who have foreign bodies in their lower airway, a high index of suspicion is required in order to make a timely diagnosis. Often the initial choking events are not witnessed, and the delayed symptoms may mimic other common conditions. Proper anticipatory guidance and education is the optimal way of reducing the tragic outcomes of choking events. This article reviews the current principles in the management of children and infants with foreign bodies in their respiratory tracts.

INTRODUCTION

Upper airway foreign body obstruction and aspirated foreign bodies are a major cause of childhood mortality and morbidity and continue to present challenges to physicians who care for children. These events are not new occurrences. In 1633, the London physician, Stephen Bradwell, wrote, "Of things that endanger stopping of the breath in swallowing, some are sharp and some blunt... I have heard of a child in Woodstreet strangled with a grape."⁽¹⁾ Then, as now, bystanders often perform prompt, effective life-saving maneuvers to children with foreign bodies in their upper airway. These life-saving maneuvers are age dependent and are usually performed in the field prior to arrival at a health care facility. Additionally, the diagnosis of a foreign body in the lower airway has added difficulty because these choking events are unwitnessed and the delayed symptoms may mimic other common conditions such as asthma, recurrent pneumonia or upper respiratory tract infections. This article reviews the clinical presentation, diagnostic work-up and appropriate management of children and infants with foreign bodies in their respiratory tract.

BACKGROUND

Hundreds of pediatric choking deaths occur every year in the United States.^(2,3,4,5,6,7,8,9,10,11) More than 300 children die annually due to foreign body upper airway obstruction. Studies show that ninety percent of deaths occur in infants and children less than 5 years of age and 65% in those less than 2 years of age. The child younger than three years of age is at greatest risk for dying from a foreign body aspiration.⁽⁴⁾ These deaths are usually attributable to

aspiration of foods, toys or other small objects. Organic debris is commonly retrieved from the upper airway by appropriate first aid maneuvers in children who have acute upper airway obstruction and by bronchoscopy in the lower airway. Hot dogs, candy, grapes and peanuts are the most common foods recovered.^(5,6,7,8,9,10,11,12,13,14) The shape and smoothness of these foods is thought to enable them to pass easily into the upper airway.

The description of the "café coronary" in 1963, heightened the awareness of the causes, prevention and emergency treatment of food-related choking events in adults.⁽¹⁵⁾ This paper reported nine cases of sudden death in adult restaurant patrons that occurred when a piece of meat acutely obstructed the victim's upper airway. The authors suggested an association in adults between choking on food, excessive alcohol intake and poorly fitted dentures. A 1984 report focused increased attention on food-related choking episodes in children.⁽¹⁴⁾ Analyzing national data on all identified food-related asphyxiation events in infants and children up to 9 years of age from 1979 to 1981, one such death occurred every five days. More than 90% took place in infants and children younger than 5 years of age and 65% in infants younger than 2 years of age. Round or cylindrical foods were the most common culprits in these events. Foods are the most common cause of choking events in toddlers. The natural curiosity of the toddler, the ubiquitous presence of small foods in the home and the lack of an efficient grinding surface before the eruption of the back molars could explain the high propensity for choking in this age group.⁽¹⁶⁾

In 1979, the US Consumer Product Safety Commission passed regulations to control marketing of nonfood items that are threats to cause choking in infants and children.⁽¹⁷⁾ Small toys, rubber balloons, nails, tacks and bolts are the main offenders.⁽¹⁸⁾ Other dangerous objects include earrings, straight pins, aluminium foil, rocks and other small metal objects.^(19,22,23) Especially worrisome are rubber balloons, now the leading cause of pediatric choking deaths from children's toy products.⁽²⁴⁾ Several features of balloons explain why they are so dangerous. Their collapsibility allows them to pass through the vocal cords and lodge in the carina. In addition, their inflatability prevents any air passing through to the lungs. Many communities have directed efforts to prevent childhood deaths from choking on balloons by banning rubber balloons in daycares, schools and hospitals. Other communities are proponents of safer non-rubber balloon alternatives.

FOREIGN OBJECTS IN THE NOSE

Foreign objects are commonly placed by young children into their nose. The classic presentation of an unexplained foul smelling nasal discharge which is unilateral and persistent is common in unwitnessed events. Other less specific symptoms include chronic sinusitis, recurrent epistaxis and halitosis.^(25,26,27,28) Frequently the presenting complaint is that the parent witnessed the young child place the object in the nose. The removal of these nasal foreign bodies is generally straightforward with adequate visualization and appropriate instruments necessary for a successful removal. Visualization may be improved with applying a topical vasoconstrictor to the nasal mucosa, using a high intensity light source and using suction to remove any nasal secretions. Complications include trauma to the nasal mucosa, potential fracture to the cribriform plate and potentiating aspiration of the foreign body into the proximal airway.^(25,26)

MANEUVERS TO REMOVE UPPER AIRWAY FOREIGN BODIES

Certain assumptions underlie the current recommendations for treatment of airway obstruction in children and infants. Although cardiac arrest with secondary airway obstruction is often seen in adults, in infants and children airway obstruction with secondary cardiac arrest is much more common. A foreign body that completely obstructs the upper airway is an immediate threat to life and must be removed immediately. However, if the child can speak or breathe or is coughing, the foreign object may dislodge spontaneously, making any first aid maneuvers potentially detrimental by

converting a partial airway obstruction into a complete airway obstruction. Partial airway obstruction with very poor air exchange, or complete airway obstruction with cyanosis requires immediate interventions to avoid permanent disability or death.

Which maneuver is used to relieve an upper airway obstruction depends on the age of the child. The pediatric and emergency medicine community, including the Committee on Pediatric Emergency Medicine of the American Academy of Pediatrics (AAP), the American Heart Association (AHA), and the Red Cross, consider the abdominal thrust maneuver the most effective method of relieving complete airway obstruction in children older than 1 year of age.^(29, 30) The utility of this maneuver based in the following principles: eighty percent of respiratory effort is from diaphragmatic contraction, abdominal inward pressure compresses the diaphragm upwards thus raising intrathoracic pressure, and a sudden rapid increase in intrathoracic pressure may expel the obstructing object.⁽³⁰⁾ As the victim becomes hypoxic from obstruction, muscle tone diminishes thus making the abdominal thrust maneuver more effective.

The AAP and AHA recommend the head-down back-blow maneuver followed by the chest-thrust maneuver for relieving airway obstruction in the child younger than 1 year of age. Some experts prefer the abdominal thrust maneuver for this age group as well as the older child and no studies deny its effectiveness. However, critics of using this maneuver in the child less than 1 year of age, cite cases of ruptured abdominal organs, pneumomediastinum and even a thrombosed aorta.^(31,32,33) Possible explanations for these complications in children less than one year of age is that 70% of abdominal thrust maneuver in infants were performed by untrained individuals and 50% of the time these maneuvers were performed by people who learn of this technique by reading newspapers and lay magazines. Additionally infants have relatively large stomachs, livers and spleens as compared to older children. This anatomical difference could contribute to the higher complication rate of the abdominal thrust maneuver in infants.

Another argument against using the abdominal-thrust maneuver in an infant or young child relates to the increased compliance of the infant's chest wall compared with that of the older child. Because of this compliance the infant's chest wall absorbs some of the energy from the abdominal-thrust maneuver, resulting in chest wall expansion at the expense of lung expansion.⁽³⁴⁾ This can make the abdominal thrust

maneuver in the infant less effective than in the older child by not producing pressure changes significant enough to expel the foreign object from an obstructed airway.⁽³⁴⁾

The head-down back-blow maneuver, the first step recommended for infants, combines the force of gravity with the force the chest compression generates to expel intrathoracic air. Some investigators are concerned that the sudden air acceleration associated with the back blows may cause an object to paradoxically travel rather into the airway. Indeed, studies indicate that the back-blow maneuver may make the object move caudally in accordance to Newton's third law of motion, "to every action there is always an equal reaction."⁽³⁵⁾ In a patient who is awake, back blows also can cause the airway to open or reconfigure slightly, forcing the foreign body further into the airway and worsening the obstruction. In the face-down infant who is unconscious, this appears to be less relevant.

In controlled experiments on a closed system comparing the abdominal-thrust maneuver with the back-blow maneuver, the generated intrathoracic pressures were 13 mm Hg for the abdominal thrust maneuver and 32.5 mm Hg for the back-blow maneuver.⁽³⁶⁾ Conversely, in another controlled experiment, the abdominal-thrust maneuver generated an average maximum pressure of 21 mm Hg compressed with 11 mm HG for the back-blow maneuver.⁽³⁶⁾ The pressure response persisted much longer with the abdominal-thrust maneuver (0.7 seconds) than with the back-blow maneuver (0.01 seconds).⁽³⁶⁾ These contradictory findings make the scientific proof of the efficacy of the abdominal thrust maneuver less clear.

The second step recommended for infants, the chest-thrust maneuver, uses sternal compression to increase intrathoracic pressure in an effort to expel the foreign object from the airway. These compressions are similar to those performed for cardiopulmonary resuscitation. The rescuer uses two or three fingers to compress the sternum approximately one-third to one-half the depth of the chest. This corresponds to a depth of about 1/2 to 1 inch, although these measurements are often not precise. The chest compressions theoretically can inflict rib and cardiac damage in infants, but a study designed to investigate this possibility found no significant rib injuries or fractures in a large review of infants who underwent chest compression, and no such injuries are reported elsewhere in the literature.⁽³⁷⁾

Blind finger sweeps of the oropharynx to remove a foreign body that cannot be visualized should not be performed.

Attempt to remove a foreign body only if it is visible.⁽²⁹⁾ In administering chest compressions or subdiaphragmatic abdominal thrust for the unconscious, nonbreathing child, open the child's mouth by grasping both the tongue and the lower jaw between the thumb and fingers and lifting, thus performing a tongue-jaw lift procedure. This action draws the tongue away from the back of the throat to eliminate the possibility of the tongue causing a partial airway obstruction.

If the back blows, abdominal thrust or chest thrust are unsuccessful in the nonbreathing, unconscious victim, removal of the foreign body should be attempted. This optimally would be performed in a controlled setting such as the emergency department under direct visualization using laryngoscopy with forceps. If the foreign body is visualized, extraction using Magill forceps would be the procedure of choice. Endotracheal intubation may force the foreign body distally enough to partially ventilate and more importantly oxygenate the child. The pop-off valve of the bag-valve system should then be occluded enough to deliver sufficient volume of oxygenation and ventilation. If these attempts fail, the physician should proceed to create a surgical airway.

LOWER AIRWAY FOREIGN BODIES

As the foreign body passes through the vocal cord into the trachea and bronchi, acute symptoms of choking, gagging and distress may resolve thus making the diagnosis more difficult. The common clinical triad of cough, wheezing and decreased breath sounds in children with foreign bodies in their lower airway is not consistent. Between 50% and 90% of children with foreign body aspiration have a suggestive history, most commonly of an acute episode of paroxysmal cough.^(3,6,7,12) Other common symptoms that occur at the time of aspiration are cyanosis, choking and dyspnea.^(3,7,38) In addition, the symptoms of these patients mimic many other pulmonary conditions including asthma, bronchiolitis and pneumonia.^(6,8,11,39) Only half of all children are diagnosed correctly in the first 24 hours after a choking episode and an additional 30% receive the correct diagnosis in the following week.^(5,10,12) The remainder may have delays in diagnosis of weeks to years.^(5,7,9,10,12,40) Specifically for this reason the clinician must have a high index of suspicion in order to diagnose an aspirated foreign body in the lower airway. The most common foreign bodies retrieved from the lower airway are foods, nuts, and seeds.⁽⁴¹⁾ The most common inert foreign objects retrieved from the lower airway are toys.⁽⁴¹⁾

RADIOGRAPHIC STUDIES

Diagnostic imaging plays a variable role in identifying foreign bodies in the lower airway. Since the majority of foreign bodies in the lower airway are food, they are not radiopaque and usually are not apparent radiographically. However, appropriate radiographic studies can aid in localizing the site of the foreign body in the lower airway.^(5,6,7,8,12,13,39,40,42,43) Radiographic evaluation in the emergency department should start with AP and lateral views of the chest and neck. Differential inflation of the affected lung is the most common abnormality identified. This differential inflation may be accentuated by fluoroscopy, lateral decubitus films or assisted expiratory views.^(3,6,7,12,13,42,44,45) The lateral decubitus film and the assisted expiratory views accentuate the ball-valve mechanism of the partial obstructive bronchial foreign body, leading to residual hyperinflation of the involved lung and sometimes mediastinal shift toward the unaffected side. Resorption atelectasis beyond the site of an airway foreign body and the presence of pulmonary infiltrates are other indirect signs of foreign bodies in the lower airway.^(3,5,6,7,12) Although CT scan, xeroradiography and ultrasonography have been advocated for foreign body imaging in the lower airway, their utility has yet to be demonstrated.^(3,46) For localizing peanuts and other seeds in the airway, MRI has recently been suggested as an appropriate imaging modality.⁽⁴⁷⁾ Given the limited sensitivity of radiographic findings in children who have aspirated foreign materials into their lower airways, clinical judgment must dictate whether children should be scheduled for diagnostic bronchoscopy in the face of negative radiographic studies.⁽⁴⁵⁾

MANAGEMENT

The majority of children presenting to the emergency department with foreign bodies in their lower airway are not in extremis. Administration of supplemental oxygen and close observation with monitoring are usually all that is required in the stable patient. As reviewed earlier if the foreign body is in the pharynx and can be easily visualized and the child's cooperation can be enlisted, the foreign body may be safely removed in the emergency department using appropriate instruments. In the vast majority of cases, airway foreign body removal is best achieved in the operating suite. Rigid bronchoscopy under general anesthesia is the procedure of choice for removal of most foreign bodies of the trachea or bronchi.^(3,5,6,7,13,40,42,48) The availability of a fiberoptic-telescope system with a ventilating bronchoscope

enables ventilation and instrumentation to occur simultaneously. Other less accepted methods include the use of a Fogarty balloon catheter in conjunction with a rigid bronchoscope to facilitate removal of the airway foreign body.^(42,44) Postural drainage with percussion is thought to be potentially dangerous and its use is discouraged.

Post bronchoscopy complication rates range from 2% to 8%, with the most common complications including subglottic edema from the endoscopic procedure and residual lung atelectasis.^(8,13,44) Flexible fiberoptic bronchoscopy is not indicated in the removal of foreign objects from the lower airway. Reasons include inability to administer anesthetic agents, inert potential for further airway compromise and greater difficulty in controlling instruments.^(38,40,45,48) Excessive pressure or biting motions upon removal of the foreign body may lead to fragmentation and further morbidity. Sharp objects should be removed within the lumen of the scope to minimize mucosal injury. In the rare occurrence that the patient is so unstable that general anesthetic is not indicated, topical anesthetic and restraints could be used to remove tracheal or bronchial foreign bodies.^(45,48,49) Open thoracotomy is indicated when rigid bronchoscopy fails or when objects are tightly impacted.^(5,7,40)

CONCLUSION

Regulatory changes and increased public awareness have reduced the number of choking deaths, but foreign bodies in the airway still remain a significant problem. Tragic outcomes will only be reduced when primary care physicians stress to their patients, their patient's families, and the communities the importance of prevention through anticipatory guidance. The appropriate maneuvers of relieving foreign body airway obstruction should be taught to parents and caretakers. Additionally, in an emergency department setting, laryngoscopy and forceps extraction must be rapidly undertaken when indicated. Aspirated foreign bodies in the lower airway require the clinician to have a high index of suspicion in order to make a timely diagnosis. Finally, it is important to remember, if one foreign body is found in the respiratory tract, always look for others.

References

1. Bradwell R: Help for Sudden Accidents Endangering Life. London: Thomas Purfoot, 1633:113-114
2. Spitz L: Management of ingested foreign bodies in children. *Br Med J* 1971; 4:469-472
3. McGuirt W, Holmes K, Feehs R, et al: Tracheobronchial foreign bodies. *Laryngoscope* 1988; 98:615-618

4. National Safety Council: Accident Facts. Itasca, IL: National Safety Council, 1992:5
5. Hamilton A, Carswell F, Wisheart J: The Bristol Children's Hospital experience of tracheobronchial foreign bodies 1977-87. *Bristol Med Chir J* 1989; 104:72-74
6. Steen K, Zimmerman T: Tracheobronchial aspiration of foreign bodies in children: a study of 94 cases. *Laryngoscope* 1990; 100:525-527
7. Laks Y, Barzilay Z: Foreign body aspiration in childhood. *Pediatr Emerg Care* 1988; 4:102-106
8. Esclamado R, Richardson M: Laryngotracheal foreign bodies in children. *Am J Dis Child* 1987; 141:259-262
9. Mantel K, Butenandt I: Tracheobronchial foreign body aspiration in childhood. *Eur J Pediatr* 1986; 145:211-216
10. Brown T, Clark C: Inhaled foreign bodies in children. *Med J Aust* 1983; 2:322-325
11. Lima J: Laryngeal foreign bodies in children: a persistent, life threatening problem. *Laryngoscope* 1989; 99:415-420
12. Wiseman N: The diagnosis of foreign body aspiration in childhood. *J Pediatr Surg* 1984; 19:531-535
13. Black R, Johnson D, Matlak M: Bronchoscopic removal of aspirated foreign bodies in children. *J Pediatr Surg* 1994; 29:682-684
14. Harris C, Baker S, Smith G, et al: Childhood asphyxiation by food: A national analysis and overview. *JAMA* 1984; 251:2231-2235
15. Haugen R: The cafe coronary: Sudden deaths in restaurants. *JAMA* 1963; 186:142
16. Rovin J, Rodgers B: Pediatric foreign body aspiration. *Pediatr Rev* 2000; 21:86-90
17. US Consumer Product Safety Commission: Method for Identifying Toys and Other Articles Intended for Use by Children Under 3 Years of Age which Present Choking, Aspiration, of Ingestion Hazards Because of Small Parts, 16 CFR 1501. Washington DC: General Services Administration, 1979
18. Tinsworth D: Analysis of Choking-Related Hazards Associated With Children's Products. Washington DC: US Consumer Product Safety Commission, September 1989
19. Becker P: Earring aspiration and other jewelry hazards. *Pediatrics* 1986; 78:494-496
20. Press S, Liberman J: Aspiration through a "sip-up" straw. *Am J Dis Child* 1986; 140:1090-1091
21. Ross M, Janik J: "Foil Tab" aspiration and retropharyngeal abscess in a toddler. *JAMA* 1988; 260:3130
22. Arnold R, Hoffman A, Brutinel W, et al: "Barbie" doll curler aspiration into the upper trachea. *Am J Dis Child* 1987; 141:1325-1326
23. Myer C: Foreign body aspiration. *Am J Dis Child* 1988; 142:485-486
24. Ryan C, Yacoub W, Paton T, et al: Childhood deaths for toy balloons. *Am J Dis Child* 1990; 144:1221-1224
25. Brownstein D, Hodge D: Foreign bodies of the eye, ear, and nose. *Pediatr Emerg Care* 1988; 4:215-218
26. Baker M: Foreign bodies of the ears and nose in childhood. *Pediatr Emerg Care* 1987; 3:67-70
27. Bennett J: An unexpected cause of halitosis. *J R Army Med Corps* 1988; 134:151-152
28. Fireman P: Diagnosis of sinusitis in children: emphasis on the history and physical examination. *J Allergy Clin Immunol* 1992; 90:433-436
29. Committee on Pediatric Emergency Medicine: First aid for the choking child. *Pediatrics* 1993; 92:477-479
30. Heimlich J: A life-saving maneuver to prevent food choking. *JAMA* 1975; 234:398-401
31. Visintine R, Baick C: Ruptured stomach after Heimlich maneuver. *JAMA* 1975; 234:415
32. Croom D: Ruptured stomach after attempted Heimlich maneuver. *JAMA* 1983; 250:2602-2603
33. Fink J, Klein R: Complications of the Heimlich maneuver. *J Pediatr Surg* 1989; 24:486-487
34. Hughes T: The choking controversy. *Emergency* 1979; 11:34
35. Day R, Crelin E, Dubois A: Choking: The Heimlich abdominal thrust vs. back blows: An approach to measurement of inertial aerodynamic forces. *Pediatr* 1982; 70:113-119
36. Gordon A, Belton M, Ridolpho P: Emergency management of foreign body airway obstruction. In: Safar P, Elam J, eds. *Advances in Cardiopulmonary Resuscitation*. New York: Springer-Verlag, 1977:39-50
37. Spevak M, Kleinman P, Belanger P, et al: Cardiopulmonary resuscitation and rib fractures in infants. *JAMA* 1994; 272:617-618
38. Banerjee A, Rao KS, Khanna S, et al: Laryngo-tracheobronchial foreign bodies in children. *J Laryngol Otol* 1988; 102:1029-1032
39. Gyi B, Austin J, Eisenberg L: Radiolucent intratracheal foreign body mistaken for croup in a 9-year-old boy. *Am J Dis Child* 1984; 138:749-750
40. Weissberg D, Schwartz I: Foreign bodies in the tracheobronchial tree. *Chest* 1987; 91:730-733
41. Metrangelo S, Monetti C, Meneghini L, Zadra N, Giusti F: Eight years' experience with foreign-body aspiration in children: what is really important in a timely diagnosis? *J Pediatr Surg* 1999; 34:1229-1231
42. Yane D, Pritchard M, Colville C, et al: Bronchoscopy for aspirated foreign bodies in children. *Arch Surg* 1988; 123:1988-1993
43. Svedstrom E, Puhakka H, Kero P: How accurate is chest radiography in the diagnosis of tracheobronchial foreign bodies in children? *Pediatr Radiol* 1989; 19:520-522
44. O'Neill J, Holcomb G, Neblett W: Pediatric gastrointestinal foreign body ingestions. *Ann Emerg Med* 1983; 13:112-115
45. Friedman E: Tracheobronchial foreign bodies. *Otolaryngol Clin North Am* 2000; 33:179-185
46. Cotton E, Yasuda K: Foreign body aspiration. *Pediatr Clin North Am* 1984; 31:937-941
47. Imaizumi H, Kaneko M, Nara S, et al: Definitive diagnosis and identification of peanuts in the airways using magnetic resonance imaging techniques. *Ann Emerg Med* 1994; 23:1379-1382
48. Witt W: The role of rigid endoscopy in foreign body management. *Ear Nose Throat J* 1985; 64:70-74
49. Holinger L: Foreign bodies of the larynx, trachea, and bronchi. In: Blueston C, Stool S, Scheetz M, eds. *Pediatric otolaryngology*. Philadelphia: WB Saunders, 1990

Author Information

Michael P. Poirier, M.D.

Associate Professor of Pediatrics, Eastern Virginia Medical School, Children's Hospital of The King's Daughters