

The Prevalence Of Atopic Eczema / Dermatitis Syndrome (AEDS) In Basrah Providence, IRAQ

I Edan Al-saimary, K Al-Hamdi, S Bakr

Citation

I Edan Al-saimary, K Al-Hamdi, S Bakr. *The Prevalence Of Atopic Eczema / Dermatitis Syndrome (AEDS) In Basrah Providence, IRAQ*. The Internet Journal of Dermatology. 2005 Volume 3 Number 2.

Abstract

A total of 484 patients suffering from atopic eczema /dermatitis syndrome (AEDS) were examined in the main hospitals of the Basrah providence, IRAQ.

It has been found that 211 (43.6%) of patients were males, and 273 (56.4%) were females. Infantile and childhood patients had a higher prevalence ratio (34.7% and 27.5%) respectively than the three adulthood groups (17.8, 11.4% and 8.7%) respectively ($P < 0.001$). Also, males has a higher prevalence ratio of AEDs in stages (1,4,5) in percentages (55.36, 43.28, and 32.84) respectively, while females were highly infected with AEDS in stages (2,3) in percentages (61.65 and 60.35) respectively ($P < 0.001$).

INTRODUCTION

The term "atopic dermatitis" AD was first proposed by wise and sulzberg in 1933⁽¹⁾, other synonyms used are atopic eczema, constitutional eczema or dermatitis, prurigo Besnier, and many other⁽²⁾. The term "atopic eczema/ dermatitis syndrome" AEDs was proposed from the EACCI nomenclature task face in 2003 as the 'umberella' term to cover the different subtypes of atopic dermatitis (AD); extrinsic and intrinsic types^(1,3). The frequency of AEDs can vary from 10% to over 25%, and up to 45% according to the population studied, i.e. pediatric or adult population, epidemiologic study, in or out patients, and the criteria used for atopy diagnosis^(4,5,6). AEDS is increasing in frequency over time. A study of Scottish school children showed a prevalence of 5.3% in 1964 and 12% in 1986⁽⁷⁾. The 12-months period prevalence estimates for the 6 to 7 year age group ranged from under 2% in Iran to over 16% in Japan and Sweden^(8,9). In the 13-to-14-year age group, disease prevalence ranged from less than 1% in Albania to over 17% in Nigeria⁽⁷⁾. Other studies found that AEDs affects up to 20% of childhood population, in 65% AEDs had resolved by 7 years of age, in 74% AEDs had resolved by 16 years of age^(10,11,12).

The aim of the present study was to determine the prevalence of atopic eczema/ dermatitis syndrome in various age groups of both sexes in Basrah governorate, IRAQ.

MATERIALS & METHODS

A total of 484 patients (211 males and 273 females) in various age groups were included in this study. The patients were suffering from atopic dermatitis attending the out patients of department of dermatology of main hospitals in Basrah governorate (out patients based study), in addition to specialized private clinics. The patients were examined, and atopic dermatitis diagnosed under supervision of dermatologists based on criteria of Hanifin & Rajka (1980)⁽¹³⁾, Spergel & Schneider (1999)⁽¹⁴⁾, and Stanway (2005)⁽¹⁵⁾.

The study was carried out during a period from November to July 2005.

The grouping of patients: The patients (males and females) were grouped into five groups according to^(7,16,17).

These groups are:

Infantile group (1): less than two years,

Childhood group (2): from 2 to 11 years.

Adulthood groups: over than 11 years, and then subdivided into: group (3): 11 to < 20 years,
group (4): 20 to < 30 years.

And group (5): over than 30 years.

STATISTICAL ANALYSIS

Chi-square test and ANOVA test were used according to SPSS program ver. 11.

RESULTS

From a total of 484 patients with AEDS, 211 (43.6%) were males, and 273 (56.4%) were females Fig (1). Patients of early age stage (1,2) recorded highly prevalence ratio of AEDS of about (34.7% and 27.5%) respectively than adulthood stage patients (3,4,5) that recorded (17.8%, 11.4% and 8.7%) respectively (P<0.001).

There are a highly significant differences between sexes of patients with AEDs (P<0.001) Fig (2). Males has a highly prevalence ratio of AEDS in stages (1,4,5) in percentages (55.36%, 43.28% and 32.84%) respectively, while females have a higher AEDS prevalence ratio in stages (2,3) and in percentages (61.65% and 60.35%) respectively. Fig (3)

{image:1}

{image:2}

{image:3}

DISCUSSION

No previous study evaluated demographical/ or epidemiological screening of atopic eczema/dermatitis syndrome in Iraqi governorates, so, the present investigation was found necessary to determine the prevalence ratio and status of AEDS.

Studies of other countries showed that AEDS is increasing in frequency over time. Reasons suggested for the increasing prevalence include: decreased rates of breast-feeding_(18,19), earlier introduction of weaning foods, wide spread use of food additives, and changes in the formulation of infant formula_(20,21,22). And environmental factors favouring the expression_(23,24,25). Many other studies supported the results of our study: a slight female preponderance has been demonstrated in some studies_(11,16). A small female preponderance was also noted in the International study of asthma and allergies in childhood (ISAAC) study with an overall female: male ratio of 1.3:1 being higher in countries with the highest prevalence of atopic dermatitis symptoms₍₇₎. Our results interested in age related-atopic dermatitis confirmed by study of Sugiura, et al. (1998) that found the atopic dermatitis is predominantly a disease of infantile and childhood, and prevalence estimates show a continuous reduction with increasing age₍₂₆₎. A recent

Norwegian study based on review of medical records found a prevalence of atopic dermatitis of 13% in patients under the age of 20 years₍₂₇₎.

CONCLUSION

We concluded that there is a high prevalence ratio of atopic eczema/dermatitis syndrome in the Basrah governorate. Females were more frequently infected with AEDS than males. Infantile and childhood stages had a higher prevalence ratio than all adulthood groups (P<0.001).

CORRESPONDENCE TO

IHSAN EDAN AL-SAIMARY; Assist. Prof. Dept. Microbiology, Coll. Medicine, Univ. Basrah, PO BOX 696 Ashar, Basrah, IRAQ E-mail: Ihsanalsaimary@yahoo.com

References

1. Wuthrich, B. and Grendelmeier, P.S. (2003). The atopic eczema/ dermatitis syndrome. *J. Invest. Allergol. Clin. Immunol*; 13 (1):1-5.
2. Werfel, T. and Kapp, A. (1998). Environmental and other major provocation factors in atopic dermatitis. *Allergy*, 53:731-739.
3. Wuthrich, B. (1989). Atopic dermatitis flare provoked by inhalant allergens. *Dermatologica*, 178: 51-53.
4. Wuthrich, B. (1999). Clinical aspects, epidemiology and prognosis of atopic dermatitis. *Ann. Allergy Asthma Immunol.*, 83:464-470.
5. Wollenberg, A. and Bieber, T. (2000). Atopic dermatitis: from the genes to skin lesions. *Allergy*, 55: 205-213.
6. Oppel, T., Schuller, E., Gunther, S., Moderer, M., Haberstock, J., Bieber, T. and Wollenberg, A. (2000). Phenotyping of epidermal dendritic cells allows the differentiation between extrinsic and intrinsic forms of atopic dermatitis. *Br.J.Dermatol.*, 143:1193-1198.
7. Charman C.R. and Williams H.C. (2002). *Epidemiology*. In: Bieber, T. and Leung, D.Y.M. (Edits). *Atopic dermatitis*. Marcel Dekker, Inc. New York, Ch. 2 PP:21-42.
8. Sugiura, H., Umamoto, N., Deguchi, H., Murata, Y., Tanaka, K., Sawai, T., Omoto, M., Uchiyama, M., Kiriya, T. and Uehara, M. (1998). Prevalence of childhood and adolescent atopic dermatitis in a Japanese population: comparison with the disease frequency examined 20 years ago. *Acta Derm. Venereol. (Stockh)*, 78: 293-294.
9. Leung, R. and Ho, P. (1994). Asthma, Allergy and atopy in three south-east Asian populations. *Thorax*, 49: 1205-1210.
10. Emerson, R.M., Williams, H.C. and Allen, B.R. (1998). Severity distribution of atopic dermatitis in the community and its relationship to secondary referral. *Br. J. Dermatol.*, 139: 73-76.
11. Marks, M., Kilkenny, M., Plunkett, A. and Merlin, K.(1999). The prevalence of common skin conditions in Australian school students: 2-atopic dermatitis. *Br. J. Dermatol.*, 140: 468-473.
12. Popescus C.M., Popescu, R., Williams, H., Forsea, D. (1998). Community validation of the United Kingdom diagnostic criteria for atopic dermatitis in Romanian schoolchildren. *Br. J. Dermatol.*, 138:436-442.
13. Hanifin, J.M. and Rajka, G. (1980). Diagnostic features of atopic dermatitis. *Acta Derm. Venereol. (Stockh)*, 92(suppl.):44-47.

14. Spergel, J.M. and Schneider, L.C. (1999). Atopic dermatitis. *inter. J. Asthma Allergy Immunol.*, 1(1):1-16.
15. Stanway, A. (2005). Atopic dermatitis. Available in <http://DermNetNZ.bookstore.Net.3October2005>.
16. Herd, R.M., Tidman, M.J., Prescott, R.J. and Hunter, J.A.A. (1996). Prevalence of atopic eczema in the community the Lothian atopic dermatitis study. *Br.J. Dermatol.*, 135:18-19.
17. Niishioka, K. (1996). Atopic eczema of the adult type in Japan. *Australas J. Dermatol.*, 37: 57-59.
18. Bleiker, T.O., Shahidullah, H., Dutton, E. and Brown, R. A.C. (2000). The prevalence and incidence of atopic dermatitis in a birth cohort: the importance of family history of atopy. *Arch. Dermatol.*, 136:274.
19. Mar, A., Tam, M., Jolley, D. and Marks, R. (1999). The cumulative incidence of atopic dermatitis in the first 12 months among Chinese, Vietnamese and Caucasian infants born in Melbourne, Australia. *J. Am. Acad. Dermatol.* 40:597-602.
20. Williams, H.C. (1992). Is the prevalence of atopic dermatitis increasing? *Clin.Exp.Derm.*, 17:385-391.
21. Kay, J., Gawkrödger, D.J., Mortimer, M.J. and Jaron, A.G. (1994). The prevalence of childhood atopic eczema in a general population. *J. Am. Acad. Dermatol.*, 30: 35-39.
22. Williams, H.C. and Stachan, D.P. (1998). The natural history of childhood eczema: observations from the 1958 British cohort study. *Br.J.Dermatol.*, 139:834-839.
23. Leung, R. (1994). Asthma, allergy and atopy in south-east Asian immigrants in Australia. *Aust NZJ. Med.*, 24: 255.
24. Dotterud, L.K., Kvammen, B., Lund, E. and Falk, E. (1995). Prevalence and some clinical aspects of atopic dermatitis in the community of Sqr-Varanger. *Acta. Derm. Venereol. (Stoch)*, 75:50-53.
25. Dotterud, L.K. and Falk, E.S. (1999). Atopic disease among adults in northern Russia, an area with heavy air pollution. *Acta Derm Venereol.*, 79: 448-450.
26. Sugiura, H., Umemoto, N., Deguchi, H., Murata, Y., Tanaka, K., Sawai, T., Omoto, M., Uchiyama, M., Kiriama, T. and Uehara, M. (1998). Prevalence of childhood and adolescent atopic dermatitis in Japanese population: comparison with the disease frequency examined 20 year ago-*Acta. Derm. Venereol. (Stockh)*, 78:293-294.
27. Falk, E. (1993). Atopic diseases in Norwegian Lapps. *Acta. Derm. Venereol. (Stockh) suppl.* 182: 10-14.

Author Information

Ihsan Edan Al-saimary

Department of Microbiology, Coll. Medicine, University of Basrah

Khalil E. Al-Hamdi

Department of Medicine, Coll. Medicine, University of Basrah

Sundis S. Bakr

Department of Microbiology, Coll. Medicine, University of Basrah