

Post Cesarean Section Caecal Volvulus: Case Report And Literature Review.

F Mohamed, M Abukhater, B Ahmad, A Mohamed

Citation

F Mohamed, M Abukhater, B Ahmad, A Mohamed. *Post Cesarean Section Caecal Volvulus: Case Report And Literature Review..* The Internet Journal of Gynecology and Obstetrics. 2009 Volume 14 Number 1.

Abstract

Caecal volvulus is a rare, but potentially fatal cause of intestinal obstruction. Accurate diagnosis is vital as delay in the surgical treatment of this condition may lead to an increased incidence of gangrene of the caecum and a higher mortality. We report a case of caecal volvulus in a 43 years old female who presented with small bowel obstruction 10 days after a cesarean section. We also review the literature for the clinical and radiological features of the condition.

INTRODUCTION

Caecal volvulus is axial twisting that occurs involving the caecum, terminal ileum and ascending colon. It is responsible for 25%–40% of all colonic volvulus and together with ascending colon volvulus accounts for only 1 per cent of mechanical bowel obstructions. Caecal volvulus after cesarean section is very rare, with only three cases of ileo-caecal volvulus reported in the English literature in the last two decades. We report a case of caecal volvulus in a 43 years old female which occurred 10 days after a cesarean section.

CASE REPORT

A 43 year female, presented to the emergency department at KFMC Riyadh, KSA with severe on and off abdominal colic of two days duration associated with frequent vomiting and constipation. She had a cesarean section 10 days before her presentation and 4 cesarean sections before. On examination she was in pain, mildly dehydrated not jaundiced or anemic. Vital signs were normal. The abdomen was distended; lax and not tender, and bowel sounds were absent. Laboratory investigations showed WBC of 14.4×10^3 . Liver function test, urea and electrolytes were normal. The plain abdominal x ray showed a dilated loop of bowel occupying the right side of the abdomen extending across the mid line to the left with absence of air in the colon in the supine film (figure 1&2) and air fluids levels in the caecum in the erect film. The CT of the abdomen demonstrated the typical bird beak sign (figure 3).

Figure 1

Figure 1: Supine plain x ray of the abdomen showing the dilated caecum occupying the right side of the abdomen extending across the mid line to the left. (Observe absence of air in the remaining colon).




Figure 2

Figure 2: The erect plain x-ray abdomen showing air fluid level in the caecum.



Figure 3

Figure 3: The abdominal CT showing severely dilated caecum with air fluid level with the progressively tapering efferent and afferent bowel loops terminating at the site of torsion demonstrating the bird beak sign.



Diagnosis of caecal volvulus was established and after fluid resuscitation the patient was taken for laparotomy under general anesthesia. Intraoperatively the caecum was found axially twisted and severely distended with mild distention of the small bowel (figure 4). The caecum was untwisted and fixed to the lateral abdominal wall (caecopexy). The patient didn't show signs of recurrence after 4 years of followed up in the outpatient clinic.

Figure 4

Figure 4: Intra operative photograph showing severely dilated caecum and mildly dilated small bowel.



DISCUSSION

Caecal volvulus is axial twisting that occurs involving the caecum, terminal ileum and ascending colon. First noted by Hildanus in the 16th century and later reviewed by Rokitsky in 1837, it is a rare condition with a reported incidence ranging from 2.8 to 7.1 per million people per year(1). It is responsible for 25%–40% of all volvulus involving the colon (2). Caecal and ascending colon volvulus accounts for only about 1 per cent of mechanical bowel obstructions (3, 4).

There are two prerequisites for caecal volvulus to occur: a segment of mobile caecum and ascending colon and a point of fixation about which torsion may occur (5, 6, 7). The mobility results from either incomplete embryologic rotation of the bowel or improper developmental fusion of the mesentery of the caecum and ascending colon with the posterior parietal peritoneum. In addition to the prerequisite of a freely mobile caecum, several additional predisposing factors have been implicated in the genesis of caecal volvulus, although the relative importance of these factors is not clear (8). These include concomitant acute medical problems, pregnancy, distal colon obstruction and a previous laparotomy (9).

Previous abdominal surgery, resulting in intraabdominal adhesions which could act as a fulcrum for caecal volvulus, has been considered important by several investigators (10, 11).

In the post-operative setting caecal volvulus has been described as a complication following left colectomy, cholecystectomy, gastric resection, incarcerated femoral

hernia repair, appendectomy and various laparoscopic procedures (12-15). It has also been reported following kidney transplantation (16) and nephrectomy (17, 18). Presumably any surgical procedure which might require some degree of medial visceral rotation or disruption of the fusion plane between the caecum and ascending colon with the lateral peritoneum could provide for sufficient mobility to allow caecal volvulus to occur (19).

In female patients previous gynaecological operations may be a more important factor in predisposing to caecal volvulus than appendectomy, which has been suggested as a major antecedent (9). This may account for the fact that caecal volvulus in western societies is considerably more common in women than in men (9, 20, 11, 21, 22). Caecal volvulus after cesarean section is rare. Three cases of ileo-caecal volvulus post cesarean section have been reported in the English literature in the last two decades (23, 24, and 25). Marren A et al reported a case of caecal volvulus associated with intestinal malrotation immediately following caesarean section (26).

The reasons for the association of caecal volvulus with pregnancy and cesarean section are not clear. Some authors suggested that pregnant patients are at high risk for caecal volvulus due to uterine displacement of the ascending colon (27, 28), together with the fact that caesarean section rapidly increases intra-abdominal space, thereby increasing the likelihood of volvulus (29).

Intestinal volvulus including caecal volvulus is also reported as a rare complication of laparoscopic procedures such as laparoscopic cholecystectomy. Ferguson et al published a literature review of bowel obstruction following laparoscopic procedures in January 2008. Of the 12 reported cases 8 cases followed laparoscopic cholecystectomy, two of them were caecal volvulus (30). Agahi et al report a case of caecal volvulus after laparoscopic adjustable gastric band surgery (31). Similarly Wales et al report a case of caecal volvulus in a patient who underwent laparoscopy-assisted sigmoid resection for sigmoid volvulus one year earlier (32).

Accurate diagnosis is vital as delay in the surgical treatment of this condition may lead to an increased incidence of gangrene of the caecum and a higher mortality (33).

As most patients with acute caecal volvulus present with clinical pictures suggestive of intestinal obstruction, abdominal radiography is frequently obtained as the initial diagnostic imaging (2).

Diagnosis can be made by plain abdominal X-ray in more than half the cases on the basis of caecal distention, proximal small bowel distention with air-fluid levels, a gasless distal colon (34), and identification of a characteristic “coffee bean deformity” directed toward the left upper quadrant.

The coffee bean deformity is seen on conventional radiographs or tomograms, as a rounded focal collection of air-distended bowel with haustral creases in the left upper quadrant that resembles a coffee bean.

The dilated caecum usually assumes a 'comma-shape', retains its haustral markings and may be located anywhere within the abdomen but is most frequently seen centrally or occupying the left upper quadrant (33).

However, given the non-specific nature of these radiological findings and the unusual occurrence of caecal volvulus, many of the patients are erroneously given the diagnoses of small bowel obstruction (35).

Barium enema shows lack of filling of the caecum, often with a “beaked” termination of the column of contrast. The use of contrast studies in the diagnosis of caecal volvulus has been debated. It has been argued that such investigations usually serve only to delay definitive surgical treatment (36). Barium enema should not be done when the patient is critically ill or when there is suspicion of gangrene or perforation (9).

Reduction of acaecal volvulus by barium enema has been reported and most probably succeeded in some patients. Also fiberoptic colonoscopy has been suggested by Ghazi et al (9, 37) as a means of preoperative decompression of volvulus of the colon occurring proximal to the sigmoid, however in contrast to management of sigmoid volvulus, attempts to treat acute caecal volvulus by non-operative decompression could be dangerous (9).

CT is not only a valuable diagnostic technique in diagnosing caecal volvulus and its complications, but it is also useful in distinguishing the three pathophysiological types of caecal volvulus (axial torsion, loop type and caecal bascule) (38).

Recognition of the CT signs of caecal volvulus is critical because the findings at clinical examination are often vague and because CT scan is usually the imaging technique of choice for patients presenting with acute abdominal pain (39). CT reveals the presence and location of the volvulus and gives the added benefit of allowing early identification of potentially fatal complications, such as ischemia and

perforation. Three-dimensional (3D) reconstructions may further improve diagnostic capabilities by allowing visualization of the entire bowel in a single image (40).

The ‘‘coffee bean’’, ‘‘bird beak’’, and ‘‘whirl’’ signs are three of the common CT findings associated with acute caecal volvulus (2). The ‘‘coffee bean’’ sign generally refers to an axial view of a dilated caecum filled with air and fluid that may be visualised anywhere within the abdominal cavity. The ‘‘bird beaks’’ are images correlating with the progressively tapering efferent and afferent bowel loops terminating at the site of torsion. The ‘‘whirl sign’’ is a description applied to the CT image of a soft tissue mass with internal architecture containing swirling strands of soft tissue and fat attenuation. In addition to the above described pathoneumonic CT signs, visualisation of a gas filled appendix has been described as a finding associated with caecal dilatation from caecal volvulus(41).

Caecal volvulus is an emergency pathology, requiring surgical treatment. Right hemicolectomy is the treatment of choice in presence of caecal gangrene. There is considerable controversy regarding the preferred operative management of caecal volvulus in the absence of gangrenous bowel.

In presence of such diversion of opinions Consorti et al suggested that the most appropriate operative strategy for a given patient can be determined only by the operating surgeon after taking into consideration the surgical expertise, patient’s physiological status, viability of the involved intestines, the potential perioperative morbidity and mortality, and the risk of volvulus recurrence (2).

Reduction of the volvulus as the only operative maneuver without caecopexy was reported to be associated with high recurrence rate (42).

There is conflicting reports about the role of caecopexy in management of caecal volvulus in absence of bowel gangrene. While some authors (43) advocate resection for all cases of caecal volvulus regardless bowel viability, others (44) advocate that resection should not be performed in patients with viable colon as it has been associated with a twofold mortality rate and increased morbidity as compared with caecopexy.

O'Mara et al (9) performed caecopexy in 18 patients in their series with no operative deaths and a low rate of postoperative complications. There were no recurrences among the all the 18 patients during a follow-up that averaged 4.8 years. They concluded that caecopexy can be

performed safely and quickly without opening the bowel and with low recurrence rate. Their result support caecopexy as the preferred operative management of caecal volvulus. Resection appears to be unnecessary except in the presence of gangrenous bowel.

SUMMARY

Caecal volvulus is axial twisting that occurs involving the caecum, terminal ileum and ascending colon. It is a rare, but potentially fatal, cause of intestinal obstruction. Caecal volvulus after cesarean section is very rare, with only three cases of ileo-caecal volvulus reported in the English literature in the last two decades.

Preoperative diagnosis depends on high index of suspicion as most patients with acute caecal volvulus present with clinical pictures suggestive of intestinal obstruction. Diagnosis can be made by plain abdominal X-ray in more than half the cases. The increasing use of CT scan in investigating patients with acute abdomen allowed more and more cases of caecal volvulus to be diagnosed pre-operatively. The three CT findings associated with acute caecal volvulus ‘‘coffee bean’’, ‘‘bird beak’’, and ‘‘whirl’’ signs are pathoneumonic. Caecal volvulus is an emergency pathology, requiring urgent surgical treatment. Right hemicolectomy is the treatment of choice in presence of caecal gangrene. There is considerable controversy regarding the preferred operative management of caecal volvulus in the absence of gangrenous bowel. Opinions were divided between right hemicolectomy and caecopexy. We believe that surgical decision should be based on many factors including patient’s general condition, presence or absence of bowel gangrene, associated co morbid factors and the potential perioperative morbidity and mortality.

References

1. Grover NK, Gulati SM, Tagore NK, Taneja OP. Volvulus of the cecum and ascending colon. *Am J Surg* 1973;125:672-5.
2. Consorti ET, Liu TH. Diagnosis and treatment of caecal volvulus. *Postgrad Med J* 2005 Dec; 81(962):772-6.
3. Krippaehne WW, Velto RM, Jenkins CC. Volvulus of the ascending colon. *Am J Surg* 1967; 114:323-32.
4. Meyers JR, Heifetz C J, Baue AE. Cecal volvulus. *Arch Surg*1972; 104:594-9.
5. Nelson TG, Bowers WF. Volvulus of the cecum and sigmoid colon. *Arch Surg.* 1956; 72:469-78.
6. Villet D. Volvulus of the proximal colon. *Ann Surg.* 1959; 150:1075-85.
7. Dixon CF, Meyer AC. Volvulus of the cecmn. *Surg Clin North Am* 1948; 28:953-63.
8. Neil DA, Reasebeck PG, Reasbeck JC, Effney DJ. Caecal volvulus: ten year experience in an Australian teaching hospital. *Ann R Coll Surg Engl.* 1987 Nov; 69(6):283-5.

9. O'Mara CS, Wilson TH Jr, Stonesifer GL, Stonesifer GL, Cameron JL. Cecal volvulus: analysis of 50 patients with long-term follow-up. *Ann Surg.* 1979 Jun; 189(6):724-31.
10. Burke JB, Ballantyne GH. Caecal volvulus, Low mortality at a city hospital. *Dis Colon Rectum* 1984; 27:737-40
11. Anderson JR, Lee D. Acute caecal volvulus. *Br J Surg.* 1980 Jan; 67(1):39-41.
12. JORDAN GL Jr, BEAHRS OH. Volvulus of the cecum as a postoperative complication; report of six cases. *Ann Surg.* 1953 Feb; 137(2):245-9.
13. Ulloa SA, Ramírez LO, Ortíz VN. Cecal volvulus after laparoscopic liver biopsy. *Bol Asoc Med P R.* 1997 Oct-Dec;89(10-12):195-6.
14. Konvolinka CW, Moore RA, Bajwa K. Cecal volvulus causing postoperative intestinal obstruction: report of a case. *Dis Colon Rectum* 2001;44:893-5.
15. McIntosh SA, Ravichandran D, Wilmsink AB, Baker A, Purushotham AD. Cecal volvulus occurring after laparoscopic appendectomy. *JLS* 2001;5:317-8.
16. Guerra EE, Nghiem DD. Posttransplant cecal volvulus. *Transplantation* 1990;50:721.
17. Ali Khan S, Desai PG, Siddharth P, Smith N. Volvulus of cecum following simple nephrectomy. *Urol Int* 1985;40:1-2.
18. Etheredge EE, Martz M, Anderson CB. Volvulus of the cecum following transplant donor nephrectomy. *Surgery* 1977;82:764-7.
19. Scott CD, Trotta BM, Dubose JJ, Ledesma E, Friel CM. A cruel twist: post-operative cecal volvulus. *Ulus Travma Acil Cerrahi Derg.* 2008 Apr; 14(2):158-62.
20. Howard RG, Catto J. Caecal volvulus. A case of nonresectional therapy. *Arch Surg* 1980; 115:273-7.
21. Todd GJ, Forde KA. Volvulus of the caecum: Choice of operation. *Am J Surg* 1979; 138:632-4.
22. Morris DM, Eisenstat T, Hall GM. Management of caecal volvulus in debilitated patients. *South Med J* 1982; 75:1069- 71.
23. Al Salamah SM, El Keyali AY. Ileo-caecal volvulus post-cesarean section: A case report. *Saudi J Gastroenterol* 2000;6:163-4
24. James Fanning, Do and C Bernard Cross, MD. Post cesarean section caecal volvulus. *Am J Obstet Gynecol* 1988;158:1200-2.
25. Pratt AT, Donaldson RC, Everton LR, Yon JL. Caecal volvulus in pregnancy. *Obstet Gynecol* 1981;57:37-9.
26. Marren A, Wong K. Caecal volvulus associated with intestinal malrotation immediately following caesarean section. *N Z Med J.* 2006 Aug 18; 119
27. Lopez Carral JM, Esen UI, Chandrashekar MV, Rogers IM, Olajide F. Volvulus of the right colon in pregnancy. *Int J Clin Pract* 1998 Jun; 52(4):270-1.
28. Alinovi V, Herzberg P, Yannopoulos D, et al. Cecal volvulus following caesarean section. *Obstet Gynecol.* 1980; 55:131-4.
29. Pal A, Corbett E, Mahadevan N. Caecal volvulus secondary to malrotation presenting after caesarean section *J Obstet Gynaecol.* 2005; 25:805-20.
30. Ferguson L, Higgs Z, Brown S, McCarter D, McKay C. Intestinal volvulus following laparoscopic surgery: a literature review and case report. *J Laparoendosc Adv Surg Tech A.* 2008 Jun;18 (3):405-10.
31. Agahi A, Harle R. A serious but rare complication of laparoscopic adjustable gastric banding: bowel obstruction due to caecal volvulus. *Obes Surg.* 2009 Aug; 19(8):1197-200.
32. Wales L, Tysome J, Menon R, Habib N, Navarra G. Caecal volvulus following laparoscopy-assisted sigmoid colectomy for sigmoid volvulus. *Int J Colorectal Dis.* 2003 Nov;18(6):529-32.
33. Anderson JR, Mills JO. Caecal volvulus: a frequently missed diagnosis? *Clin Radiol.* 1984 Jan; 35(1):65-9.
34. [Cecal volvulus]. *J Chir (Paris).* 2005 Jul-Aug; 142(4):220-4.
35. Rabinovici R, Simansky DA, Kaplan O, et al. Cecal volvulus. *Dis Colon Rectum* 1990; 33:765-9.
36. Ellis H. Pseudo-obstruction of the colon. In: Ellis H, editor. *Intestinal obstruction.* NY: Appleton-Century-Crofts; 1982. p. 81-6.
37. Ghazi, A., Shinya, H. and Wolff, W. I.: Treatment of Volvulus of the Colon by Colonoscopy. *Ann. Surg.,* 183:263, 1976.
38. Delabrousse E, Sarliève P, Sailley N, Aubry S, Kastler BA. Cecal volvulus: CT findings and correlation with pathophysiology. *Emerg Radiol.* 2007 Nov; 14(6):411-5.
39. Perrin RS, Kunberger LE. Cecal volvulus. *AJR* 1998;171:860
40. Carolyn J. Moore, Frank M. Corl, Elliot K. Fishman, CT of Cecal Volvulus Unraveling the Image. *AJR* 2001; 177:95-98
41. Moore CJ, Corl FM, Fishman EK. CT of cecal volvulus: unraveling of the image. *AJR* 2001;177:95-8.
42. Inberg, M. V., Havia, T., Davidsson, L. and Salo, M.: Acute Intestinal Volvulus: A Report of 238 Cases, *Scand. J. Gastroenterol* 7:209, 1972.
43. Rivas AA, Dennison HC. Volvulus of the caecum. *Am Surg* 1978;44:332-8.
44. Tejler G, Jibom H. Volvulus of the caecum: Report of 26 cases and review of the literature. *Dis Colon Rectum* 1988;31:445-9.

Author Information

Fatima Hassan Mohamed, MBBS, ARAB BOARD

Associate Consultant Obstetrics & Gynecology, Department of Surgical Specialties, King Fahad Medical City

Muhammad Abukhater, MBBS, FRCSI

Assistant Consultant General Surgery, Department of Surgical Specialties, King Fahad Medical City

Basher A Ahmad, MBBS, Arab Board

Assistant Consultant, Radiology Department, King Fahad Medical City

Abbas AR Mohamed, MBBS, FRCSI, FICS

Consultant General and Laparoscopic Surgeon, Department of Surgical Specialties, King Fahad Medical City