

Anesthesia for Interventional Neuroradiology: Part III: Anesthetic goals and techniques

A Rosas

Citation

A Rosas. *Anesthesia for Interventional Neuroradiology: Part III: Anesthetic goals and techniques*. The Internet Journal of Anesthesiology. 1996 Volume 1 Number 3.

Abstract

ANESTHETIC GOALS AND TECHNIQUES

The goals of anesthetic management for INR procedures are:

1. To safely render a patient immobile, painless and comfortable.
2. To provide the neuroradiologist with optimum hemodynamic and intracerebral conditions for successful performance of the procedure.
3. To allow for perioperative evaluation of neurologic function
4. To assist in treating complications

The two basic techniques are general anesthesia and monitored intravenous sedation. There are advantages and problems associated with each technique, and different centers employ various approaches successfully.

The choice of anesthetic technique is dictated not only by the physiologic status of the patient, and the duration and type of interventional procedure, but also by the particular experience of each interventional neuroradiology team.

INTRAVENOUS SEDATION

The demands placed on the anesthesiologist by an interventional neuroradiology (INR) procedure are much greater than those of simple diagnostic angiography.⁽⁵⁷⁾ The anesthesiologist must keep the patient immobile and calm, sometimes for many hours, and manage hemodynamics, cerebral perfusion, and coagulation; yet, on demand, the patient may need to be responsive enough for a neurologic examination. In selected procedures with cooperative patients, managed anesthesia care with intravenous sedation (MAC) can provide excellent conditions for performance of endovascular interventional procedures. Its main advantage

is the ease with which an awake neurologic examination can be carried out during the procedure. There are also significant disadvantages associated with this approach:

Advantages of Intravenous sedation:

1. Easier to perform neurologic testing repeatedly
2. No hemodynamic changes associated with intubation or emergence

Potential disadvantages:

1. Poor control of the airway, with potential for:
 - Hypoxemia
 - Hypercapnia
 - Stertorous breathing and movement
2. Side effects of IV sedatives
 - Dysphoria
 - Prolonged somnolence
 - Extrapyrmidal symptoms
3. Poor tolerance of induced hypotension in awake patient.
4. Nausea and vomiting
5. Vagal reactions
6. Delays and complications if neurologic emergency occurs
 - Need to intubate acutely
 - Hemodynamic changes at wrong time

7. No protection from aspiration

8. May need to interrupt the procedure and induce general anesthesia if the patient is unable to tolerate sedation.

INDICATIONS FOR MAC

There are specific procedures during which the patient must be fully awake, with minimal or no sedation:

1.- Test occlusions of major cerebral vessels:(58)

When a major vascular supply to the brain must be permanently or temporarily interrupted, a test occlusion of the vessel is performed to assess the adequacy of collateral flow and the consequences of occlusion. The patient must be awake because an ongoing neurologic evaluation is of primary importance to assess neurologic integrity during the period of occlusion.

2.- Performance of the Wada-Rasmussen test:(59)

Amobarbital, a short acting barbiturate, is directly injected into the carotid circulation while the patient undergoes electrophysiologic and neurobehavioral exams in this test used to determine cerebral dominance, cognitive, and language function, and to locate seizure foci prior to surgical ablation. Usually, no premedication or sedation is given during this procedure because it interferes with interpretation of the test.

RELATIVE INDICATIONS FOR MAC

These procedures are usually performed under MAC in our institution:

1.- Simple procedures:

When they are of short duration, the patient is amenable and absolute immobility is not crucial (e.i., diagnostic angiography)

2.- Intra-arterial chemotherapy of brain tumors with cis-platinum:

This procedure is essentially the same as a diagnostic angiogram, except that a catheter is placed in a branch of the carotid or vertebral system to infuse chemotherapy in direct proximity to an intracranial tumor. The procedure typically lasts two to three hours, but absolute immobility is not crucial.

3.- Preoperative embolization of intracranial and extracranial tumors or, arterio-venous malformations which are perfused

by the external carotid circulation.

These procedures are usually of short duration, do not often require hemodynamic management, and although sometimes painful, are usually well tolerated with MAC. However, in patients with large extracerebral to intracerebral vascular anastomosis, embolic material can pass into the brain from the extracerebral circulation. In these cases, as discussed before, Young et al advocate increasing cerebral blood flow, by increasing the PaCO₂, to establish a pressure gradient that opposes intracerebral embolization. Hypercapnia is produced by adding CO₂ to the circuit, or by decreasing the minute ventilation. Control of ventilation for production of hypercapnia is accomplished best with general anesthesia

4.- Embolization of intracranial arterio-venous malformations (AVM's):

This is a challenging procedure for both neuroradiologist and anesthesiologist. It had previously been performed under MAC to evaluate the patient's neurologic status during the procedure. However, in our institution, general anesthesia has superseded MAC for these cases because our team feels that the advantages of general anesthesia outweigh those of MAC. Superselective flow-guided catheterization of the feeding vessels of the AVM, can place the tip of the embolizing catheter in the nidus of the AVM. Using liquid adhesive with the catheter in this position, the deposition of embolic material can be precisely controlled, such that the possibility of compromising normal neural tissue is decreased.

Placement of the catheter in this position requires a motionless patient, and hemodynamic management by the anesthesiologist.

The hemodynamic and respiratory management is designed to help the radiologist "float" the superselective catheter into position and to decrease the possibility of neurologic damage. We feel that controlled hypotension or induced hypertension with PaCO₂-related adjustments of cerebral blood flow and cerebral vascular tone are best managed with the patient under general anesthesia. The complications associated with MAC, such as nausea and vomiting during the critical part of the procedure, are thus avoided. Our colleagues in neuroradiology also feel that awake neurologic assessment is no longer of prime importance because they are able to control the deposition of embolic material more exactly with the conditions provided by general anesthesia.(60) However, if the lesion cannot be reached

with a superselective catheter, or embolization with particulate material must be carried out from a more distal location, the procedure is more safely carried out with MAC.

This allows for performance of a superselective anesthesia functional examination (SAFE),(61) (62) which helps to decide whether the catheter is positioned in a hazardous position, proximal to a vessel irrigating expressive regions of the brain, or if the area can be embolized without neurologic sequelae. At TMH, SAFE is performed with minimal or no sedation. A neurophysiologic exam with EEG is performed as a baseline, followed by sodium amytal injection via the superselective catheter into the vessel. After injection the neurophysiologic exam is repeated to evaluate the area of irrigation.

Thus, the choice of anesthesia, general or MAC, depends not only on the procedure and the patient, but also on the vascular anatomy, the type of embolization material and catheter system used by the neuroradiologist.

TECHNIQUES FOR MAC

There are many “recipes” for sedation.(63) (64) (65) The following are two approaches used at our institution:

1) Neuroleptic analgesia:

This technique offers analgesia, amnesia, and cardiovascular stability. In addition, it decreases cerebral blood flow, and cerebral metabolic rate. It usually renders a quiet cooperative patient, with good airway control, and little nausea. However, excessive sedation with airway obstruction, dysphoria, and extrapyramidal movements can occur. An infusion is prepared and used as follows:

NEUROLEPTIC INFUSION SOLUTION

10 ml of fentanyl + 10 ml of droperidol mixed with D5W to make 250 ml

(10 micrograms/ml droperidol + 2 micrograms fentanyl/ml)

Suggested use:

a.- Start with 50 ml bolus, then proceed with insertion of arterial line and bladder catheters

b.- Infuse at 25ml/Hr, titrating to patient needs

2) Propofol infusion:

Propofol should be combined with fentanyl and/or midazolam for adequate amnesia. If used by itself, propofol

can cause patients to wake up and move unexpectedly as the infusion is titrated down. Patients can also develop sudden airway obstruction, (usually at a critical point in the procedure !) and one may need to insert an airway, which will usually cause the patient to move or cough.

SUGGESTED USE

a.- Begin with fentanyl, 2 - 3 micrograms/Kg IV, then add midazolam 2 - 3 mg IV, and droperidol 1.25 mg IV

b.- Insert arterial line under local anesthesia

c.- Propofol, 40 - 60 mg IV bolus can be given for bladder catheterization, and femoral arterial puncture.

d.- Propofol drip: Starting at 20 micrograms/Kg/min, titrating until patient sedated.

GENERAL ANESTHESIA

There are many advantages to this technique. With newer induction and inhalation agents, general anesthesia can be rapidly induced with minimum hemodynamic changes, the depth readily controlled, and a smooth and rapid emergence obtained.(66) (67) Under general anesthesia, PaCO₂ can be regulated, and management of induced hypotension is facilitated. Respiratory immobility is essential with certain imaging techniques; when apnea is requested by the neuroradiologist it can be induced at will with general anesthesia. When a neurologic emergency such as thrombosis or hemorrhage occurs, no time is lost in securing the airway and the anesthesiologist can immediately begin to control hemodynamics.

The main disadvantage of general anesthesia is that awake neurologic assessments cannot be readily performed, but as discussed above, we feel this is only of prime importance in selected cases. Other disadvantages are that endotracheal intubation can cause hypertension, while extubation can lead to straining, airway obstruction, and coughing. However, with careful anesthetic management, these disadvantages can be minimized.

TECHNIQUE

We do not give sedatives or narcotics if intraoperative neurophysiologic assessment with somatosensory- evoked potentials (SSEP), or electroencephalography (EEG) is contemplated until baseline studies are obtained and the anesthetic plan has been discussed with the neurophysiologist.

We obtain peripheral intravenous access with a single 16 or

18 gauge IV catheter and a multiple port extension, preferably on the arm nearest to the anesthetist. This is usually adequate, unless the neuroradiologist feels that there is a high probability of serious complications. In such a case, two peripheral IV's or a central line and a peripheral IV are inserted. Particular attention must be given to fluid balance since large amounts of irrigating solutions and contrast media are often used by the neuroradiologist. A urinary catheter is always inserted, either after induction of anesthesia or under sedation. Glucose-containing solutions are avoided because of their deleterious effect on brain ischemia. (68) In diabetics, we strictly control blood sugar levels perioperatively, with intravenous insulin, to maintain blood sugar levels between 150 and 200 mg percent.

After non-invasive monitoring is established, we induce anesthesia with propofol, 2 - 2.5 mg/kg, or etomidate, 0.2 - 0.6 mg/kg and administer esmolol, 0.5 to 1 mg/kg, or lidocaine 1 mg/kg, prior to intubation to avoid the sympathetic response to intubation. If a radial arterial line is indicated, it is usually inserted after induction, prior to intubation. A control activated clotting time (ACT) is obtained after insertion of the arterial line. We do not use eye lubricant in case a fundoscopic exam is necessary during the procedure. Central venous lines or pulmonary artery catheters are inserted via the subclavian route at this time if indicated.

For airway management, we prefer a low profile RAE endotracheal tube, or a regular endotracheal tube with the extraoral end cut short. Cutting the tube short keeps the extraoral end from being pushed in or kinked by the image intensifier, which moves close to the patient's face. We have recently used a laryngeal mask airway (LMA) for airway management. This device allows for control of the airway with minimal stimulation, hypertension, or cough, and a smooth emergence under light general anesthesia. (69) (70) In our experience, however, the head movement that sometimes occurs when using an LMA with spontaneous ventilation has impaired radiographic imaging; and the PaCO₂ levels have been higher than ideal. Muscle relaxants and controlled ventilation can be used with an LMA, and thus avoid these problems, but only as long as airway pressures are not greater than 20 cm H₂O. (71) More experience is needed in order to recommend routine use of LMA's in this setting.

For maintenance of anesthesia we use a continuous infusion of propofol plus low inspired concentrations of isoflurane or desflurane. Inhalation anesthetics allow rapid changes of

anesthetic depth and blood pressure, but increase ICP with concentrations higher than 1 MAC and normocapnia. (72) We do not advocate the routine use of nitrous oxide for these cases since nitrous oxide increases cerebral blood flow from 60 to 100 percent; and can increase ICP to a variable degree. (73) Nitrous oxide could also theoretically enlarge air bubbles that may be accidentally introduced into the arterial circulation by the radiologist. (74) Minimal amounts of intravenous narcotics should be used because these procedures are not painful, and excessive doses of narcotics will only delay emergence. The intraoperative management of PaCO₂, controlled hypotension, and induced hypertension will be discussed in more detail later. The blood pressure should be controlled during emergence and after the procedure.

After assessing the reversibility of the muscle relaxant, we extubate the patient while still under deep anesthesia, reverse the muscle relaxants, and wake the patient breathing oxygen via face mask. If an LMA was used, the patient is allowed to wake up with the device in place, to be removed once the patient responds to commands. This awake extubation with an LMA is remarkably well tolerated, with little cough, straining, or hypertension.

Postoperatively, the mean arterial pressure (MAP) should be kept 10 - 20 percent lower than the patient's basal levels. Acute elevations of blood pressure can be treated with esmolol, 1mg/kg; propofol, in bolus doses of 30 - 50 mg IV; vasodilators; or small doses of narcotics, depending on etiology.

References

- r-0. 1. Hilal SK, Sane P, Mawad ME, Michelsen WJ: Therapeutic interventional radiological procedures in neuroradiology, Angiography. 3rd edition. Edited by Abrams H. Boston, Little Brown, 1983
- r-1. 2. Brown MM: Surgery, angioplasty, and interventional neuroradiology. *Curr Opin Neurol Neurosurg* 6:66-73, 1993
- r-2. 3. Barnwell SL: *Interventional neuroradiology*. West J Med 158:162-170, 1993
- r-3. 4. Higashida RT, Hieshima GB, Halbach V: Advances in the treatment of complex cerebrovascular disorders by interventional neurovascular techniques. *Circulation* 83 (sup):I-196, 1991
- r-4. 5. Moret J, Picard L: Endovascular therapy of intracranial arteriovenous malformations; Results in 242 patients. Paper presented at the World Interventional Neuroradiological society, Val d'Isere, France, Jan., 1989
- r-5. 6. Eichhorn JH, Cooper JB, Cullen DJ et al: Anesthesia practice standards at Harvard: a review. *J Clin Anes* 1:55, 1988
- r-6. 7. Eichhorn JH, Cooper JB, Cullen DJ et al: Standards for patient monitoring during anesthesia at Harvard Medical School, *JAMA* 256:1017, 1986
- r-7. 8. Burk NS: Anesthesia for magnetic resonance imaging,

- Anesth Clin North Am 7:(3)707, 1989
- r-8. 9. Peer Review in Anesthesiology-American Society of Anesthesiologists, Park Ridge, Ill, 1989, ASA
- r-9. 10. Eichhorn JH, Cooper JB, Cullen DJ et al: Standards for patient monitoring during anesthesia at Harvard Medical School, JAMA 256:1017, 1986
- r-10. 11. Eichhorn JH: Prevention of intraoperative anesthesia accidents and related severe injury through safety monitoring, Anesthesiology 70:572, 1989
- r-11. 12. Kanal E, Shellock F, SMRI Safety Committee: Policies, guidelines, and recommendations for MR imaging safety and patient management, J Magnet Res Imaging 2:247 1992
- r-12. 13. Peden C, Menon D, Hall A et al: Magnetic resonance for the anaesthetist. Part II. Anaesthesia and monitoring in MR units. Anesthesia 47:508 1992
- r-13. 14. Gravenstein JS, Paulus DA: Clinical Monitoring practice, ed 2, Philadelphia, 1987, JB Lippincott.
- r-14. 15. Slogoff S, Keats AS, and Arlund C: On the safety of radial artery cannulation, Anesthesiology 59:42, 1983.
- r-15. 16. Young WL, Pile-Spellman J: Anesthetic considerations for interventional neuroradiology. Anesthesiology 80:427-455, 1994
- r-16. 17. Duckweiler GR, Dion J, Vinuela F, Jabour B, Martin N, Bentson J: Intravascular microcatheter pressure monitoring: Experimental results and early clinical evaluation. American J Neuroradiol 11:169-175, 1990
- r-17. 18. Gorbach MS: Considerations in the interpretation of systemic pressure monitoring p 296 In Lumb PD, Bryan-Brown CW (eds): Complications in Critical Care Medicine. Year Book Medical Publishers, Chicago 1988
- r-18. 19. Swedlow DB, Irving SM: Monitoring and patient safety. In Blitt CD, ed: Monitoring in anesthesia and critical care medicine, ed 2, New York, 1990, Churchill-Livingstone
- r-19. 20. Plum F, Posner JB: The diagnosis of stupor and coma, ed 3, Philadelphia, 1980, FA Davis
- r-20. 21. Aidinis SJ, Zimmerman RA, Shapiro HM, Bilanink LT, et al: Anesthesia for brain computed tomography. Anesthesiology 44:420, 1976
- r-21. 22. Frank SM, Parker SD, Rock P, et al: Moderate hypothermia with partial bypass and segmental repair for thoracoabdominal aortic aneurysm. J Vasc Surg 19:687-697, 1994
- r-22. 23. Busto R, Dietrich WD, Globus MY, et al: Small differences in intraschismic brain temperature critically determine the extent of ischemic neuronal injury. J Cereb BI Flow Met 7:729-738, 1987
- r-23. 24. Sise MJ, Hollingsworth P, Brimm JE: Complications of the flow-directed pulmonary catheter: a prospective analysis in 219 patients. Crit Care Med 9:315 1981
- r-24. 25. Shah KB, Rao TLK, Laughlin S, El-Etr AA: A review of pulmonary artery catheterization in 6,245 patients. Anesthesiology 61:271 1984
- r-25. 26. Harper R, Grossman R, personal communication: Rosas A
- r-26. 27. Shapiro HM, Drummond JC: Neurosurgical Anesthesia: in Miller RD, Cuchiara RF, Miller ED, et al. Anesthesia 4e Churchill Livingstone, New York 1994
- r-27. 28. Swan HJC, Ganz W, Forrester JS et al: Catheterization of the heart in man with use of a flow-directed-balloon-tipped catheter, N Engl J Med 283:477, 1970
- r-28. 29. Kassell NF, Peerles SJ, Durward QJ et al: Treatment of ischemic deficits from vasospasm with intravascular volume expansion and induced arterial hypertension, Neurosurgery 11:337, 1982.
- r-29. 30. Earnest F, Forbes G, Sandok BA et al: Complications of cerebral angiography: Prospective assessment of risk. AJR 142:247 1984
- r-30. 31. Faught E, Trader SD, Hanna GR: Cerebral complications of angiography for transient ischemia and stroke: prediction of risk. Neurology 29:4 1979.
- r-31. 32. Greenberger PA: Contrast media reactions, J Allergy Clin Immunol 74:600, 1984
- r-32. 33. Goldberg M: systemic reactions to intravascular contrast media: A guide for the anesthesiologist. Anesthesiology 60:46-56, 1984
- r-33. 34. Caro JJ, Trinidale E, McGregor M: The cost effectiveness of replacing high osmolality with low osmolality contrast media. r-34. Am J Roentgenol 159; 869-874, 1992.
- r-35. 35. Goldberg M: systemic reactions to intravascular contrast media: A guide for the anesthesiologist. Anesthesiology 60:46-56, 1984
- r-36. 36. Horrow JC: Protamine: A review of its toxicity. Anesth Analg 64:348 1985
- r-37. 37. Lowenstein E, Johnston WE, Lappas DG et al. Catastrophic pulmonary vasoconstriction associated with protamine reversal of heparin. Anesthesiology 59:470 1983
- r-38. 38. Weiss ME, Adkinson NF jr: Allergy to Protamine. in Vervloet D, ed: Clinical reviews in allergy:anesthesiology and allergy, New Brunswick, NJ, 1991, Humana Press.
- r-39. 39. Vezina D, Sheridan P, Blain R, Roberts KD, Bleau G: Safety of protamine sulfate administration in vasectomized men. Contraception 41:605-616, 1990
- r-40. 40. Pharo GH, Horrow J, Van Riper DF, Levy JH: Suspected protamine allergy : diagnosis and management for coronary artery surgery. Anesth Analg 78:181-184, 1994
- r-41. 41. Murray DJ: Evaluation of the patient with anemia and coagulation disorders. In Rogers MC, Tinker JH, Covino BG, Longnecker DE, Principles and practice of Anesthesiology. r-42. St. Louis, Mosby, 1993
- r-43. 42. Keefe D, Frishman WH: Clinical pharmacology of the calcium-channel blocking drugs. In Packer M, Frishman WH, eds: Calcium channel antagonists in cardiovascular disease, Norwalk, Conn,1984, Appleton-Century-Crofts.
- r-44. 43. Ryhanen P,Hollman A, and Horttonen L: Blood pressure changes during and after anesthesia in treated and untreated hypertensive patients, Ann Chir Gyn 67:180, 1978.
- r-45. 44. Sieber FE, Toung TJK: Hyperglycemia and stroke outcome following carotid endarterectomy. Anesthesiology 71: A1136, 1989.
- r-46. 45. Berger L, Hakim AM: The association of hyperglycemia with cerebral edema in stroke . Stroke 17:865, 1986.
- r-47. 46. Sieber FE et al: Hypoglycemia and cerebral autoregulation, Anesthesiology 71:A605, 1989.
- r-48. 47. Goldman L, Caldera DL: Risks of general anesthesia and elective operation in the hypertensive patient. Anesthesiology 50:285, 1979.
- r-49. 48. Edvinson L, Owman C, and Siesjo B: Physiological role of cerebrovascular nerves in the autoregulation of cerebral blood flow. Brain Res. 117:518, 1976.
- r-50. 49. Hunt WE, Hess RM: Surgical risk as related to time of intervention in the repair of intracranial aneurysms, J Neurosurg 28:14, 1968.
- r-51. 50. Kirsch JK, Diring MN: Evaluation of the patient with neurologic disease. In Rogers MC, Tinker JH, Covino BG, et al.,Eds. Principles and Practice of Anesthesiology, St. Louis, 1993, Mosby.
- r-52. 51. Stoelting RK: Psychological preparation and preoperative medication. In Miller RD, ed: Anesthesia, New York, 1981, Churchill-Livingstone.
- r-53. 52. Lyons SM, Clarke RSJ, and Vulgaraki K: The

- premedication of cardiac surgical patients, *Anaesthesia* 30:459, 1975.
- r-54. 53. Reves JG, Frazen RJ, Vinik HR et al: Midazolam: Pharmacology and uses, *Anesthesiology* 54:66, 1981.
- r-55. 54. Manchikanti L, Kraus JW, and Edds SP: Cimetidine and related drugs in anesthesia, *Anesth Analg* 61:595, 1982.
- r-56. 55. Stoelting RK: Responses to atropine, glycopyrrolate and Riopan on gastric fluid pH and volume in adult patients, *Anesthesiology* 48:367, 1978.
- r-57. 56. O'Sullivan G, Sear JW, Bullingham RE, et al: The effect of magnesium trisilicate, metoclopramide and ranitidine on gastric pH, volume, and serum gastrin, *Anaesthesia* 40:246, 1985.
- r-58. 57. Young WL, Pile -Spellman J: Anesthetic considerations for interventional neuroradiology. *Anesthesiology* 80:427-456.
- r-59. 58. Anon VV, Aymard A, Gobin YP, et al: Balloon occlusion of the internal carotid artery in 40 cases of giant intracavernous aneurysm: Technical aspects, cerebral monitoring, and results. *Neuroradiology* 34:245-251, 1992.
- r-60. 59. Wada J, Rassmussen T: Intracarotid injection of sodium amytal for the lateralization of cerebral speech dominance: Experimental and clinical observations. *J Neurosurg* 17:266-282, 1960.
- r-61. 60. Mawad ME, Chi L: Personal communication.
- r-62. 61. Purdy PD, Batjer HH, Samson D, et al: Intraarterial sodium amytal administration to guide preoperative embolization of cerebral arteriovenous malformations. *J Neurosurg Anesth* 3:103-106, 1991.
- r-63. 62. Rauch RA, Vinuela F, Dion J, et al: Preembolization functional evaluation in brain arteriovenous malformations: The ability of superselective amytal test to predict neurologic dysfunction before embolization. *AJNR Am J Neuroradiol* 13:309-314, 1992.
- r-64. 63. Eskridge JM: Interventional neuroradiology. *Radiology* 172: 991-1006, 1989.
- r-65. 64. O'Mahoney BJ, Bolsin SNC: Anesthesia for crossed embolization of cerebral arteriovenous malformations. *Anesth Intensive Care* 16: 318-323.
- r-66. 65. Glauber DT, Audenart SM: Anesthesia for children undergoing craniospinal radiotherapy. *Anesthesiology* 67: 801-803, 1987.
- r-67. 66. Van Hemelrijck J, Gonzales JM, White PF: Pharmacology of intravenous anesthetic agents. in Rogers MC, Covino BG, et al: Principles and practice of anesthesiology, St Louis, 1993, Mosby.
- r-68. 67. Longnecker DE, Miller FL: Pharmacology of inhalational anesthetics in Rogers MC, Covino BJ, et al, eds: Principles and practice of anesthesiology, St Louis, 1993, Mosby.
- r-69. 68. Longstreth WT, Inui TS: High blood glucose level on hospital admission and poor neurological recovery after cardiac arrest. *Ann Neurology* 15:59, 1984.
- r-70. 69. Brain AIJ, The laryngeal mask-a new concept in airway management. *Br J Anaesth.* 1983;55:801-804.
- r-71. 70. Benumof JL. Laryngeal mask airway: indications and contraindications (editorial views). *Anesthesiology.* 1992;77:843-846.
- r-72. 71. Graziotte PJ: Intermittent positive pressure ventilation through a laryngeal mask airway. *Anaesthesia.* 1992;47:1088-1089.
- r-73. 72. Adams RW, Cucchiara RF, Gronert GA, et al: Isoflurane and cerebrospinal fluid pressure in neurosurgical patients, *Anesthesiology* 54:97, 1981.
- r-74. 73. Hansen TD, Warner DS, Todd MM, et al: Relative cerebral blood flow effects of nitrous oxide and volatile anesthetics, *Br J Anaesth* 63:290, 1989.
- r-75. 74. Losasso TJ et al: The "risk" of nitrous oxide in sitting neurosurgical patients: a randomized study, *Anesthesiology* 71:A1137, 1989.

Author Information

Alejandro L. Rosas, M.D.