

Attitudes and subjective reasons of medication compliance and noncompliance among outpatients with schizophrenia in India

R Baby, S Gupta, R Sagar

Citation

R Baby, S Gupta, R Sagar. *Attitudes and subjective reasons of medication compliance and noncompliance among outpatients with schizophrenia in India*. The Internet Journal of Epidemiology. 2008 Volume 7 Number 1.

Abstract

Objective: This study aims to assess the attitudes and reasons of medication compliance and noncompliance among patients with Schizophrenia attending outpatients in India and examine the prevalence and factors effecting noncompliance. **Method:** outpatients with schizophrenia (n=75) and their accompanying family members(n=75) were assessed using standardized tools which consist reasons of medication influences (ROMI) scale, Drug Attitude Inventory (DAI) scale, Positive and Negative Symptoms Scale (PANSS). They were also evaluated for sociodemographic details; illness related and drug related variables. **Results:** The prevalence of non-compliance in the selected setting is 38.7%. Majority of patients and family members had a positive attitude towards medication and treatment. Family member is able to identify the compliance status of the patient and the reasons for the noncompliance better than the patients. The reasons which are significantly correlating with compliance are 'perceived daily benefit from medication', 'positive relationship with psychiatrist', pressure from the family and health system', 'positive family belief towards illness and treatment'. Where as the significant reasons for noncompliance are 'no perceived daily benefit from medications', 'difficulty in gaining access to treatment and medications', financial obstacles', 'embarrassment or stigma related to treatment and medications', 'medicines currently not perceived as necessary'. The factors significantly associated with noncompliance include lower educational status, rural area of stay, adjustment difficulty with their family and spouse, previous history of non-compliance, poor insight into illness, higher positive PANSS score. The patients who gave history of hospitalization in the past are more likely to be compliant with their medication. **Conclusions:** The Findings strongly recommends the need to develop a standard protocol for carrying out adherence counseling to all patients. Also there is need to create a cadre for mental health nurse practitioner in outpatient and inpatients units to ensure that all patients and family members are given psycho education / adherence counseling before discharge and during follow-ups.

Original work done by the author for the partial fulfillment of the post graduation programme in psychiatric nursing.

consistent counseling to all patients before their discharge from inpatient unit and during their follow-up

Significant outcomes

- Majority of patients and their family member have positive attitude towards antipsychotic medications.
- Patients with higher positive symptoms, poor insight into the illness, rural background, lower educational status, adjustment difficulty with their family and spouse, previous history of non-compliance are more likely to be noncompliant with the medications
- Standard medication adherence counseling protocol need to be developed and used to ensure

Limitations

- Self report methods was used to assess medication compliance
- Smaller sample size, and sample of convenience was used
- Patients with irregular follow-ups was not assessed

INTRODUCTION

Medication compliance is one of the most difficult challenges in the management of schizophrenia in India. It is

estimated that the rate of noncompliance is about 50% during first year and 75% during second year, after the patients are discharged from inpatient care unit (1). Medication noncompliance is one of the main obstacles to control relapse in schizophrenia. It significantly increases the rate of relapse, rate and length of hospitalization, and the risk for hospitalization in the future (2). The over all compliance rates for antipsychotic medication was lower than that of antidepressants and other drugs (3).

The medication compliance in patients with schizophrenia had been predicted by the patient's subjective response to treatment and attitudes towards antipsychotics (4,5). Positive attitude of patients towards treatment can improve therapeutic alliance, medication compliance and long-term prognosis (6). Attitude towards antipsychotic medication is significantly affected by the factors such as lack of insight into the illness, presence of global functioning, increased severity of illness and side effects of medications (7).

Recent studies have shown that the Subjective reasons which are significantly related to medication compliance: 'perceived daily benefit, (8,9) 'positive relationship with the therapist', 'positive attitude of significant others' (9). The reasons identified for patient's non compliance are 'distressed by side effects' (8), 'lack of insight' 'lack of acceptance of treatment' (9), 'substance abuse' 'low self esteem', 'inability to identify the stressors of life', inability to identify the symptoms of relapse' (10). Although the burden of side effects is viewed as an important reasons of noncompliance, but studies have found no association between extra pyramidal symptoms and attitude towards antipsychotic medication (6). Its generally believed that atypical antipsychotics are better tolerated by the patients but studies have not found better compliance with these drugs or better effect on the subjective quality of life of patients (6,11,12).

Several studies were conducted to find out the significant predictors of medication noncompliance. These study results have showed that the factors which predict noncompliance are high level of positive symptoms, alcohol abuse (13,14,18),, previous noncompliance (13, 14, 19), shorter duration of illness (8) ,lack of insight(13,15), non affordability of drugs, education not given by doctor(16), young age(17), lower educational level(18), and co morbidity with personality disorder(19).

Majority of these studies are done in developed countries. Very few literatures are available from India and other

developing countries regarding the attitudes and reasons of medication compliance. The current study has tried to assess the attitudes and reasons for compliance as well as noncompliance among patients with schizophrenia, which can provide basis for planning effective interventional strategies to the mental health professionals for improving compliance of the patient in the future.

Aims of the study-To assess the attitudes and reasons of medication compliance and noncompliance among patients with Schizophrenia attending outpatient services in India

Objectives of the study-The objectives of the study are: i) to assess the prevalence of medication non compliance among patients with Schizophrenia, ii) to assess the attitudes towards and reasons of medication compliance and noncompliance, and iii) to identify the relationship of medication noncompliance with demographic and selected variables.

MATERIALS AND METHODS

Subjects Seventy five adult patients with schizophrenia and accompanying 75 family members attending the outpatient services of department of psychiatry, All India Institute of Medical Sciences (AIIMS), New Delhi, India from June to November 2006 of, were recruited for the study. Those patients with duration of illness less than 2 years and who are not prescribed any antipsychotics for the past 6 months were excluded from the study. All patients were diagnosed as schizophrenia by the treating psychiatrist as per ICD 10 diagnostic criteria (WHO1992) and are registered in the psychiatry department for the past two years.

DATA COLLECTION PROCEDURE

The ethics committee of AIIMS approved the study protocol. All the tools were tried out and pilot study was done prior to actual data collection and found feasible in the proposed setting and population.

Written informed consents were obtained from the patients and their family members after the purpose of the study and risk and benefits of participation had been explained.

Measures evaluated includes sociodemographic details (age of the patient and family member, gender, educational status, marital status, religion, monthly family income, type of family, place of stay, occupation, relationship with spouse/family), illness related variables (total duration of illness, family history, history of hospitalization, duration of

hospital stay, comorbid physical illness, history of substance misuse, insight into illness, severity of illness, frequency of follow up, cost of transport) and medication related variables (number of tablets taken per day, availability of drug, cost of monthly medication, supervision of medication, type of medication, side effects of medication).

Severity of illness was measured using positive and negative syndrome scale (PANSS)(20). It is a 30-item scale (has got a total score of- 210), positive and negative symptom subscale and general psychopathology subscale. The Interrater reliability was established (r=0.82). Subjective reasons of medication compliance/noncompliance were assessed using 20 item ROMI scale (21). The subject was asked to indicate the degree of influence of each item of the scale, i.e. reasons for compliance and reasons for noncompliance. The graduated responses to each are ‘no influence’, ‘mild influence’ and ‘strong influence’ (1, 2 and 3, respectively, or 9 when it is impossible to evaluate the degree of influence). The attitude towards antipsychotic medication was assessed using 30 item drug attitude inventory (DAI) . It is a self report questionnaire in which each item ticked ‘yes’ is rated +1 and items ticked ‘no’ is rated as -1 .The positive scores and negative scores are calculated and the total score will be positive score minus negative score. The positive total score indicates positive attitude and negative total score indicates negative attitude towards antipsychotic medications. DAI and ROMI were administered also to family member accompanying the patient for comparison.

STATISTICAL ANALYSIS

The statistical package for social sciences 10(SPSS Inc., Chicago,IL, USA) program was used for statistical analysis. Data were analyzed using Chi square test, Student ‘t’ test, Paired ‘t’ test, and Mann Whitney test.

RESULTS

Out of 75 patients in the study; 46 patients are compliant and 29 patients are noncompliant to the medication and treatment. Hence the prevalence of non-compliance to medication is 38.7%. The patients were defined as compliant, if they take their medications in right doses, frequency as prescribed by the treating psychiatrist more than 23days or 75% of days in the past one month and were defined as noncompliant if they do not take their medications in right doses, frequency as prescribed by the treating psychiatrist or misses the medication for more than seven days continuously or 25% of days in the past one

month.

DEMOGRAPHIC AND CLINICAL CHARACTERISTICS

As per the demographic and clinical characteristics are displayed in table 1, the variables that are statistically significant at the level of p>0.05 or below which are significantly related to medication noncompliance are educational status of patient (p=0.03*), from rural background (p=0.02*), adjustment difficulty with the family/spouse (p=0.02*), history of hospitalization (p=0.03*), level of insight into illness (P =0.001**), PANSS positive symptoms score (p=0.02*).

COMPARISON OF ATTITUDE TOWARDS MEDICATION AND TREATMENT USING DAI SCALE

The attitudes of patients towards medication and treatment as measured by drug attitude inventory (DAI) shown in table 2. The patients in the compliant and noncompliant groups have positive overall attitude mean scores. The mean positive attitude, negative attitude and overall attitude of patients towards medication are comparable among both the groups (p=0.2) majority of the patients have positive attitude towards medication irrespective of the compliant/ noncompliance status.

Figure 1

Table 2: Comparison of attitude towards medication between the Patient groups using DAI scale(n+75)

Variable	compliant (n=46)	noncompliant (n=29)	value
Positive sub score(mean)	20.24(SD=4.86)	18.34(SD=5.14)	student t'test t=1.06 p=0.2
Negative sub score(mean)	9.28(SD=4.58)	10.66(SD=5.14)	
Total attitude score(mean)	11.02(SD=9.21)	8.55(SD=10.6)	
Range	-14 to +30	-16 to +26	

The comparison of the attitudes of patients and family members towards medication and treatment as measured by drug attitude inventory (DAI) shown in table 3. even though majority of the patients and family members, positive attitude towards medication is higher in family member when compared with patient. On the contrary the patient is having more negative attitude towards medication when compared with their family member(p=0.001**).

Figure 2

Table 3: comparison of attitude towards medication between patient and family member (n=150)

Variable	patient(n=75)	family member(n=75)	value
Positive sub score(mean)	19.8 (SD=4.9)	23.0(SD=3.8)	paired t test t =5.1 p =0.001**
Negative sub score(mean)	10.0 (SD=4)	6.9(SD=3.8)	
Total attitude score(mean)	9.8(SD=9.21)	16.1(SD=7.6)	

Figure 3

Table 1: the demographic and clinical characteristics of complaint and noncompliant groups N=75

Characteristics	compliant (n=46)	noncompliant (n=29)	value
SOEIOGEMOGRAPHIC VARIABLES			
Mean age of the patient	30.8(SD=9.4)	32.1(SD=10.4)	t=0.5 p=0.5
Mean age of family member	32.7(SD=10.4)	42.9(SD=11.5)	t=1.06 p=0.2
Sex (male)	32(69.6%)	19(65.5%)	$\chi^2=0.1$ p=0.8
Education of patient (12 th standard & above)	31(67.4%)	14(31.2%)	$\chi^2=10.6$ p=0.03*
Education of family member (12 th standard & above)	26(56.5%)	17(55.1%)	$\chi^2=3.1$ p=0.6
Marital status(married)	19(41.3%)	16(55.2%)	$\chi^2=5.4$ p=0.06
Employment status (unemployed/lost job)	30(65.2%)	20(69%)	$\chi^2=1.1$ p=0.4
Religion (Hindu)	44(95.7%)	29(100%)	$\chi^2=5.5$ p=0.5
Monthly family income (Less than Rs 5000)	27(58.8%)	19(65.5%)	$\chi^2=1.2$ p=0.7
Type of family (joint family)	27(58.7%)	19(65.5%)	$\chi^2=0.3$ p=0.6
Place of stay (rural)	11(24.9%)	15(58.6%)	$\chi^2=6.07$ p=0.02*
Relationship with family/spouse (Adjustment difficulty)	11(24.9%)	15(58.6%)	$\chi^2=6.07$ p=0.02*
ILLNESS RELATED VARIABLES			
Median years of mental illness	5years	3years	z=0.6 p=0.5
Family history of mental illness (present)	17(37.6%)	10(27.6%)	$\chi^2=1.323$ p=0.72
Co morbid physical illness(present)	12(26.1%)	3(10.3%)	$\chi^2=2.9$ p=0.2
Duration of hospitalization (present)	24(52.2%)	8(27.6%)	$\chi^2=4.3$ p=0.03*
Duration of hospital stay (less than one month)	8(17.4%)	6(20.7%)	$\chi^2=0.1$ p=0.9
Substance misuse (present)	18(39.1%)	12(41.4%)	$\chi^2=0.03$ p=0.8
Frequency of follow up (monthly)	20(43.5%)	8(27.6%)	$\chi^2=2.2$ p=0.9
Median Cost of transport per visit (in Rs)	Rs100	Rs 200	Z=1.09 p=0.2
Insight into illness			
Grade3 and below	7(15.2%)	17(58.6%)	$\chi^2=25.5$ P =0.001**
Grade4 and above	39(84.8%)	12(41.4%)	
Severity of illness (PANSS)			
Positive symptoms score	14.5(SD=8.0)	24.2(SD=9.2)	t=2.3 p=0.02*
Negative symptoms score	19.6(SD=6.4)	20.8(SD=7.2)	t=0.7 p=0.4
General symptoms score	37.2(SD=12.8)	43.0(SD=14)	t=1.8 p=0.06
MEDICATION RELATED VARIABLES			
Mean Number of medicines/day	3.28(SD=2.41)	3.34(SD=2.19)	Z=0.3 p=0.75
Availability of drug (easily available)	28(60.9%)	17(58.6%)	$\chi^2=0.13$ p=0.93
Monthly cost of medicines(median)	Rs300	Rs 400	Z = 1.107 P= 0.2
Supervision of medication (present)	27(58.7%)	16(55.2%)	$\chi^2=0.2$ p=0.9
Side effects			
EPS	13(28.3%)	11(37.9%)	$\chi^2=2.242$ p=0.691
Hematological	2(4.3%)	3(17.3%)	$\chi^2=1.028$ p=0.311
Weight gain	18(39.0%)	11(37.9%)	$\chi^2=0.011$ p=0.917
GI disturbances	15(32.6%)	9(31.0%)	$\chi^2=0.020$ p=1.000
Cardiovascular effects	7(15.2%)	2(6.9%)	$\chi^2=1.166$ p=0.468
Sedation	25(54.3%)	15(51.7%)	$\chi^2=0.001$ p=0.979
Anticholinergic effects	16(34.8%)	10(34.5%)	$\chi^2=0.01$ p=0.9
Type of medication			
Atypical antipsychotics(oral)	5(10.8%)	8(27.5%)	
Typical antipsychotics(oral)	36(79.2%)	20(68.9%)	
Depot antipsychotics	3(6.5%)	2(6.8%)	
Clozapine	1(3.5%)	4(13.6%)	

COMPARISON OF REASONS OF MEDICATION COMPLIANCE AND NONCOMPLIANCE USING ROMI SCALE

REASONS OF COMPLIANCE

the responses to the ROMI open ended question(what is your main motivation for taking the medication?) were grouped into six closed questions in the ROMI scale which is described in table. 4. it shows that the major reasons of compliance which are significantly different among

compliant and noncompliant groups are 1) perceived daily benefits due to medicines (p=0.01*) 2) positive relationship with the treating psychiatrist(p=0.03*), 3) positive family belief towards illness and treatment(p=0.01*), 4) family and health systems pressure on the subjects for taking medications(p=0.05*). Whereas the reasons of relapse prevention and fear of rehospitalization, and are not significantly different among the groups and are comparable.

Figure 4

Table 4: Rating of patient's medication influence(ROMI) scale closed questions (reasons for compliance)(n=75)

Reasons of compliance	Compliant (46)			Noncompliant (29)			Value
	Degree of influence %			Degree of influence %			
	None	mild	Strong	None	mild	Strong	
Perceived day to day benefit	8.7	26.1	65.2	24.1	37.9	37.9	$\chi^2=6.1$ p=0.01*
Positive relationship with the physician	17.4	19.6	63.0	31.1	37.9	31.0	$\chi^2=6.4$ p=0.03*
Positive family belief	13.0	19.6	67.4	24.1	17.2	37.9	$\chi^2=7.9$ p=0.01*
Relapse prevention	19.6	19.6	60.8	44.8	17.2	58.6	$\chi^2=1.5$ p=0.4
Family/health system pressure	43.5	17.4	39.1	41.4	13.8	44.5	$\chi^2=5.7$ p=0.05*
Fear of re-hospitalization	47.8	19.6	32.6	41.4	34.5	24.1	$\chi^2=2.1$ p=0.3
Chi square test							
*p<0.05							
**P<0.008 after doing Bonferroni Post Hoc correction for multiple variables							

REASONS OF NON COMPLIANCE

The responses to the ROMI open-ended question (what is your main motivation for not taking the medication?) were grouped into thirteen closed questions in the ROMI scale, which is shown table. 5. The major reasons of noncompliance which are significantly different among compliant and noncompliant groups are 1) no perceived daily benefits due to medicines (p=0.02*), 2) negative relationship with the treating psychiatrist (p=0.03*), 3) financial obstacles (p=0.03*), 5) embarrassment or stigma related to treatment and medications (p=0.01*), 6) medications currently not necessary (p=0.03*). Whereas the reasons of health professional and friends opposed to treatment and medications, substance abuse, denial of illness, distress due to medications sideeffects, desire to be rehospitalized were comparable between the two groups.

Figure 5

Table 5: Rating of patient's medication influence (ROMI) scale closed questions (reasons for noncompliance)(n=75)

Reasons of compliance	Compliant (46)			Noncompliant (29)			Value
	Degree of influence			Degree of influence			
	None	mid	Strong	None	mid	Strong	
No perceived ay to day benefit	89.1	8.7	2.2	65.5	17.2	17.2	$\chi^2=7.3$ p=0.02*
Negative relationship with the physician	89.1	8.7	2.2	65.5	17.2	17.2	$\chi^2=6.09$ p=0.03*
Health professional opposed to medication	89.1	8.7	2.2	75.9	10.3	13.8	$\chi^2=3.7$ p=0.1
Family/ friend opposed to taking medication	78.4	8.7	10.9	72.4	17.2	10.3	$\chi^2=4.7$ p=0.09
Difficulty in gaining access to treatment	78.3	13.0	8.7	65.5	10.3	24.1	$\chi^2=5.7$ p=0.05
Embarrassment or stigma associated with treatment/medication	50.0	28.3	21.7	20.7	27.6	51.7	$\chi^2=8.7$ p=0.01*
Financial obstacles	65.2	13.6	15.2	34.5	31.0	34.5	$\chi^2=7.03$ p=0.03*
Substance abuse	71.7	19.6	8.7	51.7	34.5	13.8	$\chi^2=1.2$ p=0.5
Denial of illness	71.7	19.6	8.7	62.1	10.3	27.6	$\chi^2=5.1$ p=0.07
No current need of taking medication	60.9	17.4	21.7	44.8	13.8	41.4	$\chi^2=7.3$ p=0.03*
Inconvenience caused by -							
The side effects of medication	45.7	32.6	21.7	37.9	27.6	34.5	$\chi^2=1.4$ p=0.4
Desire to be re-hospitalized	63.0	17.4	19.6	86.2	10.3	3.4	$\chi^2=5.3$ p=0.06

Chi square test
*p<0.05
**P<0.008 after doing Bonferroni Post Hoc correction for multiple variables

DISCUSSION

This study would be perhaps the first study in India, which measured the attitudes and reasons of patients with schizophrenia about medication compliance on the same subjects. Compliance in this study is defined as the degree to which the patient consistently follows the instruction given by the doctor regarding medication and treatment. For various reasons measuring compliance status accurately is a difficult process as there are different ways to measure compliant status i.e. subjective method and objective method like serum drug levels (8).

In the published literature, demographic variables have not been consistently correlated with medication noncompliance in schizophrenia. It has been found in similar previous studies that younger age group is a significant predictor of noncompliance (22,23). Results of current study showed that there is no significant relationship of the age of the patient and their family member with the compliance. Present study results are in support of the previous findings that there is no significant relationship of gender with compliance (23)

Previous study shows lower treatment compliance in highly educated patients; it could be the result of social disapproval of psychiatric morbidity (24). But in the current study the educational status of the patient is significantly correlating with the medication non compliance, as the majorities (68.8%) of patients in the noncompliant group are educated up to senior secondary school and below where as majority (67.4%) of patients in the compliant group are educated senior secondary school and above. This significant difference can be due to the fact that the educational status is

significantly correlated to the ability to understand and comprehend the psycho education given by the treating team. The educational statuses of the family member were not significantly related to the compliance as both groups are homogeneous in terms of educational status.

As per the previous studies marital status is significantly related with the compliance and found that married patients are more likely to be compliant (24) there is no significant relationship between marital statuses with the compliance.

Employment predicted a negative attitude towards antipsychotic medication; possibly related to social stigma (6), Majority of patients in current study are unemployed/lost job due to illness in the compliant group and noncompliant group and are not correlated with the noncompliance. Family income was not correlated with noncompliance in this study. This may be due the fact most of the patients are either from poor or lower middle income background, but family members spare money for their patient's treatment and medication because of the fear of relapse and majority stay in joint family where family support is high.

In the current study majority of the patients in the compliant group (76.1%) stay in urban areas whereas 58.6% of patients in the noncompliant group stay in rural areas. Rural background was significantly related with noncompliance in this study. The main draw back for the patients staying in the rural areas are there are no effective community mental health services available, there is -unavailability of the certain medications, high cost for each follow-up to at the selected setting which is located in urban area. This may be also due to the fact that less number of high education groups in the villages as the educated herds move to cities for employment.

Current study findings revealed that adjustment difficulty of the patients with their family/spouse were significantly related to noncompliance. In these families there will be proper supervision of medication administration or absence of 'watch dog effect' of family member upon the patient might partially contribute to the noncompliant behavior of the patient.

Most of the previous studies showed that substance abuse is a strong predictor of noncompliance (13,14,18,25-30), but in the current study the patients in the compliant and noncompliant groups using substances like alcohol and tobacco, which is not significantly correlated to the

noncompliance. This percentages may not be the actual percentage because of the fear of the patients that researcher could tell their family and treating psychiatrist about their abuse which may worsen their relationship with them. In addition no objective methods were used to assess the actual substance abuse.

History of noncompliance in the past significantly predicts current noncompliance (29) Previous history of noncompliance is significantly correlated to the medication noncompliance ($p=0.01$). Ninety seven percent of the noncompliant group had pervious history of noncompliance whereas only eighty-seven in the compliant group had history of noncompliance in the past. Even though the compliant group had history of noncompliance, frequent psycho education during the follow-ups might have changed their attitude towards medication and treatment

Insight into the illness is a strong predicator of compliant behaviors; lack of insight significantly predicts noncompliance (13, 15, 18, 19,30). The current study findings support the previous study findings. Majority of the non-compliant had poor insight (grade1-3)(58,6%).

Other strong predictor of noncompliance is the severity of illness. The current study has used PANSS for assessing the severity. Previous studies showed that higher positive symptom PANSS score (13-15, 18,27) and higher total symptoms score (27,32) predicts noncompliance. In the current study only positive symptom score is found significantly correlated to noncompliance. Increased positive symptoms like hallucination, suspiciousness, and hostility leads to refusal to take medications due to unawareness of the disease and subjective need to rely upon medication (7).

In the current study no significant differences found between the compliant and noncompliant group as per their demographic and selected characteristics like gender, age, employment status, monthly family income, religion, number of daily medicine, medicine availability, duration of illness, medication supervision, cost of medicines, co morbid physical illness, family history of mental illness, frequency of follow-up, and cost of transport. Previous study findings had been contradictory to some of these findings.

The attitude scores of DAI represent an indirect representation of compliance (7). There is a high level of general positive attitude towards medication among patients with schizophrenia (6,11,33). These results are supporting the current study findings, which shows that majority of

patients and family members in both the group have positive attitude towards medication. The family member's high positive belief in treatment and medication might be the reasons for the positive attitude of the patient and increasing the pressure on patient for taking medications, which results in compliance.

Previous studies suggest that the reasons which significantly predict compliance as per the ROMI items are 'perceived daily benefit' (8), positive relationship with clinicians and for noncompliance is 'inconvenience due to side effects.'(8,9) In the current study the reasons of compliance which are significantly correlated to compliance are 'perceived daily benefit from medication', 'positive relationship with psychiatrist', pressure from the family and health system', 'positive family belief towards illness and treatment'.

Where as the significant reasons for noncompliance are 'no perceived daily benefit from medications', 'difficulty in gaining access to treatment and medications', financial obstacles', 'embarrassment or stigma related to treatment and medications', 'medicines currently not perceived as necessary'. While comparing the reasons of compliance/noncompliance given by the patient and family member, there is significant difference between the reasons of patients and that of family members, i.e. the family members are more clearly identifying the patient's reasons of compliance and noncompliance. Therefore the mental health professional should interview the patient and family members separately to get the clear picture of patient's compliant behaviors towards medication and treatment.

Previous studies have indicated that fifty to sixty percent of patients with schizophrenia in the outpatient unit are noncompliant to their medication (14,24,31,34,35). In the current study 38.7% of patients are noncompliant to their medication and treatment in the outpatient unit of the selected setting. Even though majority of patients are compliant but, thirty nine percent of the patients still fail to adhere with the prescribed treatment regime, which is an alarming finding and requires great attention. This reflects that the treating team needs to be more committed and should give proper attention to this high-risk group, as they are more likely to relapse in the near future.

Majority of the noncompliant patient are lesser educated, and are staying in rural areas. The follow-up counseling for these patients has limited scope, as it is psychiatrist driven

who is overburdened with patient load. The family member and patients are reluctant to ask their queries about their medication and treatment and most of the doubts remain unanswered during the follow up. This results in the abrupt cessation of medicines as soon as the patient reaches remission leading to noncompliance and relapse. So these patients need special attention and proper adherence counseling.

At present there is no support group functioning, which may provide services for the family members to solve the issues and burden of the family member and their role in patient care for improving compliance. The study findings strongly recommend the need for starting the of support groups to tackle the problems of the family member in managing their patient effectively. The study findings suggest that standard protocol need to be developed for providing consistent medication adherence counseling for the mental health professionals, which will improve the compliant behaviour of patients towards their medication and treatment.

In conclusion, we have shown that overall patients and their family member have a positive attitude towards medication and treatment, family member is having more positive attitude and are able to identify the reasons of noncompliance better than their patients. The prevalence of noncompliance in the selected is 37.8%, and the factors which significantly related to noncompliance are lesser education status, rural area of stay, adjustment difficulty with family members, higher positive symptoms severity, poor insight and previous noncompliance. These factors should be incorporated into any programme designed to improve overall compliance.

References

1. WEIDEN PJ, DIXON L, FRANCES A, APPELBAUM P, HAAS G, RAPKIN B. Neuroleptic noncompliance in schizophrenia. *Schizophr Bull* 1990;12:312-14.
2. LINN MW, KLETT CJ, CAFFEY EM. Relapse of psychiatric patients in foster care. *Am J Psychiatry* 1982;139:778-83.
3. CRAMER JA, ROSENJECK R. Compliance with medication regimens for mental and physical disorders. *Psychiatr Serv* 1998;49:196-201
4. ADAMS SG, HOWE JT. Predicting medication compliance in a psychotic population. *J Nerv Ment Dis* 1993;181:558-560
5. NABER D, KAROW A, LAMBERT M. subjective well-being under neuroleptics treatment and its relevance to compliance. *Acta psychiatr Scand* 2005;111:232-243
6. FREUDENREICH O, CATHER C, EVINS AE, HENDERSON DC, GOFF DC. Attitudes of schizophrenia outpatients toward psychiatric medications: relationship to clinical variables and insight. *J Clin Psychiatry* 2004;65:1372-1376
7. ADEWUYA AO, OLA BA, MOSAKU SK, FATOYE FO, EEGUNRANTI AB. Attitude towards antipsychotics among outpatients with schizophrenia in Nigeria. *Acta Psychiatr Scand* 2006;113:207-211
8. ROSA MA, MARCOLIN MA, ELKIS H. Evaluation of the factors interfering with the drug treatment compliance among Brazilian patients with schizophrenia. *Rev Bras Psiquiatr*. 2005;27:178-84
9. LOFFLER W, KILLAN R, TOUMI M, ANGERMEYER MC. Schizophrenic patients subjective reasons for compliance and non compliance with Neuroleptic treatment. *Psychiatr Serv* 2004; 55:174-179.
10. MULAIK JS. Noncompliance with medication regimens in persistently mentally ill schizophrenic patients. *Issues Ment Health Nurs* 1992; 13:219-37.
11. HOFFER A, KEMMLER G, EDER U, EDLINGER M, HUMMER M, FLEISCHHACKER WW. Quality of life in schizophrenia: the impact of psychopathology, attitude towards medication, and side effects. *J Clin Psychiatry* 2004; 65:932-9.
12. DOLDER CR, LACRO JP, DUNN LB, JETSE DV. Antipsychotic medication adherence: is there any difference between typical and atypical agents? *Am J Psychiatry* 2002;159: 103-108
13. KAMALI M, KELLY BD, CLARKE M, BROWNE S, GERVIN M et al. A prospective evaluation of adherence to medication in first episode schizophrenia. *Eur Psychiatry* 2006;21:29-33
14. POGGE DL, SINGER MB, HARVEY PD. Rates and predictors of adherence with atypical antipsychotic medication: a follow-up study of adolescent inpatients. *J Child Adolesc Psychopharmacol* 2005; 15(6):901-12.
15. RITTMANNBERGER H, PACHINGER T, PETER K, JOHANNES W. Medication adherence among psychotic patients before admission to inpatient treatment. *Psychiatr Serv* 2004;55:174-9.
16. TAJ R, KHAN S. A study of reasons of noncompliance to psychiatric treatment. *J Ayub Med Coll Abbottabad* 2005;17(2):26-8.
17. CHRISTY LM, ERIC YH, CHEN CS, YIP KC, LAW CW, CINDY PY et al. Anti-psychotics adherence among outpatients with schizophrenia in Hong Kong. *J Clin Psychiatry* 2004;59(5):646-52
18. DROULOUT T, LIRAUD F, VERDOUX H. Relationship between insight and medication adherence in subjects with psychosis. *Encephale* 2003;29(5):430-7.
19. COLOM F, VIETA E, MARTINEZ-ARAN A, REINARES M, BENABARRE A, GASTO C. Clinical factors associated with treatment noncompliance in euthymic bipolar patients. *J Clin Psychiatry* 2000; 61(8):549-55.
20. KAY SR, FIZBEIN A, OPLER LA. the positive and negative syndrome scale(PANSS) for schizophrenia. *Schizophr bull* 1987;13:261-276
21. WEIDEN P, RAPKIN P, MOTT T, ZYGMUNT A, GOLDMAN D, HORVITZ-LENNON M et al. Rating of medication influences (ROMI) scale in schizophrenia. *Schizophr Bull* 1994; 20:297-310.
22. CHRISTY LM, ERIC YH, CHEN CS, YIP KC, LAW CW, CINDY PY. Anti-psychotics adherence among outpatients with schizophrenia in Hong Kong. *J Clin Psychiatry* 2004;59:646-652.
23. DIAZ E, NEUSE E, SULLIVAN MC, PEARSALL HR, WOODS SW. Adherence to conventional and atypical antipsychotic after hospital discharge. *J Clin Psychiatry* 2004; 65(3):354-60.
24. SRINIVASAMURTHY R, GHOSH A Wig NN.

Treatment acceptance patterns in a psychiatric outpatient clinic: a study of demographic variable and clinical variables. *Indian Journal of Psychiatry* 1974;16:323-324.

25. GRUNEBaum MF, WEIDEN PJ, OLSON M.

Medication supervision and adherence of persons with psychotic disorders in residential treatment settings: a pilot study. *J Clin Psychiatry* 2001; 62:394-9

26. CUFFEL BJ, ALFORD J, FISCHER EP, OWEN RR. Awareness of illness in schizophrenia and outpatient treatment adherence. *J Nerv Ment Dis* 1996;184:653-9.

27. KAMALI M, KELLY BD, CLARKE M, BROWNE S, GERVIN M, KINSELLA A et al. A prospective evaluation of adherence to medication in first episode schizophrenia. *Eur Psychiatry* 2006;21:29-33

28. POGGE DL, SINGER MB, HARVEY PD. Rates and predictors of adherence with atypical antipsychotic medication: a follow-up study of adolescent inpatients. *J Child Adolesc Psychopharmacol* 2005; 15(6):901-12.

29. LACRO JP, DUNN LB, DODLER CR, LECKBAND SG, JESTE DV. Prevalence of and risk factors for medication nonadherence in patients with schizophrenia: a comprehensive review of recent literature. *J Clin Psychiatry* 2002; 63(10):892-909.

30. OLFSON M, DAVID M, HANSELL S, CAROL

A, WALKUP J, WEIDEN PJ. Predicting medication noncompliance among patients with schizophrenia. *Psychiatr Serv* 2000;51(2):216-22.

31. GLIMMER TP, CHRISTIAN RD, PHAM D, LACRO JP, FOLSOM DP et al. Adherence to Treatment with Antipsychotic Medication and Health Care Costs among Medicaid Beneficiaries with Schizophrenia. *Am J Psychiatry* 2004; 161:692-99.

32. HUDSON TJ, OWEN RR, THRUSH CR, HAN X, PYNE JM, SULLIVAN G et al. A pilot study of barrier to medication adherence in schizophrenia. *J Clin Psychiatry* 2004;65(2):211-6.

33. RUSHER SM, WIT R, MAZMANIAN D. Psychiatric patient's attitudes about medication and factors affecting noncompliance. *Psychiatr Serv* 1997; 48(1):82-5.

34. HORAN W, BORGARO SR, STOKES JM, POGGE DL, HARVEY PD. Predictors of medication compliance after hospital discharge in adolescent psychiatric patients. *Child Psychiatry Hum Dev* 1999;30(2):103-110

35. KIKKERT MJ, KEMMLER G, EDER U, HONDER M, HUMMER M, FLEISHHACKER WW. Attitudes towards antipsychotics among outpatient clinic attendees with schizophrenia. *J Clin Psychiatry* 2006;123:49-53.

Author Information

Renju Sussan Baby

College of Nursing, All India Institute of Medical sciences

Sandhya Gupta

Associate professor, Department of psychiatry, All India Institute of medical sciences

Rajesh Sagar

Lecturer, Psychiatry nursing, All India Institute of medical sciences