

Profertility Effects Of Alcoholic Extract Of Sesame In Male Sprague-Dawley Rats

V Ukwenya, K Oluyemi, E Ashamu, C Saalu, O Oyewo, V Makanjuola

Citation

V Ukwenya, K Oluyemi, E Ashamu, C Saalu, O Oyewo, V Makanjuola. *Profertility Effects Of Alcoholic Extract Of Sesame In Male Sprague-Dawley Rats*. The Internet Journal of Nutrition and Wellness. 2007 Volume 5 Number 2.

Abstract

The effects of administration of ethanolic extract of sesame and Vitamin C as adjuvant on the reproductive system of male sprague-Dawley rats were exploited in this work. 20 rats were randomized into four groups (Control, T₁, T₂, T₃) of 5 rats each. One of the group serves as control group while the other 3 groups received a sub-lethal dose of sesame extract (3000mg/kg body weight) once in a day for 8 weeks. One of the treatment groups (T₃) also received 2.5mg/kg of Vitamin C for 8 weeks. At the end of the 8 weeks, animals in Control, T₁ and T₃ were sacrificed while T₂ was withdrawn from extract administration for 4 weeks to test for the effect of withdrawal. Sperm counts, Morphology and Motility were quantified and organ and general body weights were measured in a weighing balance. Results of the experiment revealed an increment in the sperm count and sperm percentage motility but a decreased percentage morphology distortion. There were dose-dependent increments in body and organ weight gains, withdrawal notwithstanding. This experiment thus revealed that ethanolic extract of sesame seed possesses potent profertility properties which may be beneficial to those who consume it.

INTRODUCTION

The attention given to various use of phytomedicinal products are now on the increase. Many researchers now focus on several ways of making infertility a thing of the past as well as effectively controlling unwanted pregnancies. The traditional use of Beniseed (English: Sesame; Yoruba: Ekuku; Igbira: Igorigo.) dates back to about 4000 years ago in Babylon and Assyria where it is used to produce Sesame cake, wine, brandy¹. It is an important oilseed crop believed to have originated from tropical Africa where you have the greatest genetic diversity. It was later taken at a very early date to India where a secondary center of diversity developed². The total world crop area under beniseed is about 6million hectares. 66% of this is concentrated in Asia while 25% is planted in Africa (mainly Nigeria, Ethiopia and Sudan) and 8% in America (Venezuela, Mexico, Guatemala and Columbia). The leading world producers are India, China, Mexico and Sudan in Africa. Total annual consumption is about 65% for oil extraction and 35% for food. The sesame oil produced from its seed compares favourably with olive oil as this has been substituted for the latter in many places around the world and especially in Europe. It is also used in sesame cake, salad, wine, brandy, buns and chips³. It is expedient to know that some of the potent phytochemical components are antioxidants and

steroids. Antioxidants such as sesamin (0.34 to 1.13%), sesamol (0.13 to 0.58%) and sesamol (liberated from sesamol by diluting it with mineral acids or by hydrogenation. National Cereals Research Institute have been found to have nutraceutical effects^{3,4}. Antioxidants are known to enhance fertility either directly or indirectly. Most plants rich in antioxidants have been found to increase sperm counts, motility and enhance sperm morphology^{5,6}. These compounds are capable of removing free radicals from the body system. Vitamins C and E (?-tocopherol) are free radical scavengers and they protect against lipid peroxidation. This ability has been reported to both increase peripheral testosterone level and enhance attenuate testicular toxicity⁷. Vitamins present in sesame include Thiamin (0.98mg/100g), Riboflavin (0.25mg/100g) and Niacin (5.40mg/g). These are all potent antioxidants and are capable of increasing testosterone formation by the interstitial cells of Leydig.

The seed contains fatty and non-fatty acids. The relative amount of each depends upon variety and quality of seed. Fatty acid compositions of typical beniseed oil are 7 to 11% of palmitic acid; 2 to 6% of stearic acid; 32 to 54% of oleic and 39 to 56% of linoleic acid. Other fatty acids occur in amounts less than 1%³. Fatty acids such as cholesterol formed from smaller fats are essentially needed in the

formation of testosterone ⁸ .

Sesame oil-containing diet significantly reduced atherosclerotic lesion formation and plasma cholesterol, triglyceride and low density lipoprotein (LDL) levels in LDL Receptor negative mice ⁹ .

Sesame seeds have been implicated as having anti-tumorigenic ¹⁰ , estrogenic and/or anti-estrogenic ^{11,12} and antioxidant ¹³ properties.

The seminal vesicle secretes a yellow viscous fluid containing substances that activate sperm (fructose). This fluid constitutes about 70% of the human ejaculate ¹⁴ .

Although the seminal vesicle may not contain factors that are absolutely responsible for fertilization, their secretion still plays an important role in optimizing conditions for sperm viability, motility and survival as well as sperm transport (). Testis is the male gonad that produces the spermatozoa through cascade of processes collectively referred to as spermatogenesis. This process is under the influences of testosterone, FSH and LH ^{5,15} .

The present research aims at evaluating the effects of ethanolic extract of beniseed on sperm parameters and weights of reproductive organs as indices for fertility. The organs tested are testis, epididymis and seminal vesicle.

MATERIAL AND METHODS

ANIMALS

Twenty (20) adult male rats of Sprague Dawley strain (Approx. weight of 200g) were used in this study. They were obtained from Animal house of the faculty of Basic medical sciences, Ladoko Akintola University of Technology, Ogbomoso. They were brought to the animal control room of the Anatomy Department, University of Lagos. The rats were kept in iron cages at controlled room temperature of about 30°C and photo-periodicity of 12L: 12D. They were fed on rat pellet feed obtained from Ladokun feeds, Ibadan and water made available "ad libitum". They were allowed to acclimatize for two weeks before the commencement of the experiments.

PLANT MATERIAL

Beniseed was purchased locally at Ganmo market, Ilorin, Kwara State. The seed was authenticated at Botany Department, University Lagos, Lagos. Alcoholic extraction and Lethal Dose (LD) estimation of beniseed was done at Pharmacognosy Department, College of Medicine

University of Lagos. LD₅₀ of beniseed was found to be 5000mg/kg.

EXPERIMENTAL DESIGN

The rats were divided into four (4) groups of 5 rats each. The groups were designated as control (c) and Treatments (T) with subscript 1, 2 and 3. i.e.

C = Control group, T₁ = Treatment group 1, T₂ = Treatment group 2 and T₃ = Treatment group 3.

Each of the rats in the treatment groups (T₁, T₂, T₃) received 3000mg/kg body weight of beniseed extract daily between 07.00 to 8:00 hours for a period of 8 weeks. T₃ also received 2.5mg of Vitamin C (made by Tuyil Pharmaceutical Company Ilorin, Kwara State) in addition to beniseed extract. The animals were weighed once a week. After the initial period of 8 weeks, rats in T₃ were withdrawn from beniseed extract for a period of 4 weeks. The animals were sacrificed by cervical decapitation as follows: C, T₁ and T₃ at the end of 8th week and T₂ at the end of 12th week.

SEMEN COLLECTION AND SEMEN ANALYSES

The caudal epididymis of one rat in each of the experimental group were removed and minced into several pieces on a specimen bottle containing normal saline for some few minutes to allow the sperms to become motile and swim out from the caudal epididymis ^{16,17} . The semen was then taken with 1ml pipette and dropped on a clean slide and covered with cover slips. The slides were placed under light microscope and examined for morphology and sperm motility according to Saalu et al., ^{16,17} .

DETERMINATION OF SPERM COUNTS

The new improved hemocytometer counting chamber of Neubauer's type was used for the sperm count in accordance with Oluyemi et al., ⁵ . This procedure was carried out twice on each sample to eliminate error of confidence.

MEASUREMENT OF TISSUE WEIGHTS

The weights of the testes, epididymides and seminal vesicles were taken before immersing in 10% formol saline using an electrical weighing balance.

STATISTICAL ANALYSIS

Where applicable, the data obtained were analyzed statistically by students' T-test and one way analysis of variance (ANOVA). The level of significance was at <0.05.

RESULT

Figure 1

Figure 1.0 Effects of extract administration on Mean body weight of animals

Weight	Control (g)	T1 (g)	T2 (g)	T3 (g)
Initial	217±16.3	232.8±13.1	203.3±26.6	196.6±24.0
After 4 weeks	233.2±15.5	235.7±13.1	206.6±25.9	213.7±22.6
Weight gained	>16±0.8	>2.9±0.0	>3.3±0.7	>17.1±1.4
After 8 weeks	265.7±14.7	251.2±12.4	239.6±26.6	221.9±22.3
Weight gained	>32.5±0.8	>15.5±0.7	>33.0±0.7	>8.2±0.3
After 12 week	-	-	251.3±21.2	-
Weight gained	-	-	>11.7±5.4	-

n = 5, * Significantly different from control (P<0.05); >positive weight gain. Values are recorded as Mean±S.E.M

Figure 2

Figure 1.1 Effects of extract administration on mean weight of testes of the animals after the experiment

	CONROL*	T ₁ (g)*	T ₂ (g)*	T ₃ (g)*
Testis	2.19±0.15	2.30±0.30#	2.29±0.09#	2.62±0.11*
Epididymis	0.41±0.02	0.54±0.01*	0.42±0.01#	0.82±0.02*
Seminal vesicle	0.0108±0.002	0.0164±0.003*	0.0132±0.003*	0.0170±0.003*

n = 5, * Significantly higher than control (P<0.05); # Not significantly different from control (P<0.05); Values measured as Mean ± S.E.M

▼ after 8 weeks; ▲ after 12 weeks

Figure 3

Figure 1.2 Effects of extract administration on the sperm count, motility and morphology

	CONTROL*	T ₁ *	T ₂ *	T ₃ *
Sperm Count (10 ⁶ /ml)	23.2 ± 8.9	37.6 ± 11.4	34.4 ± 11.5	41.6 ± 6.5
Sperm Motility (%)	64 ± 6.8	76 ± 13.3*	72 ± 11.2*	79 ± 3.7*
Abnormality (%)	14.6 ± 1.8	5.6 ± 7.2	5.2 ± 3.7	3.4 ± 1.7

n = 5, * Significantly higher than control (P<0.05); # Significantly lower than control (P<0.05); Values measured as Mean ± S.E.M

▼ after 8 weeks; ▲ after 12 weeks

DISCUSSION

The effects of ethanolic extract of beniseed (sesame) at 3000mg/kg body weight, with vitamin C administered as adjuvant, have shown that beniseed has a potential to increase mean body weights of rats as shown in Figure 1.0. This is most due to the high fat contents of the seed. Fats are stored in the form of Triacylglycerol (TAG) in adipose tissues in mammals via lipogenetic pathways¹⁸. The increased weight gain is in line with the work of¹⁹ Shittu et al., who also recorded a dose-dependent increase in weight gain upon administration of 14.0mg/kg and 28mg/kg body weight of aqueous extract of sesame to rats for six weeks. Beniseed contains 43.0 - 56.8 % mass of fat and about 21.6 - 25.3 %mass of carbohydrate²⁰. Carbohydrates are also converted to TAG through lipogenesis in the liver. Insulin increases the activity of acetylCoA carboxylase and provides glycerol for esterification of fatty acids to TAG. One of the most important carbohydrates concerned with fertility is cholesterol from which male steroids hormone (testosterone in humans or androstenedione in animals) are synthesized in

the leydig cells under the influence of LH (also called Interstitial cell stimulating hormone)¹⁸. Thus, increased testosterone level is responsible for the increased sperm counts noted in the treated groups when compared with the control (Table 1.2). This is because FSH binds to sertoli cells and promotes the synthesis of androgen binding protein (ABP). Thus high concentration of androgen is made available locally at the seminiferous tubule, at the site of spermatogenesis.

Seminar vesicle secretes fructose (1.5-6.5mg/mL), Phosphorylcholine, Ergothioneine, Ascorbic acid, Flavins, Prostaglandins (fatty acid derivative)²¹. These chemical components of seminal fluid, coupled with the antioxidant property of Vitamin C, are responsible for enhancing motility of sperm; hence its increased secretion by the organ will lead to increased motility. In Table 1.2, there was a significant increase in the treatment group that received Vitamin C (i.e T₃). In the absence of Vitamin C, decreased motilities were recorded in treatment groups even upon withdrawal from extract when compared with the group that received Vitamin C alongside extract. In the overall, there were increases in the motility in the treatment group in a dose-dependent manner. It is also shown that ethanolic extract of sesame enhances morphology of the spermatozoa in the epididymis at the time of sacrifice of the animals. This is in agreement of the work of Saalu et al.,⁷.

Increased cellular activities must have led to enlargement of the component cells and invariably hypertrophy of the organs (Table 1.1). There were dose-dependent increments in the weights of selected organs; but these differences are not totally significant for all treated groups. This is a positive indication that Sesame enhances fertility by production of substances necessary for the proper function of spermatozoa. This observation is also in agreement with the work of Shittu et al¹⁹. Increased cellular activities are key factor to be considered in the evaluation of organ weights. Vasudevan & Sreekumari,¹⁸ stated that androgens stimulate spermatogenesis, produce hypertrophy of prostate, seminal vesicles, muscle, bone and kidney cells. It is anabolic. This finding perfectly explains the increase weights of these organs and body weight as a whole as seen in the present experiment.

References

1. Oplinger E.S., Putnam A.R. and hyandosn E.A. (1990). Sesame. Univ. Winsconsin Ext. 140-147.
2. Purseglove, J.W. (1996): Tropical Crops Dicotyledon 430 - 435 (1) Longman, London.
3. Beckstorm - Sternberg, V.A., Kohn F.I., Stone P.U. and

- Dennilson S.P. (1994): Beniseed in Medicine 184-201 (1) Longman, London.
4. National Cereals Research Institute (NCRI) (2002). Training manual on beniseed production technology 47, Abuja, Nigeria.
5. Oluyemi Kayode A., Jimoh Olusegun R., Adesanya Olamide A, Omotuyi Idowu O, Josiah Sunday J, and Oyesola Tolulope O, (2007) Effects Of Crude Ethanolic Extract Of Garcinia Cambogia On The Reproductive System Of Male Wistar Rats (*Rattus Novergicus*), African Journal of Biotechnology, 6(10):1236-1238.
6. Adesanya Olamide, A. Oluyemi Kayode, A. Ofusori David, O. Omotuyi Idowu, U. Okwuonu Christina, O. Ukwenya Victor, A. Adesanya Rotimi: Micromorphometric and Stereological Effects of Ethanolic Extracts of Garcinia cambogia seeds on the Testes and Epididymides of Adult Wistar Rats. The Internet Journal of Alternative Medicine. 2007. Volume 5 Number 1.
7. Saalu, L. C, Oluyemi, K. A. and Omotuyi, I. O. (2007a) a-Tocopherol (vitamin E) attenuates the testicular toxicity associated with experimental cryptorchidism in rats. African Journal of Biotechnology Vol. 6 (12), pp. 1373-1377, 18 June.
8. Maximillian Ehrenstein and Edward L. Corey. Activation of Testosterone by Higher Fatty Acids and their Acid Sodium Salts. J. Biol. Chem. 1938 122: 297-302.
9. Bhaskaran S. (2006). Journal of medicinal food. Inhibition of atherosclerosis in Low-Density Lipoprotein Receptor Negative mice by sesame oil. Vol 9. No 4.
10. Hirose N, Doi F, Ueki T, Akazawa K, Chijiwa K, Sugano M, Akimoto K, Shimizu S, Yamada H (1992) Suppressive effect of sesamin against 7,12-dimethylbenz[a]-anthracene induced rat mammary carcinogenesis. Anticancer Res 12:1259-1265
11. Collins BM, McLachlan JA, Arnold S (1997). The estrogenic and antiestrogenic activities of phytochemicals with the human receptor expressed in yeast. Steroids 62: 365-372.
12. Shittu Lukeman AJ (2006). The effect of the aqueous crude leaves extract of *Sesamum radiatum* compared to Mesterolone (proviron) on the adult male Sprague Dawley rats testis and epididymis. MSc Dissertation. Lagos State University, College of Medicine, Ikeja, Nigeria.
13. Kang M.H, Kawai Y, Naito M, Osawa, T. J. (1999). Nutr. 129:1885. In: Jeng KCG, Hou RCW. Sesamin and Sesamol: Nature's therapeutic lignans. Current Enzymes Inhibition, 1: 11-20, 2005.
14. Luiz Carlos, jungueira, Jose Carneiro [2005]. Basic Histology, text and atlas. 10th Edition. McGraw-Hill Medical; 443-444
15. Akpantah, A.O, Oremosu, A.A, Ajala, M.O, Noronha, C.C, Okanlawon, A.O 2003. the effect of crude extract of Garcinia Kola seed on the histology and hormonal milieu of male Sprague-Dawley rats' reproductive organs. Nig. J. Health and Biomed. Sci. 2(1): 40-46.
16. Saalu LC, Togun VA, Oyewopo AO, Raji Y (2006). Artificial cryptorchidism and the moderating effect of melatonin in Sprague - Dawley rats. J. Appl. Sci. 6 (14): 2889 -2894.
17. Saalu LC, Adesanya AO, Oyewopo AO, Raji Y (2007b). An evaluation of the deleterious effect of unilateral cryptorchidism on the contralateral normally descended testis. Sci. Res. Essays 2(3): 074-078.
18. Vasudevan DM and Sreekumari S (2005) Textbook of Biochemistry for Medical Students. 4th Edition, Jaypee Brothers Medical Publishers (P) LTD, New Delhi, India, pp371-377.
19. Shittu, L. A. J., Bankole, M. A., Oguntola, J. A., Ajala, O., Shittu R. K., Ogundipe, O. A., Bankole M. N., Ahmed, T. and Ashiru, O. A. (2007) Sesame leaves intake improve and increase epididymal spermatocytes reserve in adult male Sprague Dawley rat. Scientific Research and Essay Vol. 2 (8), pp. 319-324, August
20. Dym, M. (1983). The male reproductive system in: Histology-cell and tissue Biology. 5th Edition. Edited by Leon weiss. NY: Elserier science publishing con.inc. 1001-1053.
21. Ganong (1999) Review of medical Physiology. 20th edition,

Author Information

Victor O. Ukwanya

Department of Anatomy, School of Basic Medical Sciences, Lagos University Teaching Hospital

Kayode Alaba Oluyemi

Department of Anatomy, School of Basic Medical Sciences, Igbinedion University

Ebenezer Ashamu

Department of Anatomy, School of Basic Medical Sciences, Ladoke Akintola University

Chia Saalu

Department of Anatomy, School of Basic Medical Sciences, Lagos State University College of Medicine

Oyebowale O. Oyewo

Department of Anatomy, School of Basic Medical Sciences, Lagos University Teaching Hospital

Victor O. Makanjuola

Department of Anatomy, School of Basic Medical Sciences, Lagos University Teaching Hospital