

Ogilvie's Syndrome: The Rectal Balloon Sign

V Naraynsingh, M Ramdass, S Teelucksingh, A Perry, D Maharaj

Citation

V Naraynsingh, M Ramdass, S Teelucksingh, A Perry, D Maharaj. *Ogilvie's Syndrome: The Rectal Balloon Sign*. The Internet Journal of Surgery. 2002 Volume 4 Number 1.

Abstract

Intestinal pseudo-obstruction as described by Ogilvie¹ in 1948 is a well recognized clinical entity of uncertain etiology and pathophysiology. Despite its well known clinical features, the diagnosis of intestinal pseudo-obstruction remains quite difficult and is often confused with mechanical obstruction. We describe three cases with an easily recognizable clinical finding, which can serve to distinguish Ogilvie's syndrome from mechanical large bowel obstruction.

CASE 1

An 89-year old female sustained a fractured femur after falling down a flight of stairs. Two days after insertion of an Austin-Moore prosthesis she began to develop gross painless distension of her abdomen. Bowel sounds remained normal and digital rectal examination (DRE) revealed a dilated, air-filled rectum without stool. Plain abdominal films showed gross distension of her entire colon and rectum. A tentative diagnosis of Ogilvie's syndrome was made. One day later she passed copious amounts of stool and her distension subsequently resolved completely.

CASE 2

A 62-year old male diabetic with chronic renal failure presented with constipation for one week. Examination revealed a grossly distended abdomen with normal bowel sounds. Plain supine abdominal X-rays revealed grossly distended colon up to the recto-sigmoid area. DRE revealed an air-filled rectum devoid of stool. Two days after presentation, the abdominal distension resolved with the passage of large amounts of watery stool and flatus.

CASE 3

A 57-year old alcoholic cirrhotic male was admitted with a two-day history of abdominal distension and constipation. There was a grossly distended, tympanic, mildly tender abdomen. DRE revealed a dilated, air-filled rectum without stool. He was managed conservatively and settled within 72 hours.

DISCUSSION

The accurate and early diagnosis of Ogilvie's syndrome remains elusive. Patients have concomitant disease in 80%

of cases², including femoral fractures and hip surgery³, myocardial infarction⁴ and renal failure⁵.

Symptoms include marked distension, colicky abdominal pain and constipation⁶. Nausea and vomiting are common complaints³. Examination reveals massive abdominal distension⁷ with normal, reduced or obstructive bowel sounds. Tenderness is usually minimal².

The features of Ogilvie's thus closely mimics mechanical large bowel obstruction^{8,9}. Since the treatment of these two clinical entities is vastly different, a firm diagnosis of Ogilvie's is paramount to avoid unnecessary surgical intervention.

The authors have noted that in all three cases of Ogilvie's syndrome, DRE revealed not only an empty rectum, but a rectum that was filled with air, such that the examining finger on passing via a snug fitting anal canal enters a massively ballooned, air-filled space, resulting in much difficulty in reaching its walls.

CONCLUSIONS

We believe that the related paralytic disorder that affects the colon should also affect the rectum, leading to distension of this portion of the gut as well. This finding is quite distinct from the empty but collapsed rectum palpated in a patient with mechanical colonic obstruction where the distal colon and rectum retains its muscular tone.

This feature may be the only frequent, easily detectable physical finding that differentiates mechanical obstruction from pseudo-obstruction, thus avoiding unnecessary surgical intervention or uncertainty in diagnosis.

CORRESPONDENCE TO

Mr. Michael J. Ramdass 100 East Drive, Champs Fleurs, St. Joseph, Trinidad, West Indies. E-mail: jimmyramdass@hotmail.com

References

1. Ogilvie H. Large-intestine colic due to sympathetic deprivation. *Br Med J* 1948; ii: 671-3
2. Dorudi S, Berry AR, Kettlewell MGW. Acute colonic pseudo-obstruction. *Br J Surg* 1992; 79:99-103
3. Adams JT. Adynamic ileus of the colon. An indication for caecostomy. *Arch Surg* 1974; 109: 503-7
4. Caves PK, Crockard HA. Pseudo-obstruction of the large bowel. *Br Med J* 1970; 11: 583-6
5. Sloger AF, Panella VS, Demas BE. Ogilvie's syndrome. Successful management without colonoscopy. *Dis Dog Sci* 1988; 33: 1301-6
6. Morton JH, Schwartz SI, Gramiak R. Ileus of the colon. *Arch Surg* 1960; 81: 425-34
7. Bullock PR, Thomas WEG. Acute pseudo-obstruction of the colon. *Ann R Coll Surg Eng* 1984; 66: 327-30
8. Dudley HAF, Paterson-Brown S. Pseudo-obstruction. *Br Med J* 1986; 292: 1157-8
9. Jones PF. Acute colonic pseudo-obstruction. In: *Emergency abdominal surgery in infancy, childhood and adult life*. Oxford Blackwell Scientific 1987: 235-9.

Author Information

Vijay Naraynsingh, FRCS FACS

Department of Surgery, University of the West Indies, General Hospital

Michael J. Ramdass, MBBS MRCS

Department of Surgery, University of the West Indies, General Hospital

S. Teelucksingh, MRCP

Department of Surgery, University of the West Indies, General Hospital

Andrew Perry, MBBS

Department of Surgery, University of the West Indies, General Hospital

Dale Maharaj, FRCS

Department of Surgery, University of the West Indies, General Hospital