

Molecular Characterisation Of Leptospiral Isolates Identified In Agricultural Labours Of Salem District Of Tamilnadu

N Jeyakumar, A Ranjitsingh, K Seenivasan, I Joseph

Citation

N Jeyakumar, A Ranjitsingh, K Seenivasan, I Joseph. *Molecular Characterisation Of Leptospiral Isolates Identified In Agricultural Labours Of Salem District Of Tamilnadu*. The Internet Journal of Microbiology. 2008 Volume 6 Number 2.

Abstract

In the present study, RAPD analysis of the two Leptospire isolates identified (K₁ and K₂) were subjected to random amplified polymorphic DNA fingerprinting analysis and compared with 5 genomospecies. Genotype characters of isolated K₁ and K₂ have poor resemblance with reference strain Panama CZ 14M, LSU 1945 and Sarmin CZ 390. The genomic profile of K₁ and K₂ were very similar and resemble the genomic profile of Rachmat belongs to the genomospecies *Leptospira interrogans*. The present study reports the prevalence of the leptospire belongs to Rachmat strain of the genomospecies *Leptospira interrogans*.

INTRODUCTION

Traditionally leptospire are classified and identified in serological techniques such as the cross-absorption tests (Gerritsen et al., 1995). However, these techniques are laborious and show poor inter-laboratory reproducibility (Kobayashi et al., 1985), but still cannot distinguish all serovars. The DNA based methods for the identification of *Leptospira* holds good results and these method include restriction fragment length polymorphism (RFLP) analysis (Terpstra et al., 1978; Hookey et al., 1987; Ellis et al., 1991); DNA-DNA hybridisation (Perolat et al., 1993); Pulsed field gel electrophoresis (PFGE) (Herrmann et al., 1992); Polymerase chain reaction followed by RFLP analysis (Ralph et al., 1993) and random amplified polymorphic DNA (RAPD) fingerprinting (Corney et al., 1993; Natarajaseenivasan et al., 2004). RFLP analysis is generally more sensitive and discriminatory than serotyping, but some serovars are not readily distinguished. (Thiermann et al., 1986), and the complex banding patterns are difficult to interpret. Hybridization of labelled DNA probes to RFLP blots greatly simplifies the visualised banding patterns facilitating interpretation. (Van Eys et al., 1991) and allows serovars with similar RFLP banding pattern to be differentiated (Zuerner et al., 1993). RFLP analysis, DNA-DNA hybridization and PFGE are slow and require large amounts of purified DNA. In contrast, PCR - based methods

are rapid and require only small amounts of DNA. (Van Eys et al., 1989; Gerritsen et al., 1991; Merien et al., 1992; Gravekamp et al., 1993). PCR-based fingerprinting systems have been well developed for a range of bacteria. (Corney et al., 1993; Giesendorf et al., 1993; Ralph et al., 1993; Van Belkum et al., 1993; Woodward et al., 1993). Because of some disadvantage in the application of cross-absorption agglutination (CAA) technique and in restriction endonuclease analysis (REA) alternative methods were searched. Corney et al. (1993) stressed the need to develop a rapid and simple typing method which can distinguish different genotype without the disadvantages of CAA and REA. Rapid identification of the isolates would allow farmers to start appropriate vaccination regimens with minimal delays. Welsh and McClelland (1990) and Williams et al. (1990) developed a DNA fingerprinting technique based on the random amplification of genomic sequences by using a single primer at low stringency in a polymerase chain reaction (PCR). Gerritsen et al. (1995) investigated the use of leptospiral and non leptospiral primers in RAPD fingerprinting of leptospirosis. Perolat et al. (1998), Brenner et al. (1999) and Levett (2001) reported more than 16 genomospecies of leptospire based on genotypic characters. The molecular analysis of new isolates from different regions may increase genomospecies number of leptospire. With this background the leptospire isolates collected in the present study were subjected to random amplification of

polymorphic DNA finger printing analysis to find out whether they are new genomospecies or related to the already existing genomospecies.

MATERIAL AND METHODS

ISOLATION OF GENOMIC DNA FROM LEPTOSPIRAL ISOLATES

Blood samples from Leptospiral infected patients were inoculated into sterile EMJH medium and the isolated leptospiral strains (K₁ and K₂) were grown at 30°C in EMJH medium for 7 days to a density of 10⁸ cells/ml and centrifuged at 10,000 rpm for 30 mins at 4°C. Pellets were collected and resuspended in 400 µl of TE buffer, vortexed and centrifuged again to remove the medium debris. After removing the supernatant, 40 µl of lysozyme (10mg/ml) was added to the pellet, vortexed well and incubated at 37°C for 24 hours. After that, 56 µl of 10% sodium dodecyl sulphate and 5 µl of proteinase K (10 mg/ml) were added and incubated in a water bath at 65°C for 30 mins. Then, 80 µl of 5 M NaCl and 64 µl NaCl/CTAB was added, vortexed and incubated at 65°C in water bath 30mins. After that, 645 µl of chloroform, Isoamylalcohol (24: 1 v/v) was added, mixed well and centrifuged at 10,000 rpm for 20mins. 300 µl of clear supernatant was taken without disturbing inter phase layer and transferred into another tube. To this tube 180 µl of isopropanol was added and kept the tube in a deep freezer for 1 hour to precipitate the DNA and centrifuged at 10,000 rpm for 20 mins at 4°C to pellet the DNA. The nucleic acid was precipitated by adding 150 µl of ice cold ethanol and centrifuged at 10,000 rpm for 5 mins. The DNA thus obtained was suspended in 20 µl of TE buffer and stored at -20°C until use.

The isolated Leptospiral DNA was analysed by agarose gel electrophoresis in 1.0% agarose with 5 µl of Ethidium bromide (10 mg/ml) and the electrophoresis was carried out in a submarine gel electrophoresis apparatus at 100 V for 4 to 5 hours in TBE buffer (pH 8.0). At the end of the electrophoresis, the EtBr stained band in gel was visualized in UV transilluminator.

ELUTION OF ISOLATED LEPTOSPIRAL DNA FROM THE GEL

The agarose gel containing the DNA was cutout with a sterile scalpel. These agarose pieces were then transferred into dialysis membrane and TBE buffer was added so that the agarose pieces were surrounded by buffer. The dialysis bag was placed at the bottom of a gel apparatus filled with buffer

in such a way that the pieces of agarose was in the same position with respect to the electrodes as it was in the gel. A current of 100-200 mV was applied for 2 hours to elute the DNA out of gel. The buffer containing the electroeluted DNA was transferred through cotton plugged 1 ml tip into an eppendorf tube and the collected DNA was centrifuged at 10,000 rpm for 2mins. The supernatant was then extracted with phenol and then with chloroform: Isoamylalcohol (24:1 v/v). The extracted DNA solution was mixed with 3 M Tris-HCl (pH 7.5) to give a final concentration of 0.5 M and two volume of isopropanol was added and allowed to stand at 20°C for 30mins and centrifuged at 15,000 rpm for 10mins for repeated time to remove EtBr. The DNA pellet was then dried under vacuum and resuspended in a TE buffer and stored at 40°C until use.

QUANTIFICATION OF DNA USING UV SPECTROPHOTOMETER

The eluted DNA (50 µl) was dissolved in 3.0 ml of 1X SSC buffer. The absorbance of DNA sample was determined at 220nm using 0.1X SSC buffer as blank in a UV spectrophotometer. The wavelength was increased by 10 nm up to 300 nm, the absorbance of the sample was measured. The absorbance ratio of 260 to 280 nm were measured.

RANDOM AMPLIFIED POLYMORPHIC DNA (RAPD) FINGERPRINTING OF ISOLATED STRAIN

DNA prepared from the isolated strain (Boom et al., 1990), the RAPD fingerprinting technique was carried out using B₁₁ and B₁₂ primer as per the method described by Gerritsen et al. (1995). The amplification of isolated leptospiral DNA was carried out in a total volume of 50 µl which contains 50 ng of isolated chlorosomal DNA, 10 mM Tris HCl (pH 9.0), 50 mM KCl, 4 mM MgCl₂, 0.1mM of each of the four dNTP mixture (dATP, dTTP, dCTP and dGTP), 300pm of each of primer (B₁₁ and B₁₂) and 0.5 U of Taq DNA polymerase.

Primer B11 - CCGGAAGAAGGGGCGCCAT

B12 - CGATTAAGAAGGACTTGAACAC

AMPLIFICATION OF DNA

The amplification of DNA was carried out in a total of 35 cycles in a MJ Thermal cycler. Each cycle consist of denaturation of DNA at 95°C for 1 min, annealing of the primers at 60°C for 1 min and extension for 3mins at 72°C with a final extension for 10mins in last cycles.

AGAROSE GEL ELECTROPHORESIS OF AMPLIFIED DNA

The amplified DNA products were analysed by agarose gel electrophoresis (1.2% Agarose with 5 l EtBr (10mg/ml)) and the electrophoresis was carried out in a submarine electrophoresis apparatus at 100 V for 4-5 hours in TBE buffer (pH 8.0). At the end of the electrophoresis, the gel was visualized under UV transilluminator and photograph was taken using polaroid camara with wratten gelatin filter. The RAPD profiles were documented by using a gel documentation system (Bio-Rad, Hercules, Calif., U.S.A) and analysed by using Quantity I-D analysis software, and the dendogram were formed with 4.0% tolerance in UPGAMA. To confirm the specificity of the primers, DNA isolated from other bacteria such as E. coil and S. aureus were subjected to the same procedure with B₁₁ and B₁₂ primers.

RESULTS AND DISCUSSION

To establish a rapid identification system based on RAPD fingerprinting, RAPD profiles of 5 genomospecies were prepared. The molecular weight of the isolated leptospiral strains (K₁ and K₂) were 23,000 bp and it has the A₂₆₀₋₂₈₀ ratio of 1.70. DNA were purified and subjected to RAPD analysis. Profiles generated from field isolates were compared with this profile to enable identification of the isolates.

Purified DNA from 12 Leptospira reference strains (Ballico, Rachmat, RGA, Wumalasena, Cynopteri 3522C, Moskva V, CZ 188, Sarmin CZ 390, Perepelitsin, Mini Sari, Panama CZ 214K, and LSU 1945) comprising five different genomospecies (*L. interrogans*, *L. kirschneri*, *L. santarosai*, *L. borgpetersenii*, and *L. noguchii*) Profiles containing up to eight intense bands and a number of fainter bands were generated when the reaction products were run on agarose gel. Although genomic profile of the isolates (K₁ and K₂) had a number of common bands, they were very close to strain Rachmat (80-90%) (Fig 1 and Table 1].



Figure 1

Table 1. Table showing the percentage of similarities as per Dendogram

Serovar	Genomospecies	% of similarity
Rachmat	<i>L. interrogans</i>	80-90
Ballico	<i>L. interrogans</i>	70-75
RGA	<i>L. interrogans</i>	65-70
Perepelitsin	<i>L. borgpetersenii</i>	60-65
Mini Sari	<i>L. borgpetersenii</i>	60-50
Wumalasena	<i>L. kirschneri</i>	40-59
Cynopteri 3522 C	<i>L. borgpetersenii</i>	40-50
Moskva V	<i>L. kirschneri</i>	40-50
CZ 188	<i>L. santarosai</i>	40-50
Sarmin CZ 390	<i>L. santarosai</i>	40-50
Panama CZ 214K	<i>L. noguchii</i>	30-40
LSU 1945	<i>L. noguchii</i>	30-40

Figure 2

Fig. 1



ACKNOWLEDGEMENT

The authors are grateful to Management and Staff Members of Department of Advanced Zoology and Animal Biotechnology, Sri. Paramakalyani College, Alwarkurichi, Tirunelveli, Tamilnadu, India for their help and co-operation.

References

1. Brenner, D.J., A.F. Kaufmann, K.R. Sulzer, A.G. Steigerwalt, F.C. Rogers, and R.S. Weyant. 1999. Further determination of DNA relatedness between serogroups and

- serovars in the family Leptospiraceae with a proposal for *Leptospira alexanderi* sp. nov. and four new *Leptospira* genomospecies. *Int. J. Syst. Bacteriol.* 49: 839-858.
2. Corney, B.G., J. Colley, S.P. Djordjevic, R. Whittington, and G.C. Graham. 1993. Rapid identification of some *Leptospira* isolates from cattle by random amplified polymorphic DNA fingerprinting. *J. Clin. Microbiol.* 31: 2927-2932.
 3. Ellis, W.A., J.M. Montgomery, and A.B. Thiermann. 1991. Restriction endonuclease analysis as a taxonomic tool in the study of pig isolates belonging to the Australis serogroup of *Leptospira interrogans*. *J. Clin. Microbiol.* 29: 957-961.
 4. Gerritsen, M.A., T. Olyhoek, M.A. Smits, and B.A. Bokhout. 1991. Sample preparation method for polymerase chain reaction based semiquantitative detection of *Leptospira interrogans* serovar Hardjo subtype hardjobovis in bovine urine. *J. Clin. Microbiol.* 29: 2805-2808.
 5. Gerritsen, M.A., M.A. Smits, and T. Olyhoek. 1995. Random amplified polymorphic DNA fingerprinting for rapid identification of *Leptospira* of serogroup Sejroe. *J. Med. Microbiol.* 42: 336-339.
 6. Giesendorf, B.A.J., A. Van Belkum, and A. Koeken. 1993. Development of species-specific DNA probes for *Campylobacter jejuni*, *Campylobacter coli*, and *Campylobacter lari* by polymerase chain reaction fingerprinting. *J. Clin. Microbiol.* 31: 1541-1546.
 7. Gravekamp, C., H. van de Kemp, M. Franzen, D. Carrington, G.J. Schoone, G.J.J.M. van Eys, C.O.R. Everard, R.A. Hartskeerl, and W.J. Terpstra. 1993. Detection of seven species of pathogenic leptospires by PCR using two sets of primers. *J. Gen. Microbiol.* 139: 1691-1700.
 8. Herrmann, J.L., E. Bellenger, P. Perolat, G. Baranton, and I. Saint Girons. 1992. Pulsed-field gel electrophoresis of Not I digests of leptospiral DNA: a new rapid method of serovar identification. *J. Clin. Microbiol.* 30: 1696-1702.
 9. Hookey, J.V., S.A. Waitkins, and R. Gahir. 1987. The use of bacterial restriction endonuclease analysis to classify icterohaemorrhagiae and autumnalis serovars. *Isr. J. Vet. Med.* 43: 265-270.
 10. Kobayashi, Y., T. Tamai, and E. Sada. 1985. Serological analysis of serogroup Icterohaemorrhagiae using monoclonal antibodies. *Microbiol. Immunol.* 29: 1229-1235.
 11. Levett, P.N. 2001. Leptospirosis. *Clin. Microbiol. Rev.* 14(2): 296-326.
 12. Merien, F., P. Amouriaux, P. Perolat, G. Baranton, and I. Saint Girons. 1992. Polymerase chain reaction for detection of *Leptospira* Spp. in clinical samples. *J. Clin. Microbiol.* 30: 2219-2224.
 13. Natarajaseenivasan, K., N. Prabhu, K. Selvanayaki, S.S. Sudalaimuthu Raja, and S. Ratnam. 2004. Human leptospirosis in Erode, South India: serology, isolation, and characterization of the isolates by randomly amplified polymorphic DNA (RAPD) fingerprinting. *Jpn. J. Infect. Dis.* 57:193-197.
 14. Perolat, P., I. Lecuyer, D. Postic, and G. Baranton. 1993. Diversity of ribosomal DNA fingerprints of *Leptospira* serovars provides a database for subtyping and species assignment. *Res. Microbiol.* 144: 5-15.
 15. Perolat, P., R.J. Chappel, B. Adler, G. Baranton, D.M. Bulach, M.L. Billingham, M. Letocart, F. Merien, and M.S. Serrano. 1998. *Leptospira fainei* sp. nov., isolated from pigs in Australia. *Int. J. Syst. Bacteriol.* 48: 851-858.
 16. Ralph, D., M. McClelland, J. Welsh, G. Baranton, and P. Perolat. 1993. *Leptospira* species categorized by arbitrarily primed polymerase chain reaction (PCR) and by mapped restriction polymorphisms in PCR-amplified rRNA genes. *J. Bacteriol.* 175: 973-981.
 17. Terpstra, W.J., G.J. Schoone, and J. ter Schegget. 1986. Detection of leptospiral DNA by nucleic acid hybridization with ³²P and biotin labelled probes. *J. Med. Microbiol.* 22: 23-28.
 18. Thiermann, A.B., A.L. Handsaker, J.W. Foley, F.W. White, and B.F. Kingscote. 1986. Reclassification of North American leptospiral isolates belonging to serogroups Mini and Sejroe by restriction endonuclease analysis. *Am. J. Vet. Res.* 47: 61-66.
 19. Van Belkum, A., M. Struelens, and W. Quint. 1993. Typing of *Legionella pneumophila* strains by polymerase chain reaction mediated DNA fingerprinting. *J. Clin. Microbiol.* 31: 2198-2200.
 20. Van Eys, G.J.J.M., C. Gravekamp, M.J. Gerritsen, W. Quint, M.T.E. Cornelissen J. ter. Schegget, and W.J. Terpstra. 1989. Detection of leptospires in urine by polymerase chain reaction. *J. Clin. Microbiol.* 27: 2258-2262.
 21. Van Eys, G.J.J.M., M.J. Gerritsen, H. Korver, G.J. Schoone, C.C.M. Kroon, and W.J. Terpstra. 1991. Characterization of serovars of the genus *Leptospira* by DNA hybridization with hardjobovis and icterohaemorrhagiae recombinant probes with special attention to serogroup Sejroe. *J. Clin. Microbiol.* 29: 1042-1048.
 22. Welsh, J., and M. McClelland. 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids. Res.* 18: 7213-7218.
 23. Williams, J.G.K., A.R. Kubelik, K.J. Livak, J.A. Rafalski, and S.V. Tingey. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids. Res.* 18: 6531-6535.
 24. Woodward, M.J., and J.S. Redstone. 1993. Differentiation of *Leptospira* serovars by the polymerase chain reaction and restriction fragment length polymorphism. *Vet. Rec.* 132: 325-326.
 25. Zuerner, R.L., W.A. Ellis, C.A. Bolin, and J.M. Montgomery. 1993. Restriction fragment length polymorphisms distinguish *Leptospira borgpetersenii* serovar hardjo type hardjo-bovis isolates from different geographical locations. *J. Clin. Microbiol.* 31: 578-583.

Author Information

N. Jeyakumar

Department of Microbiology, Kamaraj College

A.J.A. Ranjitsingh

Department of Advanced Zoology and Animal Biotechnology, Sri. Paramakalyani College

K. Nataraja Seenivasan

Biomedical Diagnostic Laboratory, Department of Microbiology, Bharathidhasan University

I. Joseph

Research scholar, Department of Advanced Zoology and Animal Biotechnology, Sri. Paramakalyani College