

Adjusting Structural Equation Modelling Of Spiritual Coping Scale: Use Of The Sattora-Bentler Method As An Alternative To Maximum Likelihood Estimation

M Khodayari, A Abadi, H Majd, M Rassouli, H Sadeghi-Bazargani

Citation

M Khodayari, A Abadi, H Majd, M Rassouli, H Sadeghi-Bazargani. *Adjusting Structural Equation Modelling Of Spiritual Coping Scale: Use Of The Sattora-Bentler Method As An Alternative To Maximum Likelihood Estimation*. The Internet Journal of Epidemiology. 2012 Volume 10 Number 2.

Abstract

Background: Structural equation modelling (SEM) is a multivariate analysis method used to investigate direct and indirect effects among several observed or latent variables. In psychology and social sciences, data are often collected through questionnaires or inventories that commonly include Likert scale questions. Multivariate normal distribution is an essential assumption that often does not hold for this kind of data. Through an experiment on spiritual coping, our study aimed to illustrate key problems associated with using the common maximum likelihood (ML) method, and to assess a way for dealing with structural equation models when variables are in categorical form and don't exhibit a normal and continuous distribution.

Methods: Data regarding measurement of spiritual coping and its predictors collected through a questionnaire were analysed. A structural model was developed and specified for spiritual coping. The model was fitted and investigated to assess common fit indices and Sattora-bentler estimators for small samples and non-normal Likert scale data. Data analysis and modelling was done using the EQS statistical software package.

Results: It was found that fit indices and parameters encountered underestimation problems when using common ML estimator method. The robust SB- χ^2 method showed the model to have a better fit to the data ((S-B χ^2 / df) = 1.82, CFI = 0.94)).

Conclusion: Structural equation modelling using a robust SB-estimator is an appropriate method for analysing complex Likert scale measurements, especially with a small sample size, specifically regarding the spiritual coping scale and similar metrics.

INTRODUCTION

Nearly everyone working in research fields of psychology and neuropsychiatric diseases is acquainted with the Likert-type scale. The Likert scale was invented by a psychologist called Likert Rensis¹. Although general applicability of Likert scales has been often questioned, this type of measurement remains an extremely popular methodology in the fields of psychology, public health and nursing research²⁻⁶. In fact, it is so widely used in scaling responses in surveys that it is sometimes used interchangeably with rating scales, even though the two are not synonymous. Analysis of data measured on Likert scales is another issue requiring special notation. A variable measured using Likert type questions exists on an ordinal scale- generally one limited to a few levels. This poses the question of whether we can safely apply statistical methods that rely on assumptions of normality. The issue gets even more

problematic when the sample size is small and a complex model needs to be developed to assess a latent variable representative of a health-related phenomenon.

Structural equation modelling (SEM) is a statistical methodology that takes a confirmatory (i.e. hypothesis-testing) approach to the analysis of a structural theory bearing on some phenomenon. Typically, this theory represents "causal" processes that generate observations on multiple variables⁷.

Review of SEM applications during the past 15 years (in psychological research, at least) reveals most measurements to be based on Likert scaled data with estimation of parameters done using maximum likelihood (ML) procedures⁸.

When the number of categories is large and the data approximate a normal distribution, failure to address the

ordinal form of the data is likely to have negligible consequences; however, this may not be the case in many studies⁹. In psychology and other social sciences, data are often collected through questionnaires which use a Likert scale. Multivariate normality is an essential assumption that may not hold for this kind of data.

In an experiment on assessing spiritual coping, the aim of our study was to illustrate the problems of using the common maximum likelihood (ML) method, and to assess a way for dealing with structural equation models when variables are in categorical form and don't follow a normal and continuous distribution.

MATERIAL AND METHODS

The study population where the data come from contained 120 adolescents in State Welfare Organizations of Tehran database. Data was derived from institutionalized orphan adolescents between 14-20 years of age enrolled in nineteen protector centres of Tehran.

A structural model was developed and specified for spiritual coping. The developed scale was called "Institutionalized adolescents spiritual coping scale". The structural statistical model was fitted and investigated using common fit indices of ML, and Sattora-bentler estimators for small samples and non-normal Likert scale data.

SATTORA-BENTLER METHOD ()

ML methods produce parameter estimators to ensure that observed sample probability is maximized. This method assumes that observed variables have multi-normal distribution. The likelihood function will be:

Figure 1

$$\log L = -c - \frac{n}{2} \log |\Sigma| - \frac{1}{2} A$$

In which:

Figure 2

$$c = \frac{n}{2} \log(2\pi)$$

&

$$A = n \text{trace} [\Sigma^{-1} (S + (\bar{y} - \mu)(\bar{y} - \mu)')]$$

Quantity of A cause estimator function minimized.

Assuming that π shows parameters, we have:

Figure 3

$$F_{ML}(\pi) = \frac{1}{2} [\ln |\Sigma| + \text{trace}(\Sigma^{-1} T) - \ln |S| - p]$$

In which:

Figure 4

$$T = S + (\bar{y} - \mu)(\bar{y} - \mu)'$$

When mean structure was unknown, assume $\mu = 0$ and then we have:

Figure 5

$$F_{ML}(\pi) = \frac{1}{2} [\ln |\Sigma| + \text{trace}(\Sigma^{-1} S) - \ln |S| - p]$$

An alternative hypothesis under this offered model is $\mu = y$ and $\Sigma = S$ which Σ get from ML function:

Figure 6

$$\log L_{H_1} = -c - \frac{n}{2} \log |S| - \frac{n}{2} p$$

Regarding that:

Figure 7

$$F_{ML}(\pi) = -\frac{\log L}{n} + \frac{\log L_{H_1}}{n}$$

Assuming that π showed the ML estimator value given the null hypothesis, the chi square likelihood proportion for

fitting model H_0 against H_1 equals $2nF_{ML}(\pi)$.

Bentler (2005) noted that adjusting test statistics for small samples may help to achieve better estimators when analysing Likert data for the purpose of hypothesis testing.

When observed variables that do not have multivariate normality, SB- χ^2 goodness of fit test benefits from the following adjustment to chi square:

Figure 8

$$T_M = 2n F(\hat{\pi}) / c$$

In which c is adjusting constant and computed from:

Figure 9

$$c = tr[UT] / d$$

In which d is degree of freedom for model and:

Figure 10

$$U = (W^{-1} - W^{-1}\Delta(\Delta'W^{-1}\Delta)^{-1}\Delta'W^{-1})$$

In which W is a weighted matrix and its optimum value is S , also:

Figure 11

$$\Delta = \frac{\partial \Sigma}{\partial \pi} \Big|_{\hat{\pi}}$$

This study was approved as a thesis research project for a degree of master in biostatistics registered in Tarbiat Modarres University in Tehran, Iran.

RESULTS

The model for spiritual coping, offered by researchers, contained seven variables. They were spiritual coping, spiritual attitude and subjective norm, which were taken as latent variables; and hopefulness, self-respect, self-efficacy and age, which were taken as observational variables. Observational variable scores were computed from the mean of their items. Because of being three latent variables in the structural model, there were three observational models. After reducing nonrelated items, spiritual attitude, subjective norms and spiritual coping were computed by 11, 10 and 16 items by respectively. Exploratory factor analyses had given the best SEM model. Figure 1 shows the structural model, which describes spiritual coping. As shown in Fig. 1, the measurement model related to subjective norms used second order factor analyses.

Figure 12

Figure 1: Structural model for spiritual coping




Table 1 shows the results of confirmatory factor analyses for measurement models. Results show that fit indices and parameters encountered the under-estimation problem by using the common ML estimator method. Indices show that application of a robust SB-estimator produced a better fit.

Figure 13


Table 1: Confirmatory factor analysis fit indices for the measurement models

Indices	Continues			categorical			
	spiritual coping	spiritual attitude	subjective norm	spiritual coping	spiritual attitude	subjective norm	
mardia	122	21	6.5	mardia	122	21	6.5
X ²	313	102	98	SB-X ²	210	95	53
(df)	(104)	(44)	(35)	(df)	(104)	(44)	(35)
NFI	0.73	0.775	0.67	NFI	0.88	0.845	0.83
CFI	0.80	0.84	0.74	CFI	0.93	0.91	0.92
GFI	0.77	0.86	0.86	GFI	0.93	0.91	0.93
RMSEA	0.13	0.13	0.13	RMSEA	0.09	0.09	0.075
AIC	105	32	34	AIC	2	7	10

The Mardia index shows that none of the variables had normal distribution. Results show that when variables aren't normal, it is better to use Robust SB-xi square method. Figure 2 shows the measurement model of subjective norms drawn by EQS software package.

Figure 14

Figure 2: second order CFA model for measuring subjective norm latent variable



Measurement models evaluation show the efficiency of Robust Sattora-Bentler correction. Fit indices for total SEM model confirm the same results. Table 2 compares SEM model fit indices in both continuous and categorical status. Study shows that fit indices and parameters encounter the under-estimation problem by using the common ML estimator. Using a Robust SB-chi square method produces a model which better fits data of this type.

Figure 15

Table 2: Fit indexes for Structural Equation Modeling of spiritual coping

Indexes	Continues		categorical	
	Total SEM model	Indexes	Total SEM model	Indexes
X ²	1.84	SB-X ²	1.82	
NFI	0.54	NFI	0.89	
CFI	0.716	CFI	0.94	
GFI	0.660	GFI	0.84	
RMSEA	0.083	RMSEA	0.06	
AIC	1.84	AIC	1.02	

Figure 16

Figure 3 shows total the SEM model for spiritual coping, when analysed with common ML method.




Chart 3: Total SEM model for spiritual coping

DISCUSSION

An important disadvantage of all normal theory methods, including ML, is that they can yield distorted conclusions about model adequacy when there are violations of the requisite distributional assumption- here, multivariate normality. That is, when data are not normal, the test statistic Maximum Likelihood is not distributed as a chi-square variate, and the standard errors obtained are generally not correct. Assuming that each categorical variable has an underlying continuous and normal distributed scale is undoubtedly a difficult criterion to meet and, in fact, may be totally unrealistic in some cases. As such, in psychological studies, care should be taken in using SEM models on Likert scales.

This study was an example of SEM with small sample size and non-normal distribution. Many studies use Likert scale data with a small sample size in psychology, psychiatry, and social sciences. As found in our experience, in case of non-normal distribution and small sample size, fit indices get problematic and under-estimation problem is encountered. This study used the robust Sattora-bentler method for adjusting under-estimated fit indices. Results show that using this method could be beneficial for researches who need to handle categorical Likert scale data similar to that used here in spiritual coping assessment.

Crawford and Henry evaluated the Positive and Negative

Affect Schedule(PANAS) in a large sample of 1003 cases. Data had high positive skew. PANAS is measured with 20 Positive and Negative items, offered by Watson and Clark. Results produced a $\chi^2 = 689.8$ (156) using the ML method and $SB - \chi^2 = 508.3$ (156)¹⁰ using the Sattora-Bentler method.

Consistent with our results, indices garnered from another study by Crawford and Henry (2003) on the Depression Anxiety Stress Scales (DASS) demonstrated that the $SB - \chi^2$ index provided a better fit than the χ^2 index¹¹.

The Personal Disturbance Scale was evaluated by Henry (2002) with regards to anxiety and depression scores being computed from 14 Delusions-Symptoms-States items. Data was derived from 758 adolescents between the ages of 16 and 19. Based on the Kolmogorov-Smirnoff test, the normality hypothesis was rejected, and the $SB - \chi^2$ index was used by for comparing deferent models although with a quite large dataset.

Beck Depression Inventory (BDI) has also been evaluated by fitting second order factor analysis model. Results of the study showed very different values between the ML method and SB estimators. CFI index for the unadjusted ML was equal to 0.76, while it was equal to 0.93 for SB robust method¹².

In the present study, the CFA model fit indices for spiritual coping with 11 items showed evidence of moderate fitting of the model to the data. The reason was that all the variables used for measuring spiritual coping were categorical variables, yet were applied as continues variables (CFI = .836). Fitting the model to the data again using a SB robust model provided acceptable results (($SB - \chi^2$)/df) = 2.16, CFI = .908).

Consistent with previous research on other types of psychological scales, we found in our study on spiritual coping that, when variables are categorical, using the $SB - \chi^2$ robust method produces better estimates than the common ML method. The method is recommended to be employed preferentially to ML when employing both Likert scales and other similar psychological scales.

References

1. Likert R. A Technique for the Measurement of Attitudes. Archives of Psychology 1932;140:1-55.
2. Canan O, Ozcay F, Ozbay-Hosnut F, Yazici C, Bilezikci B. Value of the Likert dyspepsia scale in differentiation of functional and organic dyspepsia in children. J Pediatr

- Gastroenterol Nutr 2011 April;52(4):392-8.
3. Celenza A, Rogers IR. Comparison of visual analogue and Likert scales in evaluation of an emergency department bedside teaching programme. *Emerg Med Australas* 2011 February;23(1):68-75.
 4. Jamieson S. Likert scales: how to (ab)use them. *Med Educ* 2004 December;38(12):1217-8.
 5. Ogden J, Lo J. How meaningful are data from Likert scales? An evaluation of how ratings are made and the role of the response shift in the socially disadvantaged. *J Health Psychol* 2011 August 8.
 6. Pell G. Use and misuse of Likert scales. *Med Educ* 2005 September;39(9):970.
 7. Bentler PM. Causal modeling via structural equation systems . In: Nesselroade R, Cattell RB, editors. *Handbook of multivariate experimental psychology* . 2nd ed. New York: Plenum; 1998. p. 317-35.
 8. Breckler SJ. Applications of covariance structure modeling in psychology: cause for concern? *Psychol Bull* 1990 March;107(2):260-73.
 9. Muthen B, Kaplan D. A comparison of some methodologies for the factor analysis of non-normal Likert variables. *British Journal of Mathematical and statistical psychology* 1985;38:171-89.
 10. Crawford JR, Henry JD. The positive and negative affect schedule (PANAS): construct validity, measurement properties and normative data in a large non-clinical sample. *Br J Clin Psychol* 2004 September;43(Pt 3):245-65.
 11. Crawford JR, Henry JD. The Depression Anxiety Stress Scales (DASS): normative data and latent structure in a large non-clinical sample. *Br J Clin Psychol* 2003 June;42(Pt 2):111-31.
 12. Byrne BM, Baron P, Larsson B, Melin L. The Beck Depression Inventory: testing and cross-validating a second-order factorial structure for Swedish nonclinical adolescents. *Behav Res Ther* 1995 March;33(3):345-56.

Author Information

Mohmad Taghi Khodayari, MSc, PhD student

Department of Biostatistics, Shahid Beheshti University of Medical Sciences

Alireza Abadi, PhD

Department of Biostatistics, Shahid Beheshti University of Medical Sciences

Hamid Alavi Majd, PhD

Department of Biostatistics, Shahid Beheshti University of Medical Sciences

Maryam Rassouli, PhD

Shahid Beheshti University of Medical Sciences

Homayou Sadeghi-Bazargani, MD, MPH, PhD

Department Of Statistics And Epidemiology, Faculty Of Health And Nutrition, Tabriz University Of Medical Sciences