

Reducing Surgical Site Infection In Paediatric Neurosurgical Patients

C Huang, M Ramanan, R W Chaseling

Citation

C Huang, M Ramanan, R W Chaseling. *Reducing Surgical Site Infection In Paediatric Neurosurgical Patients*. The Internet Journal of Neurosurgery. 2013 Volume 9 Number 2.

Abstract

Background

Infection of the surgical site following insertion of external ventricular drains or ventriculoperitoneal shunts in paediatric neurosurgical patients is an important preventable complication that is associated with significant morbidity and mortality. The objective of this pilot study is to determine whether the infection rate following ventriculoperitoneal shunt and external ventricular drain insertion is reduced by the introduction of an evidence-based checklist that bundles together several simple interventions, compared to standard practice.

Methods

A checklist was developed utilizing the concept of "bundling" key evidence based components. Extensive interdepartmental consultation and literature searching was carried out in the process of developing the checklist. All relevant staff were educated in the use of the checklist. Compliance was monitored closely. During the study period from January 2008 to December 2009, the checklist was used for all patients undergoing ventriculoperitoneal shunt or external ventricular drain insertion. Patients who underwent these operations between January 2007 and December 2007 were used as controls.

Results

There were a total of 197 patients with 130 in the intervention group and 67 in the control group. The infection rate dropped from 7.5% to 2.3% with introduction of the checklist ($p=0.08$). Compliance to the checklist was 88% across the 130 patients, and the number of bed days required to treat neurosurgical site infections was reduced from 65 days to 4 days per quarter.

Conclusion

In this pilot study, we have demonstrated a nonsignificant but clinically important trend towards lower infection rates for shunt and external ventricular drain surgery with the use of a simple checklist. The proposed mechanisms for this observation include improved communication between healthcare staff and readily access to useful written information on infection reduction strategies.

INTRODUCTION

Surgical site infections (SSI) are one of the most common complications occurring in hospitalised patients (Safer Systems Saving Lives

METHOD

A review of the medical literature was performed to identify gaps in the current practice. The Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic

review, MEDLINE and EMBASE were searched to identify strategies that reduced SSI

RESULTS

There was a total of 197 patients who underwent VPS or EVD insertion during the study period (Table 2). 67 patients formed the control group prior to introduction of the checklist and 130 patients were in the intervention group after introduction of the checklist. There were a total of 8 shunt or EVD infections in the cohort of 197 patients

representing an overall rate of 4.1 %. The infection rate was 7.5% (5 out of 67) in the control group and 2.3% (3 out of 130) in the intervention group. The absolute risk reduction of 5.2% did not reach statistical significance (p=0.08) with a 95% confidence interval ranging from a 12% reduction to a 1.6 % increase. The odds ratio for developing a shunt or EVD infection following exposure to the checklist was 0.29 (95% confidence interval 0.068 to 1.265). The low number of patients and infections meant that this study was underpowered to detect a statistically significant effect.

Table 2

Summary of results

Group	Infection (%)	No infection (%)	Total
Checklist	3 (2.3)	127 (97.7)	130
No checklist	5(7.5)	62 (92.5)	67
Overall	8 (4.1)	189 (95.9)	197

chi-squared=3.016, df=1, p=0.08
odds ratio=0.29 (95% confidence interval 0.07 to 1.27)

It was also found that the number of unplanned returns to the operating theatre dropped from 34 per year in the control period to 2 per year during the intervention period. The total number of additional days of hospitalisation due to SSI decreased from 65 days to 4 days per quarter. Overall compliance with the checklist was 88%.

DISCUSSION

The

CONCLUSION

The introduction of an evidence-based checklist resulted in a decrease in the rate of SSI in paediatric patients undergoing VPS or EVD insertion surgery when compared to historical controls treated without the checklist. Due to small numbers, the benefit did not reach statistical significance. The

References

1. Australia. Melbourne, Victoria. Victoria Government Department of Human Services, Rural and Regional Health and Ages Care Services Division, Safer Systems Saving Lives – Preventing Surgical Site Infections Version 4; 2005 Dec. 17p.
2. Fry, DR, The Economic Costs of Surgical Site Infection. Surgical Infections, 2002, Volume 3, Supplement S37-43.
3. Hollenbeak CS, Murphy D, Dunagan WC, Non random selection and the attributable cost of surgical site infections.

- Infection Control Hospital Epidemiology, 2002;23:177-182
4. Kulkarni AV, Drake J, Lambert-Pasculli M, Cerebrospinal fluid shunt infection: a prospective study of risk factors. Journal of Neurosurgery 94: 195-201, 2001
5. Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ. The impact of surgical-site infections in the 1990s: attributable mortality, excess length of hospitalization, and extra costs? Infect Control Hosp Epidemiol. 1999;20:725-730.
6. Bratzler DW, Houck PM, Richards C, Steele L, Dellinger P, Fry DE, Wright C, Ma A, Carr K & Red L, Use of antimicrobial prophylaxis for major surgery: Baseline results from the national surgical infection prevention project, Arch Surg 2005, vol 140, pp.174-182.
7. Eymann R, Chehab S, Strowitzki M, Steudel WI, Kiefer M, Clinical and economic consequences of antibiotic-impregnated cerebrospinal fluid shunt catheters. J Neurosurg Pediatrics 1: 444-450, 2008
8. World Health Organization. Implementation Manual, WHO Surgical Safety Checklist 2009, Safety Surgery Saves Lives, ISBN 978 92 4 159859 0, 20p
9. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AHS, Dellinger EP, Herbosa T, Joseph S, Kibatala PL, Lapitan MCM, Merry AF, Moorthy K, Reznick RK, Taylor B, Gawande AA, A Surgical Safety Checklist to Reduce Morbidity and Mortality in a Global Population. The New England Journal of Medicine, 2009, 360;5, 491-499
10. Neily J, Mills PD, Young-Xu Y, Carney BT, West P, Berger DH, Mazzia LM, Paull DE, Bagian JP. Associated between Implementation of a Medical Team Training Program and Surgical Mortality. JAMA, 2010; 304(15): 1693-1700
11. Pronovost PJ, Goeschel CA, Colantuoni E, Watson S, Lubomski LH, Berenholtz SM, Thompson DA, Sinopoli D, Cosgrove S, Sexton JB, Marsteller JA, Hyzy RC, Welsh R, Posa P, Schumacher K, Needham D. Sustaining reductions in catheter related bloodstream infections in Michigan intensive care units: observational study, BMJ, 2010;340:c309
12. Lucey MA, Myburgh JA, Antibiotic Prophylaxis for External Ventricular Drains in Neurosurgical Patient: An Audit of Compliance with a Clinical Management Protocol. Critical Care and Resuscitation 2003;5: 182-185
13. Lingaard L, Regehr G, Cartmill C, Orser B, Espin S, Bohnen J, Reznick R, Baker R, Rotstein L, Doran D, Evaluation of a preoperative team briefing: a new communication routine results in improved clinical practice. BMJ Qual Saf February 8, 2011 doi: 10.1136/bmjqs.2009.032326
14. Conley DM, Singer SJ, Edmondson L, Berry WR, Gawande AA, Effective Surgical Safety Checklist

Implementation. J Am Coll Surg,
doi:10.1016/j.jamcollsurg.2011.01.052

15. Maxfield D, Grenny J, Lavandero R, Groah L, The Silent Treatment: Why Safety Tools and Checklists Aren't Enough, VitalSmarts, AORN & AACN;2011

Author Information

Christopher Huang

Neurosurgery Department The Canberra Hospital

Australia

rachliszt@hotmail.co

Mahesh Ramanan

Sydney Australia

maheshvarasharma@hotmail.com

Raymond W Chaseling, MBBS FRACS GradDipMed?

Sydney Australia

RaymonC2@chw.edu.au