

Color Coding Of Ampoule Labels In Anesthesia

K Ramaswamy

Citation

K Ramaswamy. *Color Coding Of Ampoule Labels In Anesthesia*. The Internet Journal of Anesthesiology. 2006 Volume 11 Number 2.

Abstract

To the Editor

It is well known that drug errors contribute to significant number of mishaps in anesthesia (1,2,3). This is a scanned photograph (Photograph 1) of similar looking ampoules of suxamethonium (Antigen Pharmaceuticals Ltd) and lignocaine (Hameln Pharmaceuticals Ltd), which were being used in our hospital recently. Timely reading of the label prevented the author from drawing up lignocaine instead of suxamethonium in an emergency.

Figure 1



This incident was highlighted in our hospital critical incident reporting system and resulted in the hospital procuring suxamethonium from a different manufacturer. This emphasises that the safest way to avoid these mishaps is to meticulously read the contents of an ampoule.

However, most of our anesthetic colleagues would agree that, they pick up an ampoule from anesthetic cupboard by looking at its colour and shape, and then read the label before drawing it up in a syringe. If all the ampoules were of the same colour and shape it would make identification difficult and similar looking ampoules especially in emergencies could cause errors. It is in just such circumstances, when stress and tired, that attention to details such as this can slip, even for the most conscientious anesthetist.

If the wrong drug is drawn up previously labelled syringe, a disaster could result.

Colour coding is used in anesthesia for cylinders, pipelines, flow meters, vaporizers, lines and syringe stickers because it adds to the safety. Why shouldn't labelling of ampoules also

be colour coded? It could only help to prevent drug errors.

CORRESPONDENCE TO

Dr. KK. Ramaswamy, MB BS, Diploma in Anesthesia.
Senior House Officer, SHO-Department of Anesthesia
Northampton General Hospital, U.K. Email:
kramaswamy@doctors.org.uk Phone: +44 01604 545671
Fax: +44 01604 545670

References

1. Bowdle, TA: Drug Administration Errors from the ASA Closed Claims Project. ASA Newsletter 2003; 67(6): 11-13.
2. Evidence-based strategies for preventing drug administration errors during anaesthesia, Anaesthesia May 2004; 59 (5): 493.
3. The Australian Incident Monitoring Study. The "wrong drug" problem in anaesthesia: an analysis of 2000 incident reports. Currie M, Mackay P, Morgan C, Runciman WB, Russell WJ, Sellen A, Webb RK, Williamson JA. Anaesth Intensive Care. 1993 Oct; 21(5):596-601.

Author Information

K.K. Ramaswamy, MBBS, Diploma in Anesthesia.

Senior House Officer, SHO-Department of Anesthesia, Northampton General Hospital