

Relationship between capillaroscopic alterations and bone ultrasound parameters in patients with Raynaud phenomenon

A Lurati

Citation

A Lurati. *Relationship between capillaroscopic alterations and bone ultrasound parameters in patients with Raynaud phenomenon*. The Internet Journal of Rheumatology. 2007 Volume 4 Number 2.

Abstract

The aim of this study was to evaluate phalangeal bone quantitative ultrasound (QUS) parameters in patients with Raynaud phenomenon (RP) and relate it with nailfold capillaroscopic findings. patients referring to our Rheumatology Unit with RP were enrolled and studied for capillaroscopic alterations; bone quality profile was measured by QUS of the phalanges: AD-SoS (Amplitude Dependent Speed of Sound) UBPI (Ultrasound Bone Profile Index), UBI (ultrasound Bone Index), Z score and T score were collected. One hundred thirty six females with RP had investigated for age, height, weight, Body Mass Index, previous diseases and therapies, menopausal age were enrolled. Nailfold capillaroscopy revealed minor alterations (borderline capillary dilatation, no capillary loss) in 36.8% (Group I), major alterations (capillaries definitely dilated, avascular areas, microbleeding) in 37.5% (Group II) and no significative alterations in 25.7% of patients (Group 0). A higher frequency of low QUS parameters in phalanges was observed in group II when compared to group I or 0 (72.5% vs. 54% vs 18%; $p < 0.01$). With an ANOVA analysis we found a significant difference between the three groups in terms of Ad-SOS (Group II 1750 ± 140 ; Group I 1890 ± 132 ; Group 0 1990 ± 167 , $p < 0.001$), UBPI (Group II 0.21 ± 0.17 ; Group I 0.36 ± 0.21 ; Group 0 0.51 ± 0.24 , $p < 0.001$), UBI (Group II 1.2 ± 0.43 ; Group I 1.4 ± 0.32 ; Group 0 1.5 ± 0.41 , $p < 0.001$), Z-scores (Group II -2.8 ± 1.45 ; Group I -1.85 ± 1.27 ; Group 0 -1.1 ± 1.39 , $p < 0.001$) and T-scores (Group II -4.8 ± 2.1 ; Group I -3.2 ± 1.8 ; Group 0 -1.8 ± 2.4 , $p < 0.001$). A standard linear regression analysis revealed an association between the capillaroscopic findings and QUS ($R 0.47 \pm 0.8$, $p < 0.01$). In our study patients with capillaroscopic alterations showed reduced phalangeal quantitative ultrasound parameters, more markedly in patients with scleroderma pattern or other major capillaroscopic alterations, independently from confounding variables.

INTRODUCTION

Raynaud's phenomenon (RP) is a vasospastic disorder characterized by episodic color changes of blanching, cyanosis, and hyperemia in response to cold and/or emotional stress. RP may be a primary or a secondary process, primarily connective tissue diseases (CTDs) and scleroderma.

Nailfold videocapillaroscopy can describe and quantify the morphological characteristics of capillaries and distinguish different forms of connective tissue disease. Furthermore, low bone mass has been seen to occur in a significant percentage of patients with scleroderma or CTDs but few data are available up to date on the usefulness of quantitative ultrasound (QUS) in the study of phalangeal bone quality and structure in primary and secondary RP. Aim of our study was to relate nailfold capillaroscopic findings to QUS phalangeal parameters.

MATERIALS AND METHODS

SUBJECTS

We studied females with RP, referring to our Rheumatologic division, at the Fornaroli Hospital of Magenta (Italy), from September 2006 to April 2007. A detailed history was taken of each patient, with particular reference to age, RP onset age, menopausal status, disease duration, current or previous treatments, and current or previous diseases; their height and weight were measured and related by the body mass index ratio (BMI). The following serological markers were determined: ESR, CRP, antinuclear antibodies, anticentromere antibodies, anti-extractable nuclear antigen, including anti-Scl70, Sm, RNP, SSB, SSA, and Jo1. Subjects were excluded if they had any history of a medical disorder known to affect bone metabolism or were on any medication that could affect bone mass.

METHODS

We measured bone ultrasound parameters at phalanges: the amplitude-dependent speed of sound (AD-SoS), broadband ultrasound attenuation (BUA), UBPI (Ultrasound Bone Profile Index), UBI (ultrasound Bone Index), fast wave amplitude (FWA), which is the maximum amplitude of the fastest peak of the received US signal; Z score and T score values (referred to healthy Italian population) were also collected. Furthermore we performed a nailfold capillaroscopy on four fingers of both hands; all examinations were performed by the same experienced rheumatologist (MGM).

STATISTICAL ANALYSIS

All values were expressed as mean \pm SD. An analysis of variance, simple (ANOVA) or with adjustment for some covariates (ANCOVA), was applied to compare the parameters measured between groups. Linear regression analysis and stepwise multiple regression analysis were used to assess the relationships between variables collected. A p value lower than 0.05 was considered statistically significant. All statistical analyses were carried out with SPSS software.

RESULTS

One hundred thirty six females with RP were enrolled. In whole population mean (\pm SD) age was 53.3 years (\pm 14.1), mean BMI was 24.8(\pm 7.8), mean menopausal age was 46.9 (\pm 5.4). The mean QUS parameters values were: AdSOS 1881 (\pm 159), FWA 28.7 (\pm 9.8), UBI 1.32 (\pm 0.38), UBPI 0.34 (\pm 0.22). The mean Z score and T score were -1.98 and -3.4 respectively. Nailfold capillaroscopy revealed minor alterations (defined as: some tortuosity, borderline capillary dilatation, no capillary loss) in 36.8% (Group I), major alterations (defined as: capillaries definitely dilated, avascular areas, megacapillaries, microbleeding and capillary thrombosis, angiogenesis) in 37.5% (Group II) and no significant alterations in 25.7% of patients (Group 0). No differences were present in age, menopausal age, BMI between groups ($p > 0.2$). In group II we found: Ad-SOS 1750 ± 140 , UBPI 0.21 ± 0.17 , UBI 12 ± 0.43 , Z-score -2.8 ± 1.45 and T-score -4.8 ± 2.1 ; with a standard ANOVA with post hoc test all these values were significantly lower than group I and group 0, also correcting for all confounding variables collected. ($p < 0.01$) (Figure 1-3).

With a linear logistic test we found a Pearson R correlation index of: 0.37 ± 0.7 for Z-score, 0.41 ± 0.5 for UBPI and

0.47 ± 0.8 for AdSOS ($p < 0.001$).

Figure 1

Figure 1: Ad-SOS mean values related to capillaroscopic pattern (p




Figure 2

Figure 2: UBPI mean values related to capillaroscopic pattern (p





Figure 3

Figure 3: Z score mean values related to capillaroscopic pattern (p)



DISCUSSION

Raynaud's phenomenon (RP) is a common clinical condition in rheumatology, with microvascular involvement as key feature. LeRoy and Medsger suggested criteria for primary RP: symmetric attacks, the absence of tissue necrosis, ulceration or gangrene, the absence of a secondary cause, negative antinuclear antibodies, normal nailfold capillaroscopy and a normal erythrocyte sedimentation rate. Secondary RP is characterized by an age of onset of more than 30 years, painful and asymmetric attacks, ischemic skin lesions, positive autoantibodies, capillaroscopic major abnormalities and/or clinical features suggestive of CTDs. Among the CTDs, scleroderma has the highest frequency of RP. Peripheral microangiopathy may be recognized by nailfold capillaroscopy, a non-invasive and safe technique that is reported to have both diagnostic and prognostic value. The peripheral microvascular damage in scleroderma is characterised by increasing structural alterations of the capillaries (giant capillaries and microhaemorrhages) with a progressive decrease in their density (scleroderma pattern). Generalised radiological osteopenia has been seen to occur in a significant percentage of patients with connective diffuse diseases and scleroderma. Bone mineral content was found to be reduced at the radius, lumbar spine, and the total body. No data are available up to date on QUS evaluation of phalangeal bone in patients with primary or secondary RP^{7,8,9}. Recently there has been a growing scientific and clinical

interest in QUS as an alternative method to dual-energy X-ray absorptiometry (DXA) for assessing skeletal status. QUS is thought to reflect not only bone density, but also some structural properties of bone, such as elasticity and trabecular stiffness and connectivity, which could be closely connected to bone strength^{10,11}.

CONCLUSION

Data collected in our study suggest that RP, primarily in patients with scleroderma or major alterations pattern suggestive for a secondary RP, modify phalangeal bone mass, bone density, and bone quality with a good correlation between nailfold patterns and QUS, independently from confounding variables.

References

1. Wigley FM. Raynaud's phenomenon. *N Engl J Med* 2002; 347: 1001-8.
2. Maricq HR Harper FE, Khan MM, Tan EM, LeRoy EC Microvascular abnormalities as possible predictors of disease subset in Raynaud's phenomenon and early connective tissue disease *Clin Exp Rheumatol* 1983;1:195-205
3. Kabasakal Y, Elvins DM, Ring EFJ, McHugh NJ Quantitative nailfold capillaroscopy findings in a population with connective tissue disease and in normal healthy controls. *Ann Rheum Dis* 1996;55:507-12 .
4. Le Roy EC, Black CM, Fleischmajer R, Jablonska S. Scleroderma (systemic sclerosis); classification, subsets and pathogenesis. *J Rheumatol* 1988;15:202-5
5. Grassi W, Medico PD, Izzo F, Cervini C. Microvascular involvement in systemic sclerosis: capillaroscopic findings. *Semin Arthritis Rheum* 2001;30:397-402
6. Bollinger J, Fagrell B. *Clinical capillaroscopy, a guide to its use in clinical research and practice*. Hogrefe & Huber Publishers. Toronto 1990
7. Borges CT, Takayama L, Aldrighi JM, Pereira RM, Souza RB. Systemic sclerosis and bone loss: the role of the disease and body composition. *Scand J Rheumatol*. 2006 Sep-Oct;35(5):384-7
8. ampaio-Barros PD, Costa-Paiva L, Filardi S, Sachetto Z, Samara AM, Marques-Neto JF Prognostic factors of low bone mineral density in systemic sclerosis. *Clin Exp rheumatol* 2005 Mar-Apr;23(2):180-4
9. Frediani B, Baldi F, Falsetti P, Acciai C, Filippou G, Spreafico A, et al. Bone mineral density in patients with systemic sclerosis *Ann Rheum Dis* 2004;63:326-327.
10. Barkmann R, Lusse S, Stampa B, Sakata S, Heller M, Gluer CC. Assessment of the geometry of human finger phalanges using quantitative ultrasound in vivo. *Osteoporos Int* 11:745-755
11. Wuster C, Albanese C, De Aloysio D, Duboeuf F, Gambacciani M, Gonnelli S, et al Phalangeal osteosonogrammetry study: age-related changes, diagnostic sensitivity, and discrimination power. The Phalangeal Osteosonogrammetry Study Group. *J Bone Miner Res* 15:1603-1614

Author Information

Alfredomaria Lurati, MD

Ospedale Fornaroli, Rheumatology Unit, Via Donatore di Sangue