

Virulence factors of Campylobacter

S Bhavsar, B Kapadnis

Citation

S Bhavsar, B Kapadnis. *Virulence factors of Campylobacter*. The Internet Journal of Microbiology. 2006 Volume 3 Number 2.

Abstract

Campylobacter is the most commonly reported bacterial cause of food-borne infection in the United States. Guillian-Barré syndrome and reactive arthritis are the associated complications with Campylobacter infection. Besides, an increasing proportion of human infections caused by *C. jejuni* are resistant to antimicrobial therapy. Cross contamination in raw poultry and consumption of undercooked poultry are the major risk factors for human campylobacteriosis. Various virulence related mechanisms viz., motility, oxidative stress defense, toxin production, invasive properties, iron acquisition and viable but non-culturable stage shield them in the human body. Since they are actively involved in the food chain, efforts to prevent Campylobacter human illness are needed.

INTRODUCTION

Campylobacter are Gram negative, slim, motile, spirally curved rods. The width and length of the cells varies from 0.2-0.9 μm and 0.5-5 μm respectively. The motility of *C.jejuni* is mediated by single polar flagellum. They are microaerophilic, do not metabolize sugars and use intermediates of the TCA cycle as source of energy. It needs 5-7% oxygen and 10% carbon dioxide with optimal growth temperature 37-42°C. [1] *C.jejuni* is recognized world wide as an important food borne pathogen. Human gastroenteritis caused by the organism is often associated with consumption of red meat and poultry. [2] The natural habitat for Campylobacter spp. is the intestine of birds and warm-blooded animals including sea gulls and other wild animals. Campylobacter may enter the environment through the feces of animals, birds or infected humans. [3]

REVIEW OF THE VIRULENCE FACTORS IN

Animals hosts of Campylobacter spp. have been indicated as a source of contamination and are associated with survival of the organism in nature, since fecal material is shed directly into aqueous environment. [4] This microorganism causes gastroenteritis, further leading to serious neurological disease like Guillain-Barre Syndrome and eventually death. [5] Various virulence factors in campylobacters contribute to survival and establishment of disease in the host.

MOTILITY AND CHEMOTAXIS

Motility and chemotaxis have been shown to play an

important role in bacterial colonization of several environments. [6] Chemotaxis towards urine has been demonstrated in *Escherichia coli*, a common urinary tract pathogen, suggesting that bacterial chemotaxis is involved in the pathogenesis of urinary tract infections. [7]

Flagellum has been suggested to play a vital role in adherence to epithelial cells. Campylobacter *jejuni* binds to the epithelial cells with the help of flagellin as an adhesin and colonizes the intestinal tracts of rabbits, suckling mice, hamsters etc. The genes coding for flagellin have been isolated from *C.jejuni* strains. [8] Fla A gene typing studies are done universally to identify the strains of *C.jejuni* and *C.coli* isolated from the environment. [9]

Bacterial chemotaxis is a complex signal transduction system by which bacteria are able to sense environmental stimuli and respond to them by flagellar rotation. In 1986, Paster and Gibbons studied the role of chemotaxis in oral colonization in selected Campylobacter and *Wolinella* species isolated from subgingival plaque, by quantitative method. Studies revealed that cells of *C.concisus* were attracted towards formate and to no other compounds tested (lactate, amino acids and sugars). Positive chemotaxis towards formate may enable Campylobacter to locate the oral cavity and further colonize plaque laden periodontal pockets. [6] Chemotaxis of Campylobacter was studied towards chemicals like, bile and mucin, which are secreted by gall bladder and the intestinal tract (primary sites of

Campylobacter infection in hosts). Other chemicals experimented with were carbohydrates, inorganic ions, organic acids and amino acids. These chemicals were qualitatively tested by hard agar plug method. Dense zones of bacterial accumulation around the agar plugs containing the attractant were observed. Mucin was the only strong chemoattractant in case of *C.jejuni* and was verified by flagellar rotation studies. Organic acid intermediates of the tricarboxylic acid cycle were chemoattractive to *Campylobacter*, *Salmonella typhimurium* and *Pseudomonas aeruginosa* all of which possess a tricarboxylic acid cycle uptake system. Chemoattraction of *Campylobacter* towards bile, mucin and L-fucose proves its ability to colonize the intestine and gall bladder of animals. [7]

INVASION AND ADHESION

One of the most important aspects of virulences in *Campylobacter* is its nature of interaction with intestinal cell lines. [10] Penetration of the intestinal mucosa play a vital role in pathogenesis of *Campylobacter* mediated gastroenteritis. *Campylobacter* adheres to the human intestinal cell lining and then becomes internalized within the cells. The translocation of *C.jejuni* across epithelial cell barrier reflects the virulent mechanism by which the organism gets access to sub mucosal tissue causing tissue damage, inflammation and thereby gastroenteritis. The virulence factors by which *Campylobacter* adheres to epithelial cells are proteins, flagella and lipopolysaccharide. [8]

The association of *C.jejuni* and *C.coli* with Hela cells was evaluated. Studies revealed that diarrhea and fever occurred more frequently in patients infected with invasive *Campylobacter* strains than in those infected with non invasive strains. Viable counts and transmission electron microscopy done after killing of extracellular bacteria by gentamycin strongly supports the fact that associated *Campylobacter* are adherent to cell membrane and are internalized into cytoplasmic vacuoles. [10] Role of flagella in adherence of *Campylobacter* to human intestinal epithelial cells was studied. *C.jejuni* flagellar mutants were constructed and examined for the ability to invade cultured human epithelial cells and also to cross the polarized cell monolayer of human epithelial origin. Findings indicated that either motility or *flaA* gene product or both are essential for translocation of *Campylobacter* across the polarized monolayer since mutants were not able to cross the cell barrier. In case of *Campylobacter*, flagellin and adhesion both are required for irreversible binding of bacteria to cells.

[8] Graham investigated the ability of clinical isolates and one reference strain of *C. fetus* to adhere to and invade the human intestinal epithelial cell line, INT 407. During an initial 4-h infection period, all *C. fetus* strains were detected intracellularly which indicated that *C. fetus* was capable of adhering, entering, and surviving within the nonphagocytic epithelial cell line, INT 407. [11]

CELL WALL

The cross section of the outer cell membrane of *Campylobacter* is typical like that of gram-negative cells. Cell wall is three layered, with an outer lipoprotein layer, a middle lipopolysaccharide layer and an inner mucopeptide layer. On enzymatic treatment it was revealed the murein was made up of muramic acid, glucosamine, alanine, glutamic acid and diaminopimelic acid. The disaccharide tetrapeptide were prescribed as Glc NAc-MurNAc-L-ALa-D-Glu-meso-Dpm-D-Ala. Eighty percent of glucose was found in cell wall polysaccharide. [12] Lipopolysaccharides (LPS) are an abundant surface component of the outer membrane of gram-negative bacteria. The LPS consisted of three distinct regions, lipid A moiety which was anchored in the outer membrane and is the endotoxic part of the LPS molecule. Other is the core, which is attached to the lipid, and at last is the O antigen attached to the outer core. The LPS molecules of *Campylobacter* are involved in adherence and play a role in antigenic variations, as *Campylobacter* has the ability to shift the LPS antigenic composition. Surprisingly N-acetyl neuraminic acid (sialic acid) is present in the core oligosaccharide, not frequently found in prokaryotes. These sialic acid residues appeared like gangliosides in structure, when attached to -D galactosidase. This molecular mimicry is involved in the neuropathological autoimmune diseases like Guillains Barre' Syndrome and Miller-Fisher Syndrome which eventually leads to the death of the patient. [13]

PROTEINS AND ENZYMES

In *Campylobacter*, the important virulent factors include cytolethal distending toxin and hemolysin. Cells' major defense mechanisms include superoxide dismutase which breaks down superoxide molecules to hydrogen peroxide and dioxygen thereby protecting several cell components including cytoplasmic enzymes, DNA and membrane factors. [14] *Campylobacter* resides within epithelial cells lining the gut lumen as well as the granulocytes and parenchyma cells within the lamina propria. The intracellular existence provides the asaccharolytic, slow growing organism a suitable niche when the microbial competition is

either less or absent. Hydrogen peroxide which is generated during aerobic metabolism reacts with myeloperoxidase, reduced iron or products of nitric oxide synthase leading to formation of more toxic intermediates such as hypochlorous anion, hydroxyl radicals, hydroxide anions, nitrogen dioxide and peroxydinitrite. Production of catalase that inactivates hydrogen peroxide interrupts the production of these toxic species and aid persistence and survival within host cells and tissues. Catalase is required for *Campylobacter* hydrogen peroxide resistance as well as persistence in macrophages. KatA is gene coding for catalase in *C.jejuni* and mutants lacking this gene showed significantly decreased viability than the wild type strain. Catalase plays minor role in intraepithelial cell survival but a significant role in intramacrophage survival by counteracting the effects of nitric oxide synthesis as well as the respiratory burst. [15]

Flagellin proteins are well known to be involved in colonization of *Campylobacter* in the host. Another virulent protein is Cad F later on termed as *Campylobacter* adhesion after identification which too is involved in host colonization. Lipoprotein which is a component of a protein binding dependent transport system for the siderophore enterochelin of both *C.jejuni* and *C.coli*, is coded by *ceu E* gene and this too is identified and characterized. Many other cytotoxins in *Campylobacter* are identified but cytolethal distending toxin (CDT) is the one which is fully characterized. The *cdt* genes cause cellular distention and eventually death of the cell lines. [16]

ANTIMICROBIAL SUSCEPTIBILITY

Campylobacter resistance to a number of antibiotics, such as tetracycline, erythromycin, ciprofloxacin, kanamycin, nalidixic acid and chloramphenicol has been reported. [17] The increasing rate of human infections caused by antimicrobial resistant strains of *C.jejuni* makes clinical management of campylobacteriosis cases more difficult. Human and healthy chicken isolates of *C.jejuni* were investigated for tetracycline resistance. High-level resistance was observed ranging from 32-256 mg/lit. Plasmids were detected in 74% isolates (30-40kb) and all of them carried the tet (O) gene. In only 6 strains resistance was found to be plasmid coded by conjugation experiments. [18] Majority of clinical isolates were found to carry foreign *dfr* genes coding for resistant variants of dihydrofolate reductase enzymes, the target of trimethoprim. The frequent trimethoprim resistance occurrence in *C.jejuni* strains was related to the high level exposures of food animals to antimicrobial drugs which would further lead to acquisition of foreign resistance genes

in naturally transformable *C.jejuni* strains. [19] In similar work, commercial broiler chicken flocks were treated with enrofloxacin. *Campylobacter*s were isolated from individual fecal samples before, during and after treatment. Rapid increase in the resistant *Campylobacter* was observed in the fluoroquinolone treated broilers. [20] During the 10 years (1991-2001), the rate of resistance to ciprofloxacin, ampicillin and tetracycline increased significantly amongst human *Campylobacter* strains. Different resistance rates to tetracycline amongst chicken isolates suggested the development of resistance during antimicrobial treatment in food animals. [21]

C.jejuni were found to contain undescribed plasmids each one approximately 35 Kb size viz., p Tet plasmid carrying tet O gene coding for tetracycline resistance and p Vir coding for virulent proteins. Mutations in these plasmids reduced adherence and invasion to INT407 cell line suggesting that plasmids are involved in virulence of *C.jejuni* pathogens. [22] A small plasmid (PCJ01) from a poultry isolate of *C.jejuni* was identified by Luo and Zhang (2001) which was further sequenced and characterized. Results indicated that PCJ01 was a 3212 bp circular molecule with 33.5% G+C content. This plasmid consisted of four open reading frames (ORFs) which share homology with RepA and RepB proteins in *C.coli* plasmids and which in turn coded for some virulent transmembrane proteins. [23]

River water is often contaminated with detergents due to the daily human activities like personal cleansing, laundry, utensil washing, household cleaning etc. and also with trace amounts of metals via the industrial effluent discharge. [24] Kazmi and colleagues (1985) showed that not only *Campylobacter* spp. but other pathogens like *E.coli*, *Vibrio*, *Salmonella* and *Shigella* were also susceptible to cadmium (20-40mg per disk). However, a few isolates of *Acinetobacter* being resistant to higher levels of silver (1024 mg/lit) and to the metals like aluminium, bismuth, boron and molybdate. [25]

IRON ACQUISITION

The symptoms of campylobacteriosis are variable in humans, ranging from mild to watery stools often accompanied with blood. The ability of pathogenic bacteria to acquire iron in the animal host is important in establishing infection. [26] A major problem for *C.jejuni* in establishing infection is less availability of free iron in mammalian fluids. Most of the iron in the cells is in the form of heme or ferritin and iron binding glycoproteins, transferrin and

lactoferrin, which are extracellular in plasma or body fluids. [27] Iron sulphur proteins participate in electron transport, anaerobic respiration, amino acid metabolism and energy metabolism. In human immune system, *Campylobacter* are exposed to the reactive oxygen species. This is important when *Campylobacter* is internalized by the phagocytes. *C.jejuni* can use a few siderophores viz., enterochelin and ferrichrome. Other iron compounds that support the growth of *C.jejuni* are heme compounds like hemin and hemoglobin, ferric iron and ferrous iron. *Campylobacter* scavenge siderophores produced by other enteric bacteria. Ferrous iron transport is important for bacterial virulence and can be related to low oxygen tension and variable pH in stomach and intestine. [28] *C.jejuni* has many systems for siderophore-mediated uptake of iron viz., enterochelin transport system, siderophore systems encoded by *cfr A* and gene *cj0718*. Periplasmic binding protein-dependent system is another iron uptake systems present in *C.jejuni*. This includes a ferrous iron uptake system (Feo B protein) which is important in assimilation of iron under microaerobic conditions. There are 3 sets of *exbBD* genes, each one linked to *ton B* gene that codes for Ton B protein, an important energy translocation protein for iron uptake systems and *ExbBD* which acts as assembly factors for TonB operation. In *C.jejuni* this iron uptake system is controlled by a global regulator, Fur. [27]

Iron acquisition studies revealed that *C.jejuni* readily obtains iron from hemin and hemoglobin. Two strains of *C.jejuni* were tested for their ability to acquire iron from hemin-hemopexin and hemoglobin-haptoglobin. Iron acquisition spot assays of the mutants showed that in addition to being unable to utilize heme, they were all incapable of using hemoglobin, hemo-hemopexin or hemoglobin-haptoglobin as iron sources. [26] Further characterization of these mutants allowed the identification of a 70 KDa outer membrane protein, which was lacking in mutants. This was later identified as Fur A iron repressed protein and the corresponding gene (*chuA*). On the *C.jejuni* NCTC11168 genome, the *chuA* gene is followed by *chub*, *chuC* and *chuD* genes which encode the components of an ABC transporter system. *Chu B* is a cytoplasmic membrane permease, *Chu C* an ATPase and *Chu D* a periplasmic binding protein. [28] Palyada and co-workers used DNA microarrays to identify the *C. jejuni* genes that were affected by iron availability. The transcript levels of 647 genes were affected after the addition of iron to iron-starving *C. jejuni* cells. Several classes of genes specific to iron acquisition and metabolism were revealed within 15 min. Directed mutagenesis of these

genes identified by the microarray analyses allowed the characterization of the ferric enterobactin receptor, previously named CfrA. Chick colonization assays indicated that mutants defective in enterobactin-mediated iron acquisition were unable to colonize the gastrointestinal tract. This work emphasized the complex response of *C. jejuni* to iron availability and provided insight regarding the role of iron in *C. jejuni* colonization in vivo. [29]

OXIDATIVE STRESS DEFENSE

Iron and oxidative stress are interconnected because iron on reacting with oxygen generates reactive oxygen species like superoxide anions (O₂⁻), peroxide (RO₂) and hydroxyl radicals (OH). Toxic oxygen species can also be generated by burst of free radical production. These toxic species damage cell lipids, proteins and DNA by oxidation. [27]

Superoxide dismutase removes superoxide by catalyzing the dissimilation of superoxides into hydrogen peroxide and oxygen (2O₂⁻ + 2H⁺ → H₂O₂ + O₂). Further catalase or peroxidase remove the hydrogen peroxide produced. Superoxide (Sod B) dismutases are classified based on the metal co-factor required. Sod B protein is located in cytoplasm and is fully characterized. [28] *Campylobacter* converts hydrogen peroxide to oxygen and water with the help of catalase and heme cofactor. *Campylobacter* adapts to hydrogen peroxide by showing increased response of catalase. *C.jejuni* and *C.coli* express a single catalase designated Kat A which is iron repressed and Per R is the regulator. [28]

Alkyl hydroxide reductase (Ahp) is a second peroxidase found in bacteria which converts reactive oxygen hydroperoxides to corresponding alcohols. *C.jejuni* produces Ahp C homolog in the absence of which it becomes more sensitive to cumene hydroperoxide. [27] Thioredoxin linked thiol peroxidase designated as Tpx belongs to bacterial antioxidant enzymes which protects the enzymes like glutamine synthetase against inactivation by oxidative stress. *C.jejuni* contains *tpx* ortholog. In *Campylobacter*, hydrogen peroxide can also be produced in the periplasm. Catalase is solely cytoplasmic so removal of periplasmic hydrogen peroxide is done via periplasmic cytochromic peroxidases. *C.jejuni* contains two such peroxidases, *Cj0020c* and *Cj0358*. Cft ferritin is an iron binding protein present in *C.jejuni* and its absence leads to sensitivity to superoxide and peroxide stress inducers. Similarly ferredoxin Fdx A absence leads to reduced aero tolerance in *C.jejuni*. Nitric oxide generated by immune system, reacts with superoxide

to give peroxy nitrite which can lead to the formation of toxic radicals. Bacteria use flavohemoglobins or single domain hemoglobin to protect themselves from such nitrosative stress. *Campylobacter* has a single domain called *Campylobacter globin* (Cgb), mutation in which, leads to hypersensitivity of *Campylobacter* to nitrosylating agents S-nitroso-glutathione. [28]

VIABLE BUT NON-CULTURABLE FORM

The ability to enter viable but non culturable (VBNC) state has been described for several enteric pathogens, including *Salmonella enteritidis*, *Escherichia coli*, *Vibrio vulnificus*, *Vibrio cholerae* and *Campylobacter jejuni*. The VBNC state represents the response to survival stress by not being cultured on laboratory media and also being capable of retaining virulence. Therefore in this form survival can prove to be an important potential public health threat. VBNC state in *Campylobacter* was 1st reported by Rollins and Colwell, who examined the ability of *C.jejuni* strain HC, from human campylobacteriosis patient, to survive in sterile stream water microcosm system. Plate counts were compared with direct viable count and acridine orange direct count methods to determine whether non-culturable cells of *Campylobacter* retain viability. Effects of temperature and aeration were evaluated in terms of transition of the organism to VBNC state. Morphological transition of spiral cells to coccoid cells was monitored by dark phase and electron microscopy as well as density gradient centrifugation. Culturability of *Campylobacter* was tested by taking plate counts of natural stream water flask microcosms (rotary shaker as well as held stationary) and these were compared with those of broth cultures (stationary biphasic and rotary shaken). The kinetics of decline in culturability of *Campylobacter* in shaken microcosm and broth were similar. When the microcosm flasks were held stationary the rate of decline was moderate. Transition to the non culturable form was accelerated at higher incubation temperatures so at 37°C the coccoid forms of *C.jejuni* spp was studied. These forms maintain intact and asymmetric membrane structure while the cell shape and size varies. Production of an extracellular viscous material by *Campylobacter* is the adaptation made in aqueous environments. The benefits of which are related to organism being able to control oxygen, nutrient and metabolite concentrations. This viscous mat formed at the agar/broth interface remains until it is mechanically disrupted, thereby providing a microenvironment for prolonged survival. This study showed that the VBNC stage is important for understanding the epidemiology of campylobacteriosis. [30] Some authors have reported the

possibility of recovering VBNC cells of *C.jejuni* by animal passage. Cappelier and co-workers chose three human isolates of *C.jejuni* Bf, 79 and 85 for testing the VBNC state when incubated in filtered, sterilized surface water. After starvation for 30 days, VBNC cells were inoculated in the yolk sacs of embryonated eggs. Culturable cells were further classified as pathogenic, as the maintenance of the adhesion potential indicated that VBNC state of *Campylobacter* constitutes a public health concern. [31] Solid phase cytometry (SPC) in conjunction with fluorescent viability staining was investigated as a tool to detect viable but non-culturable *Campylobacter jejuni* in drinking water. SPC distinguished between low numbers of dividing and non-dividing cells of *Campylobacter*, which had the potential to monitor resuscitation of VBNC cells. [32]

DISCUSSION

The virulence related mechanisms of *Campylobacter* discussed above throws light upon the insights of the mode of infection in the human body and its survival strategies. Each virulence factor has been studied in detail and based on this, the combinatorial antibiotic therapy can be designed.

CORRESPONDENCE TO

Swati P. Bhavsar Department of Microbiology, University of Pune Ganeshkhind, Pune 411007 INDIA Tel : 91-20-25690643 Fax : 91-20-25690087 Email: swatipbhavsar@yahoo.com

References

1. Beuchat L R. Efficacy of media and methods for detecting and enumerating *Campylobacter jejuni* in refrigerated chicken meat. *Appl Environ Microbiol*, 1985; 50: 934-39.
2. Baserisalehi M, Bahador N, Kapadnis B P. Effect of heat and chemical preservatives on survival of *Campylobacter* in food products. *Res J Microbiol*, 2006; 1: 512 -16.
3. Blaser M J, Hardesty H L, Powers B, Wang W L. Survival of *Campylobacter fetus* subsp. *jejuni* in biological milieus. *J Clin Microbiol*, 1980; 27; 309-13.
4. Rollins D M, Colwell R R. Viable but non culturable stage of *Campylobacter jejuni* and its role in survival in the natural aquatic environment. *Appl Environ Microbiol*, 1986; 52: 531-38.
5. Wilma C H, Jeroen A W, Frank M R, Tjakko A. Physiological activity of *Campylobacter jejuni* far below the minimal growth temperature. *Appl Environ Microbiol*, 1998; 64: 3917-22.
6. Bruce J P, Ronald J B. Chemotactic response to formate by *Campylobacter concisus* and its potential role in gingival colonization. *Infec Immun*, 1986; 52: 378-83.
7. Hughdahl M B, John T B, Michael P D. Chemotactic behavior of *Campylobacter jejuni*. *Infec Immun*, 1988; 56: 1560-66.
8. Grant C C R, Michael E, K, Witold C JR, Lucy S T. Role of flagella in adherence, internalization and translocation of *Campylobacter jejuni* in non polarized and polarized epithelial cell cultures. *Infec Immun*, 1993; 61: 1764-71.

9. Gondo T, Sekizuka T, Manaka N, Murayama O, Millar BC, Moore J E, Matsuda M. Demonstration of the shorter flagellin (flaA) gene of urease-positive thermophilic *Campylobacter* isolated from the natural environment in Northern Ireland. *Folia Microbiol*, 2006; 51: 183-90.
10. Fauchere J L, Rosenau A, Veron M, Moyon E N, Richard S, Pfister A. Association with HeLa cells of *Campylobacter jejuni* and *Campylobacter coli* isolated from human feces. *Infec Immun*, 1986; 54: 283-87.
11. Graham L L. *Campylobacter fetus* adheres to and enters INT 407 cells. *Can J Microbiol*, 2002; 48: 995-1007.
12. Smibert R M. The Genus *Campylobacter*. *Ann Rev Microbiol*, 1978; 32: 673-709.
13. Fry B N, Shi Feng, Yuen-Yuen Chen, Diane G N, Peter J C, Victoria K. The gale gene of *Campylobacter jejuni* is involved in lipopolysaccharide synthesis and virulence. *Infec Immun*, 2000; 68: 2594-3601.
14. Pesci, E C, Daniel L C, Pickett C L. Genetic, enzymatic and pathogenic studies of the iron superoxide dismutase of *Campylobacter jejuni*. *Infec Immun*, 1994; 62: 2687-94.
15. William A D, Jaime L S, Todd M P, Lynn A J. Role of catalase in *Campylobacter jejuni* intracellular survival. *Infec Immun*, 2001; 65: 6337-45.
16. Bang D D, Neilsen E M, Scheutz F, Pedersen K, Handberg K, Madsen M. PCR detection of seven virulence and toxin genes of *Campylobacter jejuni* and *Campylobacter coli* isolates from Danish pigs and cattle and cytolethal distending toxin production of the isolates. *J Appl Microbiol*, 2003; 94: 1003-14.
17. Gaudreau C, Gilbert H. Antimicrobial Resistance of *Campylobacter jejuni* subsp. *jejuni* Strains Isolated from Humans in 1998 to 2001 in Montréal, Canada. *Antimicro Agent Chemo*, 2003; 47: 2027-29.
18. Pratt A, Korolik, V. Tetracycline resistance of Australian *Campylobacter jejuni* and *Campylobacter coli* isolates. *J Antimicro Chemo*, 2005; 55: 452-60.
19. Gibreel A, Skold O. High level resistance to trimethoprim in clinical isolates of *Campylobacter jejuni* by acquisition of foreign genes (*dfr1* and *dfr 9*) expressing drug insensitive dihydrofolate reductases. *J Antimicro Chemo*, 1998; 42: 3059-64.
20. Humphrey, T. J., Jørgensen, F., Frost, J. A., Wadda, H., Domingue, G., Elviss, N. C., Griggs D J, Piddock. J V. Prevalence and Subtypes of Ciprofloxacin-Resistant *Campylobacter* spp. in Commercial Poultry Flocks before, during, and after Treatment with Fluoroquinolones. *Antimicro Agent Chemo*, 2005; 49: 690-98.
21. Luber P, Wagner J, Hahn H, Bartelt E. Antimicrobial resistance in *Campylobacter jejuni* and *Campylobacter coli* strains isolated in 1991 and 2001-2002 from poultry and humans in Berlin, Germany. *Antimicro Agent Chemo*, 2003; 47: 3925-30.
22. Bacon D J, Alm R A, Burr D H, Hu L, Kopecko D J, Ewing C P, Trust T J and Guerry P. Involvement of a plasmid in virulence of *Campylobacter jejuni* 81-176 *Infec Immun*, 2000; 68: 4384-90.
23. Luo, N. and Q. Zhang. Molecular characterization of a cryptic plasmid from *Campylobacter jejuni*. *Plasmid*, 2001; 45: 127-33.
24. Patil R V. Corrosion of metals and alloys: Analytical studies of some natural water, effluents and soil environment in Pune region. Ph.D. Thesis, Dept of Environmental Science, University of Pune, 1977; 50-70.
25. Deshpande L M, Chopade B A. Plasmid mediated silver resistance in *Acinetobacter baumannii*. *BioMetals*, 1994; 7: 49-56.
26. Pickett C L, Auffenberg T, Pesci E C, Sheen V L, Jusuf S D. Iron acquisition and haemolysin production by *Campylobacter jejuni*. *Infec Immun*, 1992; 60: 3872-77.
27. Kelly D J. The physiology and metabolism of *Campylobacter jejuni* and *Helicobacter pylori*. *J Appl Microbiol*, 2001; 90: 16-24.
28. Vliet A H M, Ketley J M Park S F, Penn C W. The role of iron in *Campylobacter* gene regulation, metabolism and oxidative stress defense. *FEMS Microbiol Rev*, 2002; 26: 173-86.
29. Palyada K, Threadgill D, Stintzi A. Iron acquisition and regulation in *Campylobacter jejuni*. *J Bacteriol*, 2004; 186: 4714-29.
30. Rollins D M, Colwell R R. Viable but non culturable stage of *Campylobacter jejuni* and its role in survival in the natural aquatic environment. *Appl Environ Microbiol*, 1986; 52: 531-38.
31. Cappelier J M, Minet J, Magras C, Colwell R R, Federighi M. Recovery in embryonated eggs of viable but nonculturable *Campylobacter jejuni* cells and maintenance of ability to adhere to HeLa cells after resuscitation *Appl Environ Microbiol*, 1999; 65: 5154-57.
32. Cools I, D'Haese E, Uyttendaele M, Storms E, Nelis H J, Debevere J. Solid phase cytometry as a tool to detect viable but non-culturable cells of *Campylobacter jejuni*. *J Microb Meth*, 2005; 63: 107-14.

Author Information

Swati P. Bhavsar, M.Sc.

Balu P. Kapadnis, M.Sc., Ph.D.