

Impact of an external energy on *Yersinia enterocolitica* [ATCC –23715] in relation to antibiotic susceptibility and biochemical reactions: An experimental study

M Trivedi, S Patil

Citation

M Trivedi, S Patil. *Impact of an external energy on Yersinia enterocolitica [ATCC –23715] in relation to antibiotic susceptibility and biochemical reactions: An experimental study*. The Internet Journal of Alternative Medicine. 2008 Volume 6 Number 2.

Abstract

Background :

While spiritual and mental energies are known to man, their impact has never been scientifically measurable in the material world and they remain outside the domain of science. The present experiments on *Yersinia enterocolitica* [ATCC –23715], report the effects of such energy transmitted through a person, Mr. Mahendrakumar Trivedi, which has produced an impact measurable in scientifically rigorous manner.

Methods:

Yersinia enterocolitica strains in revived and lyophilized state were subjected to spiritual energy transmitted through thought intervention and/or physical touch of Mr. Trivedi to the sealed tubes containing strain and were analyzed within 10 days after incubation.

Results:

The results indicated that Mr.Trivedi's energy has changed 20 of 33 biochemical characteristics of *Yersinia enterocolitica* along with significant changes in susceptibility pattern in 15 of 32 antibiotics. The Biotype number has changed from the original control strain giving rise to 2 different biotypes in treated samples while the external energy /treatment given was the same for all treated samples suggestive of random polymorphism as analyzed through an automated machine.

Conclusions:

These results cannot be explained by current theories of science, and indicate a potency in Mr.Trivedi's energy, providing a model for science to be able to investigate the impact of spiritual energy in a rigorous manner.

In lyophilized state, biochemical and enzymatic characteristics could be altered.

INTRODUCTION

All living organisms contain large complex molecules within the cells. Some of these are proteins and enzymes including DNA / RNA. The nature of the organisms is dependent on the biochemical reactions, which can be characterized and identified.

Antibiotic resistance, regardless of antibiotic and bacteria, will occur with sufficient time and drug use. Widespread antibiotic use causes selection pressure: resistant strains survive while susceptible ones are eliminated. Antibiotic resistance is progressive, increasing from low to

intermediate to high levels [1]. Bacterial strains are also characterized by their relative resistances to a range of antibiotics. However, such a change does not take place spontaneously in a strain over a 10-day period of normal incubation without continuous exposure to any drugs.

In this paper we report the impact of spiritual energy on *Yersinia enterocolitica*, in revived and lyophilized state, with respect to its antibiotic susceptibility pattern along with biochemical properties analyzed within a period of 10 days. The said energy was transmitted through thought intervention of an individual, Mr. Mahendrakumar Trivedi who has been interacting with a large number of people as a

healer over the last decade. As responses by humans can be accounted for by the placebo effect, these experiments on lower organisms were designed in order to directly test the impact through scientific studies to rule out the placebo effect.

It is widely accepted that lyophilization is the method most commonly used to store and transport microbial cultures as change in the biochemical and enzymatic characteristics of an organism cannot be carried out in this state.

MATERIAL AND METHODS

Two strains of *Yersinia enterocolitica* [ATCC –23715] were procured from MicroBioLogics in sealed packs bearing the same ATCC number and stored according to the recommended storage protocols until needed for experiments. The study was grouped as per the following.

GROUP I

One of the two sealed packets was handed over to Hinduja Microbiology Lab and was revived by them in two separate tubes, of which one was the control. The control tube was analyzed for identification, antibiotic susceptibility and biochemical reactions as per the standard protocols of sample processing in the microbiology lab.

The second tube, (ATCC 'A') having viable bacterial culture, was handed over to Mr.Trivedi for treatment, after sealing by parafilm. It was assessed on the 5th and 10th days after treatment.

TREATMENT

Mr.Trivedi held this tube in his hand under ambient conditions for between 0.5 to 3 minutes while treating it through his thought intervention process by communicating and instructing the experimental object within the tube in order to undergo the change. The tube was returned to the lab in the sealed condition itself.

GROUP II

The second sealed packet of *Yersinia enterocolitica* [ATCC –23715] (ATCC 'B') was treated by Mr. Trivedi directly in the sealed lyophilized state, using the same treatment process as above. The sealed tubes were broken and the strains were revived and analyzed on the 10th day for identification, susceptibility testing and biochemical reactions. All tests were performed with the help of automation on the Microscan Walkaway System (Dade Behring Siemens) using NBPC-30 panels. Antimicrobial

susceptibility was determined using the Minimum Inhibitory Concentration (MIC) method as per the latest CLSI guidelines.

MICROSCAN

After inoculation and rehydration with a standardized suspension of organism and incubation at 35 °C for 16 hrs, the minimum inhibitory concentration (MIC) or the qualitative susceptibility (susceptible, intermediate or resistant) was determined by observing the lowest antimicrobial concentration showing inhibition of growth. The results of susceptibility testing were expressed in millimeters of growth inhibition with disk testing and in mcg/ml in MIC testing.

QUALITY CONTROL

The acceptability of the identification media and antimicrobial agents was checked prior to the study by ATCC control organisms, *S. Aureus* ATCC 29213, & *E. Coli* ATCC 25922.

Panel used: NBPC 30

LIST OF ANTIBIOTICS TESTED

Amikacin, Amoxicillin / k clavulanic acid, Ampicillin / Sulbactam, Ampicillin, Aztreonam, Azithromycin, Cafazolin, Cefepime, Cefotaxime, Cefotetan, Cefoxitin, Ceftazidime, Cefuroxime, Ceftriaxone, Cephalothin, Chloramphenicol, Ciprofloxacin, Gatifloxacin, Gentamicin, Imipenem, Levofloxacin, Meropenem, Moxifloxacin, Nitrofurantoin, Norfloxacin, Pip/Tazo, Piperacillin, Tetracycline, Ticarcillin / k clavulanic acid, Tobramycin and Trimethoprim / Sulfa .

FOLLOWING BIOCHEMICAL TESTS WERE PERFORMED

Glucose, Sucrose, Sorbitol, Raffinose, Rhaminose, Arabinose, Inositol, Adonitol, Melibiose, Urea, Hydrogen Sulfide, Indole, Lysine, Arginine, Ornithine, Tda, Esculin Hydrolysis, Voges – Proskauer, Citrate, Malonate, Galactosidase, Colistin, Cephalothin, Oxidase, Acetamide, Cetrimide, Nitrofurantoin, Kanamycin, Nitrate, Oxidation, P4, Tartarate And Tobramycin.

SETTING

Microbiology Laboratory of P.D. Hinduja National Hospital & Medical Research Centre, Mumbai, India – accredited by The College of American Pathologists.

RESULTS

In this paper, only the changes / variations in antibiotic susceptibility and biochemical reactions that were observed before the treatment (control) and after the treatment (treated) have been reported in a tabular form attached herewith.

The results of the antibiotic susceptibility tests, identification tests and biochemical tests can be seen in Table 1.

Details of MIC values have been presented in Table 2.

ANTIBIOTIC SUSCEPTIBILITY

From tables 1 & 2, it can be seen that changes have occurred in both the revived and the lyophilized forms after treatment.

GROUP I

For two of the fifteen antibiotics shown here, Ampicillin and Cefazolin, the MIC values which were ≤ 8 in the control strain have remained the same by the 5th Day and then further increased to 16 in the treated sample by the 10th day, changing the susceptibility from sensitive in the control sample to a intermediate strain in the treated sample by the 10th day.

For twelve of fifteen antibiotics shown here, Amikacin, Aztreonam, Cefepime, Cefotaxime, Cefotetan, Cefoxitin, Ceftazidime, Cefuroxime, Chloramphenicol, Gentamicin, Tetracycline and Tobramycin, the MIC values have also shown significant increase, changing the susceptibility from sensitive in the control sample to a resistant strain in the treated sample by the 10th day .

GROUP II

Although treated directly in the lyophilized form, a change may be noted in the susceptibility to Ampicillin, whose MIC values increased after treatment from ≤ 8 in the control to 16 in the treated sample, changing the susceptibility from Sensitive in the control to intermediate in the treated sample by Day 10.

For Cephalothin, MIC values increased after treatment from 16 in the control to >16 in the treated sample, changing the susceptibility from intermediate in the control to resistant in the treated sample by Day 10.

Figure 1

Table 1: Analysis of Antibiotic susceptibility patterns, Biotype Number, Organism Identification Name and Biochemical reactions of all samples.

Name of the bacteria & ATCC no.	Antibiotics & Biochemical Reactions	Control	GROUP - I		GROUP -II
			Treated 5 th day	Treated 10 th day	Lyophilized - Treated Directly, 10 th day
			ATCC 'A'	ATCC 'A'	ATCC 'B'
<i>Y. enterocolitica</i>					
ATCC - 23715	Amikacin	S	S	R	S
	Ampicillin	S	S	I	I
	Aztreonam	S	S	R	S
	Cefazolin	S	S	I	S
	Cefepime	S	S	R	S
	Cefotaxime	S	S	R	S
	Cefotetan	S	S	R	S
	Cefoxitin	S	S	R	S
	Ceftazidime	S	S	R	S
	Cefuroxime	S	S	R	S
	Chloramphenicol	S	S	R	S
	Cephalothin	I	I	I	R
	Gentamicin	S	S	R	S
	Tetracycline	S	S	R	S
	Tobramycin	S	S	R	S
	Biotype	7101 1002	7141 1002	7736 7766	7141 1002
	Organism Identification Name	Y. entero group	Y. entero group	Y. entero group	Y. entero group
	Indole	+	+	-	+
	Inositol	-	+	-	+
	Raffinose	-	-	+	-
	Urea	-	-	+	-
	Lysine	-	-	+	-
	Tda	-	-	+	-
	Citrate	-	-	+	-
	Colistin	-	-	+	-
	Kanamycin	-	-	+	-
	Rhamnose	-	-	+	-
	Adonitol	-	-	+	-
	Hydrogen Sulfide	-	-	+	-
	Arginine	-	-	+	-
	Esculin	-	-	+	-
	Hydrolysis				
	Malonate	-	-	+	-
	Nitrate	-	-	+	-
	Melibiose	-	-	+	-
	Voges - Proskauer	-	-	+	-
	Nitrofurantoin	-	-	+	-
	Tobramycin	-	-	+	-

Key: S = susceptible, I = intermediate, R = resistant, + = positive, - = negative

Control refers to ATCC strains that were analyzed before the treatment.

Treated refers to -ATCC 'A' on 5th & 10th day after the treatment &

- ATCC 'B' lyophilized that were treated directly

Impact of an external energy on *Yersinia enterocolitica* [ATCC –23715] in relation to antibiotic susceptibility and biochemical reactions: An experimental study

Figure 2

Table 2: Minimum Inhibitory Concentration values of antibiotics in mcg/ml.

Antibiotics	Control	GROUP - I		GROUP - II
		Treated 5 th day ATCC 'A'	Treated 10 th day ATCC 'A'	Lyophilized - Treated Directly ATCC 'B'
Amikacin	≤16	≤16	>32	≤16
Ampicillin	≤8	≤8	16	16
Aztreonam	≤8	≤8	>16	≤8
Cefazolin	≤8	≤8	16	≤8
Cefepime	≤8	≤8	>16	≤8
Cefotaxime	≤8	≤8	>32	≤8
Cefotetan	≤16	≤16	>32	≤16
Cefoxitin	≤8	≤8	>16	≤8
Cefazidime	≤8	≤8	>16	≤8
Cefuroxime	≤4	≤4	>16	≤4
Chloramphenicol	≤8	≤8	>16	≤8
Cephalothin	16	16	16	>16
Gentamicin	≤4	≤4	>8	≤4
Tetracycline	≤4	≤4	>8	≤4
Tobramycin	≤4	≤4	>8	≤4

ORGANISM IDENTIFICATION

Biotype numbers of particular organisms were arrived at after interpreting the results of the biochemical tests. The Biotype numbers then led to the particular Organism Identification. Surprisingly, the biotypes have changed in all the treated samples and did not match with the control sample.

BIOCHEMICAL REACTIONS

GROUP I

In Group I, For Indole, the reaction remained positive in the treated sample by 5th Day as in control, but later found to have changed to negative by the 10th day.

For Inositol, the reaction was negative in the control sample and had changed to positive by Day 5 in the treated sample and later on changed to negative by 10th day.

In group I, eighteen of the biochemical reactions shown here, Raffinose, Urea, Lysine, Tda, Citrate, Colistin, Kanamycin, Rhaminose, Adonitol, Hydrogen Sulfide, Arginine, Esculin Hydrolysis, Malonate, Nitrate, Melibiose, Voges – Proskauer, Nitrofurantoin and Tobramycin, were unchanged by treated Day 5 in Group I, but later on the 10th day were found to have changed to positive.

GROUP II

In one of the biochemical tests, change may be noted in the Group II sample, which was treated directly in the lyophilized form. The reaction with Inositol has changed from a negative result in the control to positive by the time it

was tested on the 10th day.

DISCUSSION

While misidentifications and errors in assessment of the above characteristics are known to have occurred in such tests, the existence of changes in so many properties and tests cannot be dismissed as a machine error. The findings of alterations in antibiotic susceptibility patterns as well as in several biochemical reactions as a result of the treatment applied are unexpected and unprecedented based on the literature so far, and currently unexplained by science. The biotypes of the organisms have also changed after treatment by Mr. Trivedi. This appears to indicate the presence of an energy to which the organisms were exposed. And most important is that the particular microbe reacted to this energy, which has never been reported yet.

In separate studies, the effect of Mr.Trivedi's energy on bacterial strains of ATCC through DNA fingerprinting has indicated DNA polymorphism ranging from 4% to 79% [2] (un published). Further, 16S rDNA tests show a change in the bacterial species. [3]. Such observations are indicative of alterations in genotype. The similarity of effect of Mr.Trivedi's unique energy on biochemical changes and susceptibility patterns of *Yersinia enterocolitica* [ATCC –23715] and other bacteria referenced above [2,3] lead us to postulate that *Yersinia enterocolitica* [ATCC –23715] has also gone through a genotype change. Thus we expect to see DNA polymorphism in these strains, which may be responsible to cause such changes in antibiotic susceptibility as well as in biochemical reactions.

The results obtained in the present experiment indicate the existence of an energy that is still not known to contemporary physical and life sciences and which can effectively alter the characteristics of the microorganisms such as the bacteria studied in this work. It is not possible to find an explanation of such observations based on the current scientific knowledge. These results may give rise to better insight into the pathways that lead to mutation and may also help to understand the possible biochemical pathways that take place in the life cycle of the particular microbe.

While science has so far remained unable to enter into questions regarding 'consciousness / Spiritualism' and ordered forces or energies which can interconnect materials on a plane of higher information, the results of the current experiments prove the existence of a direct model for

investigation into the impact of a spiritual energy in a rigorous manner.

CONCLUSIONS

1 Mr. Trivedi's energy has altered the biochemical reactions along with the antibiotic susceptibility of Yersinia enterocolitica only within the period of 10 days.

2. The changes occurred in one biochemical reaction in the lyophilized state is important as per the present knowledge and would be considered significant.

3. Lyophilization is the method most commonly used for the preservation of microorganisms, as it does not alter the Chemical and enzymatic characteristics of the microbes. On contrary, the presented results have observed the changes in few characteristics of Yersinia enterocolitica in the lyophilized state itself indicating the need for superior method for storage of particular microorganism.

4. These results cannot be explained by current theories of science, and indicate a potency in Mr.Trivedi's energy, providing a model for science to be able to investigate the impact of spiritual energy in a rigorous manner.

5. In lyophilized state, biochemical and enzymatic characteristics could be altered.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Ms. Nandini Altekar for helping us in writing the paper.

References

1. Changing Antibiotic Susceptibility Patterns at a University Hospital [South Med J 94(6): 619-620, 2001. © 2001 Southern Medical Association]
2. http://www.divinelife.us/Genetics/staphylococcus_aureus.htm
3. http://www.divinelife.us/Genetics/B_Nocardia_otitidis_Treated_A.htm

Author Information

Mahendrakumar Trivedi, B.E.

Founder, Society for Divine Life

Shrikant Patil, B.D.S.

Associate, Society for Divine Life