
Justification of use of some medicinal plants in treatment of various diseases in Khulna, Bangladesh

M Faysal

Citation

M Faysal. *Justification of use of some medicinal plants in treatment of various diseases in Khulna, Bangladesh*. The Internet Journal of Third World Medicine. 2008 Volume 7 Number 2.

Abstract

11 locally used plants were selected and significant literature review was done to find out about their therapeutic activity and chemical constituents to judge their local effects.

INTRODUCTION

Plants are the natural reservoir of many antimicrobial, anticancer agents, analgesics, anti-diarrheal as well as various therapeutic activities. Bangladeshi people have traditional medical practice as an integral part of their culture. A lot of medicinal plants are available for the treatment of various diseases. However, scientific studies have been conducted only to a limited extent with few medicinal plants. In this investigation, 11 locally used plants were selected and significant literature review was done to find out about their therapeutic activity and chemical

constituents to judge their local effects.

MATERIALS AND METHODS

A field survey was done during the month June to July in Khulna district of Bangladesh and a list of various medicinal plants and their local use was collected. Then a significant literature review was done to find out about their therapeutic activity and chemical constituents in various journals, Books and Internet. This study provides some fundamental information for researchers.

RESULT AND DISCUSSION

Figure 1

Table 1: Local use and their therapeutic activity and chemical constituents of some medicinal plants of Bangladesh

Plant (Family)	Uses in traditional medicine	Reported major constituents;
<i>Aglaia roxburghiana</i> (Meliaceae)	Dysentery, leucoderma, leprosy, fever, thirst, Tumors, vomiting ¹⁶	24, 25-epoxy-29-norcycloartan-3-ol, 29-norcycloart-23-ene-3, 25-diol, 24,25-epoxy-29-nor-24-cycloarten-3 β -ol, roxburghiline, hydroxyroxburghiline, aglaroxin-A, roxburghadiol A. 10 acid ²
<i>Amoora rohıtuka</i> (Meliaceae)	Cancer, tumors, spleen and liver disease, Rheumatism. ⁹	6b,7b-epoxyguai-4-en-3-one, 6b,7b-epoxy-4b,5-dihydroxyguaiane, 11-stigmasta-5,24(28)-dien-3 β -O- β -D-glucopyranosyl-O- α -L-rhamnopyranoside, 7-keto-octadec-cis-I 1-enoic acid. ²
<i>Buchanania lanzan</i> (Anacardiaceae)	fever, diarrhea, Itch. ²¹	Myricetin 3'-rhamnoside-3-galactoside. ³
<i>Chukrasia tabularis</i> (Meliaceae)	As an astringent and antidiarrhoeic. ¹⁶	tabulalides A-E, ⁵ tabularin ⁶ , scopoletin, melanone, ⁷ chukrasin A-E. ⁸
<i>Ficus indica</i> (Moraceae)	Relieve toothache, rheumatism, humpago, inflammations, diarrhoea, dysentery, vomiting, biliousness. ⁹	Bengaloneside, leucoanthocyanidins, leucoanthocyanin glycoside, betasitosterol glycoside, mesoinositol, friedelin, beta-sitosterol, quercetin-3-galactoside and rutin, tiglic acid ester of gamma-tarzerol, cyanidin rhamnoglycoside, ficusin and bergaptin. ⁹
<i>Lannea coromandelica</i> (Anacardiaceae)	Leprous and obstinate Ulcers, toothache, mouth, Sores, impotency. ¹¹	(2R,3S)-(+)-3',5'-dihydroxy-4',7'-dimethoxydihydroflavonol, (2R,3R)-(+)-4',5,7-trimethoxydihydroflavonol, (2R,3R)-(+)-4',7-di-O-methylidihydroquercetin, (2R,3R)-(+)-4',7-di-O-methylidihydrokaempferol and (2R,3R)-(+)-4'-O-methylidihydroquercetin [21]. Quercetin-3-arabinoside, ellagic acid, α -sitosterol, physcion, physcion anthranol B, leucocyanidin. ¹⁰
<i>Nephelium litchi</i> (Sapindaceae) Syn. <i>Litchi chinensis</i>	Neurological disorders, Smallpox, throat Infection. ⁹	Folic acid, L-ascorbic acid, cyanidin-3-glucoside, cyanidin-3-rutinoside, malvidin-3-acetylglucoside, alpha-[methylene cyclopropyl] glycine. ⁹
<i>Pongamia glabra</i> (Leguminosae)	Bleeding piles, fistulous sores, bronchitis, gonorrhoea, whooping cough, tonic. ¹¹	Karanjin, ovalitenone, pongachromene, lanceolatin, betulinic acid, caffeic esters, pongapin, glabrachromene, desmethoxykanugin, (-)-isoglabrachromene, kanugin, glabra-II, fisetin tetramethyl ether, 5-methoxy-3',4'-methyleneedioxy-2'',2''-do(7,8-6'',5'')flavone ¹² , glabone. ¹³ , pongagallone-a, pongagallone-b, isopongachromene, pongamol, kanjone, pongaglabol (Glabrachalcone) ¹⁴ , isopongaglabol and 6-methoxyisopongaglabol, 5-methoxyfurano(8,74'',5'')flavone, 5-methoxy-3',4' methylenedioxyfurano (8,7-4'',5'') flavone, ovalichromene B, cycloart-23-ene-3 β ,25-diol, friedelin, and β -sitosterol- β -D-glucoside, 31-pongaglabol, aurantiamide, acetate, pongaglabol. ¹⁵
<i>Quisqualis indica</i> (Combretaceae)	Diarrhea, fever, rickets in children, boils, ulcers, helminthiasis. ⁹	Quisqualic acid, 35 quisqualin A. ⁴
<i>Semecarpus anacardium</i> (Anacardiaceae)	Scrofulous, venereal and leprosy infections, Nervous debility. ¹⁶	Anacardic acid, semicarpol, bhilawanol, monolefin I, diolefin II, bhilawanol-A, bhilawanol-B, amentoflavone tetrahydroamentoflavone, tetrahydrobustaflavone, semecarpufilavanone, gallufilavanone anacardufilavanone, anacardoside, ³⁶ semecarpetin, nallafilavanone ¹⁷ , bjeedifilavanone ¹⁸ , semecarpufilavanone ¹⁹ , gallufilavanone ²⁰ , Otrimethyl bifilavanone A1, O-trimethyl bifilavanone A2, O-Tetramethyl bifilavanone A1.
<i>Shorea robusta</i> (Dipterocarpaceae)	Ulcers, wounds, gonorrhoea, leprosy, helminthiasis. ⁸	9,10-dihydroxystearic acid, 3,25-epoxy-1,2,3,11-tetrahydroxy-12-ursen-28-oic acid, 7 28-nor-12-ursen-3-ol, shorea phenol, 2,3,23-trihydroxy-11-methoxy-12-ursen-28-oic acid. ⁴

CONCLUSION

From the study it is clear that there is more need of research in these plants to isolate and identify the active compound which works as active drug.

ACKNOWLEDGEMENT

I would like to thank the faculty members of Pharmacy Discipline, Khulna University.

References

1. Rashid, M.A., Hasan, C.M., Choudhury, S.A.R., Begum, B. and Rahman, S. 1997. Ethnopharmacological investigation of medicinal plants of Bangladesh.
2. Daulatabad, C.D. and Jamkhandi, S.A.M. 1997. A keto fatty acid from Amoorah rohituka seed oil. *Phytochemistry* 46, 155-156.
3. Arya, R., Babu, V., Ilyas, M. and Nasim, K.T. 1992. Myricetin 3'-rhamnoside-3-galactoside from *Buchanania lanzan* (anacardiaceae). *Phytochemistry* 31, 2569-2570.
4. Dictionary of Natural Products, Chapman and Hall, 2002.
5. Nakatani, M., Abdelgaleil, S.A.M., Mona, M.G., Saad, M.M.G., Huang, R.C., Doe, M. and Iwagawa, T. 2004. Phragmalin limonoids from *Chukrasia tabularis*. *Phytochemistry* 65, 2833-2841.
6. Purushothaman, K.K., Sarada, A., Saraswathi, G. and Connolly, J.D. 1977. 5,7-dihydroxy-6,2',4',5'-tetramethoxyflavone from the leaves of *Chukrasia Tabularis*. *Phytochemistry* 16, 398-399.
7. Chatterjee, A., Banerjee, B., Ganguly, S.N. and Sircar, S.M. 1974. Triterpene and coumarins from *Chukrasia tabularis*. *Phytochemistry* 13, 2012-2013.
8. Chowdhury, R. and Islam, N. A. 2004. Hydroxylated mansumbinen-28-oic acid from *Combretum coccineum*. *Biochemical Systematics and Ecology* 32, 443-445.
9. Ghani, A. 1998. Medicinal plants of Bangladesh. 2nd ed. Dhaka; Asiatic Society of Bangladesh.
10. Subramanian, S.S. and Nair, A.G.R. 1971. Polyphenols of *Lannea coromandelica*. *Phytochemistry* 10, 1939-1940.
11. Yusuf, M., Chowdhury, J.U., Wahab, M.A. and Begum, J. 1994. Medicinal Plants of Bangladesh. 1st ed. Dhaka; Bangladesh Council of Scientific and Industrial Research (BCSIR).
12. Murari, M.S., Uttam, K.M. and Asok K.M. 1991. A chromenoflavanone and two caffeic esters from *Pongamia glabra*. *Phytochemistry* 30, 3834-3836.
13. Kanungo, P.D., Ganguly, A., Guha, A., Bhattacharyya, A. and Adityachaudhury N. 1987. Glabone, a new furanoflavone from *Pongamia glabra*. *Phytochemistry* 26, 3373-3374.
14. Arya, R., Babu, V., Ilyas, M. and Nasim, K.T. 1992. Myricetin 3'-rhamnoside-3-galactoside from *Buchanania lanzan* (anacardiaceae). *Phytochemistry* 31, 2569-2570.
15. Khanna, R.N. and Seshadri, T.R. Pongaglabrone, a new component of the seeds of *Pongamia glabra*: its constitution and synthesis. *Tetrahedron* 19, 219-225.
16. Chopra, R.N., Nayar, S.L. and Chopra, I.C. 1956. Glossary of Indian Medicinal Plants, New Delhi, CSIR.
17. Murthy, S.S.N. 1988. Semecarpetin, a biflavanone from *Semecarpus anacardium*. *Phytochemistry* 27, 3020-3022.
18. Murthy, S.S.N. 1985. Jeediflavanone - a biflavonoid from *Semecarpus anacardium*. *Phytochemistry* 24, 1065-1069.
19. Murthy, S.S.N. 1983. A biflavonoid from *Semecarpus anacardium*. *Phytochemistry* 22, 1518-1520.
20. Murthy, S.S.N. 1983. A biflavanone from *Semecarpus anacardium*. *Phytochemistry* 22, 2636-2638.
21. Kirtikar, K.R., Basu, B.D., Blatter, E., Caius, J.F. and Mahaskar, K.S. 1980. Indian medicinal plants. 2nd ed. Singh B & Singh MP publishers.
22. Haque, N., Choudhury, S.A.R., Nutan, M.T.H., Rahman, G.D.S. and Rashid, M.A. 2000. Antibacterial screening of some medicinal plants of Bangladesh. *Bangladesh Journal of Physiology and Pharmacology* 15, 52-54.

Author Information

Mehedi Faysal, B.Pharm.

Pharmacy Discipline, Life Science School, Khulna University