

A Hypothetical Index for Adiposity “Body Mass Abdominal Index”- That will predict Cardiovascular disease risk factors in Children.

P Kumar

Citation

P Kumar. *A Hypothetical Index for Adiposity “Body Mass Abdominal Index”- That will predict Cardiovascular disease risk factors in Children.*. The Internet Journal of Pediatrics and Neonatology. 2008 Volume 11 Number 1.

Abstract

Obesity in preschool children is a risk factor for cardiovascular diseases in adolescents and adults. The adiposity in preschool children is measured by using weight for length, waist-to-height index and body mass index. These ratios itself have some pros and cons and mathematical complexities. This hypothesis has coined a new term- “Body Mass Abdominal Index” which is derived by the product of two independent ratios- weight for length and waist-to-height. In another way this index is also a product of body mass index and waist circumference, which signifies that it’s a function of both lean mass and fat compartments of the body. Thus this index measures adiposity and will be related as a risk factor for cardiovascular disease in adolescents and adults.

INTRODUCTION

The prevalence of overweight in children in United States has increased over the last 20 years [1]. Rapid weight gain during infancy is a risk factor for obesity [2] and hypertension [3] in children and adults. The pediatric growth charts includes weight for age, weight for length and BMI for age [4]. Currently increase in weight gain and obesity in preschool children are measured independently either by weight for length index [5], waist-to-height ratio [6] or BMI for age [5]. Age is a major criteria for adjusting weight for height index [7]. In one of the study, weight for length was used during infancy but later at 3 years of age BMI was the criteria for measurement of obesity [5]. Some other studies state that BMI is a poor indicator for assessing adiposity in the child [8]. Another index, waist-to-height ratio relates to abdominal obesity, but recent investigations states that this ratio should also be adjusted with optimal power of height [9]. Therefore all these ratios have mathematical complexity.

THE HYPOTHESIS

This Hypothesis combines the two indices- weight for height and waist-to-height ratio. They are two independent variable ratios, so product of these ratios will be more accurate for measuring adiposity and thus will be an important cardiovascular disease risk factor in children. Interestingly the final derivation gives rise to BMI, along with waist circumference as a cofactor. Therefore it suggests that BMI

can be used in preschool children but waist circumference will be influencing the final results of this index. The new term coined for this index is “Body Mass Abdominal Index” (BMAI).

Mathematically-

$$\text{Weight} / \text{Height} \times \text{Waist Circumference} / \text{Height} = \text{Weight} / (\text{Height})^2 \times \text{Waist Circumference}$$

$$= \text{BMI} \times \text{Waist Circumference} = \text{Body Mass Abdominal Index (BMAI)}$$

Where Weight is in Kg; Waist Circumference and Height is in Meters.

For example in a hypothetical case comparison, there are 2 male children, 2 years of age with some what similar growth measurements. First child’s measurements are Weight- 14 kg, Height- 90 cm, Waist Circumference- 49 cm. Second child’s measurements are Weight- 13.5 kg, Height- 88cm, Waist Circumference- 43 cm.

$$\text{First child's BMI} = 17.3, \text{Weight/ Height} = 15.5, \text{Waist / Height} = 0.54; \text{BMAI} = 8.5$$

$$\text{Second Child's BMI} = 17.4, \text{Weight/ Height} = 15.3, \text{Waist / Height} = 0.49; \text{BMAI} = 7.5$$

The above comparisons shows that BMI, Waist / Height and

A Hypothetical Index for Adiposity “Body Mass Abdominal Index”- That will predict Cardiovascular disease risk factors in Children.

Weight/ Height values are close but BMAI value clearly separates the 2 cases. The BMI includes lean mass and fat components of the body. In this scenario, although BMI value is very close but the lean mass and fat components may be totally different. The BMAI is mostly influenced by Waist circumference which will mostly include fat compartment. This signifies measurement of adiposity is better reflected in BMAI rather than BMI or Waist / Height ratio alone.

EVALUATION OF THE HYPOTHESIS AND DISCUSSION

Recent research on rapid increase of weight for length (WFL) in first 6 months of life is associated with increased risk of obesity at 3 years of age [5]. Previous studies are limited by weight criteria alone [10]. In fact, Benn RT has proposed mathematical properties of weight for length as a measurement of adiposity [11].

Another index, waist circumference and waist-to-height ratio is directly proportional to abdominal obesity. The central fat distribution is a risk factor for metabolic syndrome and cardiovascular diseases [12]. In children, it had been studied that waist-to-height ratio is a better predictor of cardiovascular disease than Body Mass Index (BMI) [13].

The BMI is calculated as body weight (Kg) divided by square of the height (Meters). The BMI is dependent on lean body mass, fatness and age [14]. Although one of the studies shows that the total body fat and percentage body fat have moderate to high correlations to BMI [15], but some other studies shows that percentage body fat and BMI depends on the gender [8]. In addition, Schaefer et al showed in boys that obesity is predicted appropriately by BMI in upper 3rd of percent body fat percentiles, not in lower 2/3rd of sample [16]. In children with similar BMI doesn't have similar levels of adiposity. Thus BMI is a poor indicator for assessing adiposity in the child.

Cole TJ, has already shown that weight for height index should be adjusted for Age (Weight /Height^p) by determining appropriate power of height (p) [7]. The optimal value of 'p' is 2, 3 and 2 in preschool children, at 11 years of age and at puberty respectively. Therefore 'p' is variable through out the infancy and childhood. Similarly recent research on waist-to-height index (Waist /Height^p) shows that optimal power of height (p) increases from age 2 to 8 years [9]. Its <1, 1 and >1 in ages <6 years, 6 years and 6 – 13 years respectively. There is another Index; Conicity Index

which is a function of weight, height and waist circumference, but it has been shown in one of the studies that BMI is better than conicity index in predicting coronary artery disease [17].

The BMAI will take into account both BMI and Waist Circumference. That will include both lean body mass and fat components of the body. The denominator of BMAI is (Height)² which includes optimal value of 'p'; that is 2 in preschool children. This will adjust weight for height index for age. There are a lot of mathematical interpretations for calculation of percentage body fat and adiposity. Every index is complicated by complex mathematical interpretations. BMAI is a very simple index to use and all the three main body measurements – weight, height, waist circumference are included. The measurement of adiposity is included in BMAI in the form of waist-to-height ratio, therefore BMAI will be a important tool in assessing cardiovascular risk factors in children. This hypothesis could be tested in cohorts of infants and preschool children. The correlation could be established with the help of DXA scanning (Dual energy Xray Absorptiometry).

Assessing adiposity in children is difficult. Recent research projects are focusing on weight for length and waist-to-height ratios in preschool children. The assessment bears direct relationship with cardiovascular risk factors in early age. Appropriate early interventions can prevent obesity related diseases in adolescents and adults. Therefore this index-”Body Mass Abdominal Index” will be useful in measuring adiposity and will predict cardiovascular risk factors in preschool children.

References

1. Ogden CL, Troiano RP, Briefel RR et al. Prevalence of overweight among preschool children in the United States, 1971 through 1994. *Pediatrics*. 1997; 99(4)
2. Stettler N, Kumanyika SK, Katz SH, Zemel BS, Stallings VA. Rapid weight gain during infancy and obesity in young adulthood in a cohort of African Americans. *Am J Clin Nutr*. 2003; 77(6):1374-1378
3. Belfort MB, Rifas- Shiman SL, Rich-Edwards J, Kleinman KP, Gillman MW. Size at birth, infant growth, and blood pressure at three years of age. *J Hypertens*. 2000;18(7):815-831
4. National Centre for Health Statistics. 2000 CDC growth charts: United States. Available at : www.cdc.gov/nccdphp/dnpa/growthcharts/sas.htm Accessed June 19,2003
5. Taveras EM, Rifas-Shiman SL, Belfort MB, Kleinman KP, Oken E, Gillman MW. Weight status in the first 6 months of life and obesity at 3 years of age. *Pediatrics* 2009;123:1177-1183
6. Li C, Ford ES, Mokdad AH, Cook S. Recent trends in waist circumference and waist-height ratio among US

A Hypothetical Index for Adiposity "Body Mass Abdominal Index"- That will predict Cardiovascular disease risk factors in Children.

children and adolescents. *Pediatrics* 2006;118:e1390-e1398

7. Cole TJ. Weight/height compared to weight/height² for assessing adiposity in childhood: influence of age and bone age on p during puberty. *Ann Hum Biol.* 1986;13(5):433-451

8. Ellis KJ, Abrams SA, Wong WW. Monitoring childhood obesity: Assessment of the weight/height² index. *Am J Epidemiol.* 1999;150:939-946

9. Tybor DJ, Lichtenstein AH, Dallal GE, Must A. Waist-to-height ratio is correlated with height in US children and adolescents aged 2 – 18 years. *Int J Pediatr Obes.* 2008; 3(3):148-151

10. Monteiro PO, Victora CG. Rapid growth in infancy and childhood and obesity in later life: a systematic review. *Obes Rev.* 2005; 6(2):143-154

11. Benn RT. Some mathematical properties of weight-for-height indices used as measures of adiposity. *Br J Prev Soc Med.* 1971; 25(1):42-50

12. Despres JP, Lemieux I, Bergeron J, Pibarot P, Mathieu P, Larose E, et al. Abdominal obesity and the metabolic

syndrome: contribution to global cardiometabolic risk. *Arterioscler Thromb Vasc Biol.* 2008; 28:1039-49

13. Savva S. Waist circumference and waist-to-height ratio are better predictors of cardiovascular disease risk factors in children than body mass index. *Int J Obes Related Metab Disord.* 2000;24:1453-8

14. Garn SM, Leonard WR, Hawthorne VM. Three limitations of the body mass index. *Am J Clin Nutr.* 1986;44:996-7

15. Maynard LM, Wisemandle W, Roche AF, Chumlea WC, Guo SS, Siervogel RM. Childhood body composition in relation to body mass index. *Pediatrics.* 2001;107:344-350

16. Schaefer F, Georgi M, Wuhl E, Scharer K. Body mass index and percentage fat mass in healthy German school children and adolescents. *Int J Obes.* 1998;22:461-469

17. Kim KS, Owen WL, Williams D, Adams-Campbell LL. A comparison between BMI and Conicity index on predicting coronary heart disease: The Framingham heart study. *Ann Epidemiol* 2000;10:424-431

Author Information

Praveen Kumar, MD

Department of Pediatrics University of Illinois- College of Medicine Chicago, IL 60612 USA