

Tubercular pericardial effusion: a case report

I Wani, H Singh, I Alam, F Shah, J Matto, M Jan, R Sofi

Citation

I Wani, H Singh, I Alam, F Shah, J Matto, M Jan, R Sofi. *Tubercular pericardial effusion: a case report*. The Internet Journal of Third World Medicine. 2008 Volume 8 Number 1.

Abstract


A case of tubercular pericardial effusion is reported in a 38 year old male. There were no stigmata of tuberculosis anywhere in the body or the any evidence of immuno-compromised state .Computed tomography scan with polymerase chain reaction was diagnostic in this case. Patient was treated with anti tubercular therapy and responded well.

CASE HISTORY

A 37 year old patient presented with fever, night sweats and weight loss of 3 months duration. He was being treated with broad spectrum antibiotics but was not responding .No history of dyspnoea , hemoptysis was elicited by patient. There was nothing significant in past history and no family history of tuberculosis. General examination showing temperature of 38 degree Celsius , blood pressure of 120/80 mm Hg. respiratory rate of 22/min., JVP was normal .No cyanosis ,clubbing or lymphadenopathy was seen. Heart sounds were muffled. Other systemic examination was normal. Moutax test was positive of more than 15 mm.HIV profile was negative. AFB of sputum and pericardial fluid were also negative . The blood culture did not yield any growth .Polymerase chain reaction (PCR) was positive for mycobacterium tuberculosis . X –ray chest was showing prominent hilar shadows. Computed tomography scan of chest showed pericardial thickening and pericardial effusion(Figure 1)

Figure 1

Fig.1 showing pericardial thickening with pericardial effusion



Doppler echocardiogram showing pericardial effusion. The patient was put on antitubercular drugs and responded well . He is regularly attending our follow up clinics.

DISCUSSION

Tubercular pericardial effusion is a un common form of extra-pulmonary tuberculosisbeing increasingly found, especially in immuno-suppressed persons . Tuberculosis is the cause of 7% of cardiac tamponade cases .The pericardium may be involved in number of ways in tuberculosis. Route of spread is usually from mediastinal or hilar nodes or from lung and rarely as part of miliary tuberculosis. In rare cases, there may be direct spread from tuberculous pneumonia. Recurrent pericardial effusion

without any manifestations of tuberculosis can be seen occasionally in tuberculous pericarditis. A large effusion is most often found with adenitis; when the effusion is due to hematogenous seeding, little pericardial fluid is found. Approximately 10% of patients with tuberculous pericarditis develop constrictive pericarditis. Early cardiac tamponade is a good predictor of subsequent constrictive pericarditis presenting in up to 50% of patients ²

A high index of suspicion is to be maintained in every case for diagnosis. Computed tomography scans findings are fibrotic thickening and, frequently, pericardial calcification. On echocardiography, thickened pericardium and fibrinous strands are seen ³. Analysis of pericardial fluid analysis in with large effusion, without tamponade or suspected purulent pericarditis, has a very low (7%) diagnostic yield ⁴. Lymphocytes predominant and high adenosine-deaminase activity in pericardial fluid is useful in diagnosing tuberculous effusion ⁵. Polymerase chain reaction (PCR) is the gold standard diagnostic test in cases when other tests are negative ⁶. In the appropriate clinical context, marked nonhilar mediastinal lymphadenopathy on chest computed tomographic studies along with a strongly positive tuberculin skin test could be of value in the noninvasive diagnosis of pericardial effusion due to tuberculosis

Treatment of tuberculous pericarditis is by standard antituberculous chemotherapy. Corticosteroids are used where rapid improvement of symptoms is deemed desirable

reduce mortality, but it does not influence the resolution of pericardial effusion ⁷. Pericardiocentesis is often used for relief of symptoms, sometimes where complete drainage of fluid by pericardiocentesis is not possible, transcatheter intra pericardial urokinase given in early course causes fibrinolysis and provides relief. ⁸

References

1. Appukuttan S, Jawahar I, Sarin, Simon c, Grant talal Bazar A A case of tuberculous pericardial Effusion ;Age and Ageing .2003; 32: 450–452
2. Suwan PK, Potjalongsilp S. Predictors of constrictive pericarditis after tuberculous pericarditis. Br Heart J 1995;73: 187–9.
3. Liu PY, Li YH, Tsai WC et al. Usefulness of echocardiographic intrapericardial abnormalities in the diagnosis of tuberculous pericardial effusion. Am J Cardiol 2001; 87: 1133–5.
4. Merce J, Sagrista J, Permanyer G, Soler J. Should pericardial drainage be performed routinely in patients who have a large pericardial effusion without tamponade? Am J Med 1998; 105: 106–9.
5. Zayas R, Anguita M, Torres F et al. Incidence of specific etiology and role of methods for specific etiologic diagnosis of primary acute pericarditis. Am J Cardiol 1995; 75: 378–82
6. Rana S, Jones A, Simpson A. Recurrent pericardial effusion: the value of polymerase chain reaction in the diagnosis of tuberculosis. Heart 1999; 82: 246–7.
7. Hakim JG, Ternouth I, Mushangi E et al. Double blind randomised placebo controlled trial of adjunctive prednisolone in the treatment of effusive tuberculous pericarditis in HIV seropositive patients. Heart 2000; 84: 183–8.
8. Winkler B, Karnik R, Slany J. Treatment of exudative fibrinous pericarditis with intrapericardial urokinase. Lancet 1994; 344: 1541–2.

Author Information

Imtiaz Wani

Post graduate Scholar, Post graduate Department of surgery, S.M.H.S Hospital Srinagar

Harbinder Singh

Post graduate Scholar, Post graduate Department of surgery, S.M.H.S Hospital Srinagar

Inthikab Alam

Registrar, Post graduate Department of surgery, S.M.H.S Hospital Srinagar

Firdous Shah

Post graduate Scholar, Post graduate Department of surgery, S.M.H.S Hospital Srinagar

Javeed Matto

Registrar, SKIMS Medical College, Srinagar

Masooda Jan

Post graduate, Scholar Post graduate Department of surgery, S.M.H.S Hospital Srinagar

Rayees Sofi

SKIMS Medical College, Srinagar