

Investigation on animal and human rabies in cases from Mazandaran and Golestan provinces referred to the Amol Research Center, Northern Iran in 2003–2005

S Abolfazl Hashemi, B Esfandiari, A Fayaz, H Nahrevanian, R Behzadi, S Kavosian, S Payman Ziapour, S Ghobadi-rad, F Maghsoudi

Citation

S Abolfazl Hashemi, B Esfandiari, A Fayaz, H Nahrevanian, R Behzadi, S Kavosian, S Payman Ziapour, S Ghobadi-rad, F Maghsoudi. *Investigation on animal and human rabies in cases from Mazandaran and Golestan provinces referred to the Amol Research Center, Northern Iran in 2003–2005*. The Internet Journal of Microbiology. 2006 Volume 4 Number 1.

Abstract

Rabies has been endemic in Mazandaran and Golestan provinces for long time and it was a cause of some outbreaks of disease in the north of Iran especially in Golestan province. This study has been carried out in northern part of Iran for three years from 2003 to 2005. Eighty six animal bite cases were investigated and out of them, %20 was female and %80 was male. Average ages of samples were 26.5, 32 and 29.3 among female, male and in all samples respectively. According to the location of bites, 48 cases (%55.8) were in hands, 26 cases (%32.2) in feet, 11 cases (%12.8) in body and one case (%1.2) in head. Animals were involved in biting were dog (%73), mouse (%6), cat (%5), cow and jackals (%3.5), donkey (%2.5) and fox, sheep, tiger (%1.3 each). Results of immuno-fluorescent assay revealed that 36 out of 86 dogs, 0 out of 5 mice, 0 out of 4 cats, 2 out of 3 Jackals, 3 out of 3 cows, 1 out of 2 donkeys, 1 out of 1 sheep, 1 out of 1 fox and 0 out of 1 tiger were positive. In this study, data analysis indicated no human cases of rabies in the collected samples. The results confirmed a high prevalence rate of animal form of rabies in northern district of Iran, which is required an extra precaution strategy.

INTRODUCTION

Rabies is severe and fatal encephalitis caused by bite of infected animals. This is a health problem especially in the eastern Mediterranean countries (1). Infection is started by a virus from the Rhabdoviridae family, genus Lyzzaviruses and class Mononegavirales (11). The virus is sensitive strong acids and bases (14). Transmission is occurred by animal saliva by any form of skin injuries (bite, scratch, or cut) (16).

The incubation period of disease is reported to be at least 10 days and at last 5 years in Iranian cases (18). The pathogenesis of disease is commenced by entrance of viruses to the nerve cells (15) and spread out among body, however no viruses have been found in blood and stool of patients (17).

According to WHO reports, ten millions of people are bitten by animals around the world, considered for prophylaxis and treatment against rabies and almost 50,000 people die from this disease annually (1, 2). The highest rates of mortality and morbidity in Asia are observed among developing countries including Bangladesh, Pakistan and India. The real rates of

infection are also doubtful in above countries (3).

This will assist to design a new strategy for prevention in different areas of Iran. According to statistics the number of rabies infected cases is seven times higher in men than women (5). In a study in 1997 in Mazandaran province, the majority of bites were reported in feet (6). In another study between 1981 to 1985 in districts of Caspian sea and Persian Gulf, foot and hand infections were %46.2 and %33.4 respectively (8). Between 1995 to 1999 %88 of human biting was reported in male (7). According to data reported by Pasteur Institute of Iran in 2003, the number of anti-rabies treated cases were 99,861 in Iran; from the 88,466 people (%5.9) were bitten by dog (5). WHO in 1997 published an eradication report of rabies in 56 countries in the world (13). Although the number of injured and suspected person are increasing in recent years, the treatment strategy is also expanding, e. g. in 1998 the number of treated people were 65,632 (8, 9).

MATERIALS AND METHODS

Human cases were referred to Amol Research Center (ARC,

Pasteur Institute of Iran, Amol, Mazandaran, Iran(from Golestan and Mazandaran provinces. The collected data were statistically analyzed using descriptive softwares.

RESULTS

Results of this study are indicated no human cases of rabies in Gilan and Mazandaran provinces. From 86 animal bites, %20.9 was women and %79.1 was men. The average age in women was 26.5, in men 32 and in total 29.3 years old. All 48 cases (%55.8) of injuries were occurred in hands, 26 cases (%30.2) in foot, 11 cases (%12.8) in body and 1 case (%1.2) was observed in head. According to seasonal distribution, 17 cases were occurred in spring, 27 cases in summer, 18 cases in autumn and 22 cases in winter.

DISCUSSION

According to studies were carried out by the Department of Rabies, Pasteur Institute of Iran, the number of anti-rabies treated people was increased from 1990 to 2002 (4, 5). People awareness and their contribution were also increased significantly (12).

There are some reasons could explain seasonal variation rates of disease including; a) In the winter the wild animals immigrate from highlands to the human residential areas for food. b) In the warm weathers the biting increases including %75 in dogs, %6 in mice, %5 in cats, %3.5 in cow and jackals, %2.5 in donkeys and %1.3 in other animals.

Rabies as a dangerous disease is expanding due to the high number of stray dogs and human cases of dog biting (19). However, it is surprising; no cases of human rabies were reported in northern provinces of Iran during last tree years. This is because of increasing people knowledge and upgrading of health and treatment centers. The high rate of disease was observed in male, which is related to more contact and availability in natural fields. It is emphasized here the importance role of dogs for transmission of rabies in Iran. This study also clarified some epidemiological aspects of disease in the Northern part of this Middle Eastern country.

CORRESPONDENCE TO

Dr. H. Nahrevanian, Department of Parasitology, Pasteur Institute of Iran, Paster Avenue, Tehran 13164, Iran, Tel / Fax: +98-21-66968855, E-mail: mobcghn@pasteur.ac.ir

Dr. B. Esfandiari, Pasteur Institute of Iran, Amol Research Center, Amol, Mazandaran, Iran, PO Box 139, E-mail: behzadesfandiari@yahoo.com

References

1. WHO in the Eastern Mediterranean Region. Annual reports of Regional Director (1950-2000), Alexandria World Heath Organization, Regional Office for Eastern Mediterranean Region, 2000.
2. Frederik A., Murph PE. Veterinary Virology, 3th Ed., Academic Press, 1999.
3. Strategies for the control and elimination of rabies in Asia. Report of WHO Interregional Consultation: Geneva, Switzerland, 2001.
4. Simani S. Rabies in Iran. Iran. J. Vet. Fac. University of Tehran, 2003; 8: 175-178.
5. Simani S. Rabies Disease. 1th Ed., Pasteur Institute of Iran Publication, Tehran, Iran, 2004.
6. Simani S. Epidemiological study on human rabies in Iran from 1995 to 2002. Iran. J. Infect. Trop. Dis. 2002; 7(16): ?-?
7. Manshouri K. A retrospective study on Rabies in Isfahan, Iran between 1994 to 1998. DVM thesis, Shahid Bahonar University of Kerman, Iran, 2000.
8. Nowrouzian I., Fayaz A., Simani S., Ossoli M. Animal bites: a statistical analysis of cases in Persian Gulf and Caspian Sea areas, Iran. J. Vet. Fac. University of Tehran, 1988; 43:115-47.
9. Fayaz A., La Rage en Iran Rabies in tropics. Springer-Verlag, Berlin, Germany, 1985.
10. Animal Report of WHO-Collaborating Center for Reference Research on Rabies, Pasteur institute of Iran, 1998.
11. Wagner RR. and Rose JK. Rhabdooviridea: The virus and Their Replication Chapter 37, Knipe DM. and et al. Fields Virology, 3th Ed., Vol. 1, published by liffincott Raven, Philadelphia, New York.
12. Baer GM. The natural history of Rabies. Ed., Boca Raton, FL: CRC Press, 1991.
13. World Survey of Rabies No.33 for the year 1997, World Health Organization, Department of Communicable Disease Surveillance and Response, 1999, pp: 1-29.
14. Mandel G. and et al. Infectious disease. 3th Ed., 1990.
15. Platkin SA, Clark HF, Rabies virus. Feigin RD, Cherry JD, Text book of Pediatric infection disease, 4th Ed, Philadelphia, WB Saunders, 1998, P: 2111-25.
16. WHO/ Rabies: Fact sheet 99, revised June 2001.
17. Simani S. The Rabies Transmission. Abstract book of The 1th National Virology Congress of Iran, Tehran, Iran, Feb 2002.
18. Fayaz A. The diseases with vector or reservoir: Rabies. In: Epidemiology and control of prevalent diseases in Iran. Azizi F, Hatami H, Janghorbani M. (Eds.), Chapter 9 (Part 4), 2th Ed., Khosravi Publications, Tehran, Iran, 2004; pp: 542-557.
19. Simani S, Fayaz A, et al. A guideline in management of Iranian control program for Rabies, Center for Diseases Control (CDC) of Iran. Iranian Ministry of Health and Medical Education, Seda Publication, Tehran, Iran, 2004.

Author Information

Sayed Abolfazl Hashemi

Amol Research Center, Pasteur Institute of Iran

Behzad Esfandiari

Amol Research Center, Pasteur Institute of Iran

Ahmad Fayaz

Department of Rabies, Pasteur Institute of Iran

Hossein Nahrevanian

Department of Parasitology, Pasteur Institute of Iran

Ramezan Behzadi

Amol Research Center, Pasteur Institute of Iran

Saeed Kavosian

Amol Research Center, Pasteur Institute of Iran

Seyyed Payman Ziapour

Amol Research Center, Pasteur Institute of Iran

Sara Ghobadi-rad

Amol Research Center, Pasteur Institute of Iran

Farid Maghsoudi

Amol Research Center, Pasteur Institute of Iran